


# Searching for Lepton Jets

Itay Yavin

New York University




M. Baumgart, C. Cheung, J. T. Ruderman, L. T. Wang and I. Y. 0901.0283 [hep-ph]

C. Cheung, J. T. Ruderman, L. T. Wang and I. Y. 0909.0290[hep-ph]


# Part I

## Introduction





# Motivation 1



# Motivation 1






# Motivation 1




# Motivation 1



# Motivation 1



# Motivation 1






# “A Theory of Dark Matter”

Arkani-Hamed, Finkbeiner, Slyter and Weiner suggested a unified description:

# “A Theory of Dark Matter”

Arkani-Hamed, Finkbeiner, Slyter and Weiner suggested a unified description:


ATIC & PAMELA & HAZE




# “A Theory of Dark Matter”

Arkani-Hamed, Finkbeiner, Slyter and Weiner suggested a unified description:

## ATIC & PAMELA & HAZE




## INTEGRAL



# “A Theory of Dark Matter”


Arkani-Hamed, Finkbeiner, Slyter and Weiner suggested a unified description:

ATIC & PAMELA & HAZE




Lepton Jets

INTEGRAL



GGI

Maybe even DAMA




Itay Yavin


# “A Theory of Dark Matter”

Arkani-Hamed, Finkbeiner, Slyter and Weiner suggested a unified description:


ATIC & PAMELA & HAZE



INTEGRAL




Maybe even DAMA




TeV DM charged under a (non)-abelian gauge theory, weakly mixed with SM.

# Resolution of PAMELA

So dark matter annihilates to dark photons first. The dark photons then decay into leptons.



- 1) **Protons** are kinematically disallowed.
- 2) The leptons are **direct products** of the annihilations.
- 3) **Sommerfeld enhancement** of the cross-section due to light particle exchange.




# Motivation 2

Aside from the recent astrophysical observations, there can be another motivation for looking for such objects.

# Motivation 2


Aside from the recent astrophysical observations, there can be another motivation for looking for such objects.






# Motivation 2

Aside from the recent astrophysical observations, there can be another motivation for looking for such objects.




# Motivation 2

Aside from the recent astrophysical observations, there can be another motivation for looking for such objects.




# Motivation 2

Aside from the recent astrophysical observations, there can be another motivation for looking for such objects.





# Hidden Valleys

Strassler and Zurek's proposal of hidden valleys share some of the phenomenology and lepton jet searches can in principle be sensitive to these type of models as well,



\* Taken (without permission) from Strassler's talk.

# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>

Excited DM - [Finkbeiner and Weiner \(astro-ph/0702587v1\)](#)

Secluded DM - [Pospelov, Ritz, and Voloshin \(0711.4866\)](#)


A Theory of DM - [Arkani-Hamed, Finkbeiner, Slatyer, and Weiner \(0810.0713\)](#)

Lepton Jets

GGI


Itay Yavin

# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>



$$\mathcal{L}_{\text{DS}} \supset i\bar{\chi}\gamma^\mu D_\mu\chi + M\bar{\chi}\chi$$

Excited DM - [Finkbeiner and Weiner \(astro-ph/0702587v1\)](#)

Secluded DM - [Pospelov, Ritz, and Voloshin \(0711.4866\)](#)


A Theory of DM - [Arkani-Hamed, Finkbeiner, Slatyer, and Weiner \(0810.0713\)](#)

Lepton Jets

GGI


Itay Yavin

# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>



$$\mathcal{L}_{\text{DS}} \supset i\bar{\chi}\gamma^\mu D_\mu\chi + M\bar{\chi}\chi$$

$$-\frac{1}{4}f_{\mu\nu}f^{\mu\nu} + \frac{1}{2}m^2 b_\mu b^\mu$$

Excited DM - Finkbeiner and Weiner (astro-ph/0702587v1)

Secluded DM - Pospelov, Ritz, and Voloshin (0711.4866)


A Theory of DM - Arkani-Hamed, Finkbeiner, Slatyer, and Weiner (0810.0713)

Lepton Jets

GGI


Itay Yavin

# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>



$$\mathcal{L}_{\text{DS}} \supset i\bar{\chi}\gamma^\mu D_\mu\chi + M\bar{\chi}\chi$$

$$-\frac{1}{4}f_{\mu\nu}f^{\mu\nu} + \frac{1}{2}m^2 b_\mu b^\mu$$

$$+|D_\mu h_i|^2 - V(h_i)$$

Excited DM - Finkbeiner and Weiner (astro-ph/0702587v1)

Secluded DM - Pospelov, Ritz, and Voloshin (0711.4866)

A Theory of DM - Arkani-Hamed, Finkbeiner, Slatyer, and Weiner (0810.0713)


Lepton Jets

GGI

Itay Yavin




# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>



$$\mathcal{L}_{\text{DS}} \supset i\bar{\chi}\gamma^\mu D_\mu\chi + M\bar{\chi}\chi$$

$$\frac{1}{4}f_{\mu\nu}f^{\mu\nu} + \frac{1}{2}m^2 b_\mu b^\mu$$

$$M \sim \text{TeV}$$

$$+ |D_\mu h_i|^2 - V(h_i)$$

Excited DM - Finkbeiner and Weiner (astro-ph/0702587v1)

Secluded DM - Pospelov, Ritz, and Voloshin (0711.4866)


A Theory of DM - Arkani-Hamed, Finkbeiner, Slatyer, and Weiner (0810.0713)

Lepton Jets

GGI


Itay Yavin

# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>



$$\mathcal{L}_{\text{DS}} \supset i\bar{\chi}\gamma^\mu D_\mu\chi + M\bar{\chi}\chi$$

$$\frac{1}{4}f_{\mu\nu}f^{\mu\nu} + \frac{1}{2}m^2 b_\mu b^\mu$$

$$M \sim \text{TeV}$$

$$m \sim \text{GeV}$$


$$+ |D_\mu h_i|^2 - V(h_i)$$

Excited DM - Finkbeiner and Weiner (astro-ph/0702587v1)

Secluded DM - Pospelov, Ritz, and Voloshin (0711.4866)

A Theory of DM - Arkani-Hamed, Finkbeiner, Slatyer, and Weiner (0810.0713)

# A Tale of Two Sectors



$$\mathcal{L}_{\text{SM}} = \dots$$

<http://pdg.lbl.gov>

$$\mathcal{L}_{\text{DS}} \supset i\bar{\chi}\gamma^\mu D_\mu\chi + M\bar{\chi}\chi$$

$$-\frac{1}{4}f_{\mu\nu}f^{\mu\nu} + \frac{1}{2}m^2 b_\mu b^\mu$$

$$M \sim \text{TeV}$$

$$m \sim \text{GeV}$$

$$+|D_\mu h_i|^2 - V(h_i)$$


Excited DM - Finkbeiner and Weiner (astro-ph/0702587v1)

Secluded DM - Pospelov, Ritz, and Voloshin (0711.4866)

A Theory of DM - Arkani-Hamed, Finkbeiner, Slatyer, and Weiner (0810.0713)

# Limits on Kinetic Mixing

The kinetic mixing with the photon is bounded by low energy experiments, in particular the muonic g-2 ratio (Pospelov 0811:1030):



Notice that this measurement does not bound  $\epsilon_2$

# Dark Spectrum

$$\text{TeV} \frac{\quad}{M\bar{\chi}\chi}$$

# Dark Spectrum

$$\text{TeV} \quad \overline{M\bar{\chi}\chi}$$

$$\text{GeV} \quad \overline{m_b^2 b_\mu b^\mu}$$

# Dark Spectrum



TeV  $\overline{\hspace{10em}}$   $M\bar{\chi}\chi$   $\begin{matrix} \text{---} \\ \text{---} \\ \text{---} \end{matrix}$   $\sim \text{MeV}$

$yh\chi\chi + \text{h.c.}$

GeV  $\overline{\hspace{10em}}$

$m_b^2 b_\mu b^\mu$

# Dark Spectrum





# Dark Spectrum

TeV  $\overline{\hspace{1.5cm}}$   $M\bar{\chi}\chi$   $\xrightarrow{\hspace{1cm}}$   $\overline{\hspace{1.5cm}}$   $\sim \text{MeV}$

$yh\chi\chi + \text{h.c.}$


Can produce at colliders!!!

GeV  $\overline{\hspace{1.5cm}}$   $m_b^2 b_\mu b^\mu$   $\xrightarrow{\hspace{1cm}}$   $\overline{\hspace{1.5cm}}$   $\sim \text{GeV}$

$+V(h_1, h_2)$

# Part II

## Production and Evolution of Dark States



# Coupling to the Standard Model

# Coupling to the Standard Model

In general the dark gauge-boson can mix with both the photon and the  $Z^0$ ,

$$\begin{aligned}\mathcal{L}_{\text{gauge mix}} &= -\frac{1}{2}\epsilon_1 b_{\mu\nu} A^{\mu\nu} - \frac{1}{2}\epsilon_2 b_{\mu\nu} Z^{\mu\nu} \\ &= -\frac{1}{2}\epsilon'_1 b_{\mu\nu} B^{\mu\nu} - \frac{1}{2}\epsilon'_2 b_{\mu\nu} W_3^{\mu\nu}\end{aligned}$$

# Coupling to the Standard Model

In general the dark gauge-boson can mix with both the photon and the  $Z^0$ ,

$$\begin{aligned}\mathcal{L}_{\text{gauge mix}} &= -\frac{1}{2}\epsilon_1 b_{\mu\nu} A^{\mu\nu} - \frac{1}{2}\epsilon_2 b_{\mu\nu} Z^{\mu\nu} \\ &= -\frac{1}{2}\epsilon'_1 b_{\mu\nu} B^{\mu\nu} - \frac{1}{2}\epsilon'_2 b_{\mu\nu} W_3^{\mu\nu}\end{aligned}$$

If **supersymmetry** is only softly broken in the dark sector, then there is also an important mixing of the electroweak gauginos with the dark gaugino:

$$\mathcal{L}_{\text{gaugino mix}} = -2i\epsilon'_1 \tilde{b}^\dagger \bar{\sigma}^\mu \partial_\mu \tilde{B} - 2i\epsilon'_2 \tilde{b}^\dagger \bar{\sigma}^\mu \partial_\mu \tilde{W}_3 + \text{h.c.}$$

# Coupling to the Standard Model

In general the dark gauge-boson can mix with both the photon and the  $Z^0$ ,

$$\begin{aligned}\mathcal{L}_{\text{gauge mix}} &= -\frac{1}{2}\epsilon_1 b_{\mu\nu} A^{\mu\nu} - \frac{1}{2}\epsilon_2 b_{\mu\nu} Z^{\mu\nu} \\ &= -\frac{1}{2}\epsilon'_1 b_{\mu\nu} B^{\mu\nu} - \frac{1}{2}\epsilon'_2 b_{\mu\nu} W_3^{\mu\nu}\end{aligned}$$

If **supersymmetry** is only softly broken in the dark sector, then there is also an important mixing of the electroweak gauginos with the dark gaugino:


$$\mathcal{L}_{\text{gaugino mix}} = -2i\epsilon'_1 \tilde{b}^\dagger \bar{\sigma}^\mu \partial_\mu \tilde{B} - 2i\epsilon'_2 \tilde{b}^\dagger \bar{\sigma}^\mu \partial_\mu \tilde{W}_3 + \text{h.c.}$$

All in all we have the following couplings (after diagonalization and etc.), which act as a portal to the dark sector

$$\mathcal{L}_{\text{portal}} = \epsilon_1 b_\mu J_{\text{EM}}^\mu + \epsilon_2 Z_\mu J_b^\mu + \epsilon'_1 \tilde{B} \tilde{J}_{\tilde{b}} + \epsilon'_2 \tilde{W}_3 \tilde{J}_{\tilde{b}}$$


# Prompt (Dark) Photon Production

The coupling to the electromagnetic current allows for the production of the dark photon in a similar manner to a prompt photon ([Baumgart et al. 0901.0283](#)) :



# Prompt (Dark) Photon Production

The coupling to the electromagnetic current allows for the production of the dark photon in a similar manner to a prompt photon (Baumgart et al. 0901.0283) :




**Too much background!!!**




# Prompt (Dark) Photon Production

The coupling to the electromagnetic current allows for the production of the dark photon in a similar manner to a prompt photon ([Baumgart et al. 0901.0283](#)):




# Prompt (Dark) Photon Production

The coupling to the electromagnetic current allows for the production of the dark photon in a similar manner to a prompt photon ([Baumgart et al. 0901.0283](#)):




# Prompt (Dark) Photon Production

The coupling to the electromagnetic current allows for the production of the dark photon in a similar manner to a prompt photon ([Baumgart et al. 0901.0283](#)):




# Prompt (Dark) Photon Production


The coupling to the electromagnetic current allows for the production of the dark photon in a similar manner to a prompt photon ([Baumgart et al. 0901.0283](#)):




Lepton-Jet recoiling against a QCD jet:



# Cross-Sections at the LHC




# Cross-Sections at the LHC




# Rare $Z^0$ Decay

The neutral vector-boson couples directly to the dark current (Baumgart et al. and Cheung et al.). Therefore, the dark higgses and can be directly produced:



# Rare $Z^0$ Decay

The neutral vector-boson couples directly to the dark current (Baumgart et al. and Cheung et al.). Therefore, the dark higgses and can be directly produced:






# Rare $Z^0$ Decay

The neutral vector-boson couples directly to the dark current (Baumgart et al. and Cheung et al.). Therefore, the dark higgses and can be directly produced:


**Lepton Jets** - A collimated collection of energetic leptons with a small opening angle.



# Rare $Z^0$ Decays - Reach


At LEP :  $\text{BR}(Z \rightarrow f\bar{f}) = \frac{\epsilon_2^2 g_{\text{dark}}^2}{12\pi} \frac{M_{Z^0}}{\Gamma_{Z^0}} \rightarrow \mathcal{O}(100)$  events for  $\epsilon_2 = 10^{-2}$

At Tevatron and LHC :



# Neutralino Decay


The bottom of the SUSY cascade is no longer stable ([Arkani-Hamed and Weiner](#)). It will decay into the dark sector. A clean channel is electroweak-ino production ([Cheung et al.](#))



# Neutralino Decay


The bottom of the SUSY cascade is no longer stable ([Arkani-Hamed and Weiner](#)). It will decay into the dark sector. A clean channel is electroweak-ino production ([Cheung et al.](#))

$$+ \epsilon'_1 \tilde{B} \tilde{J}_{\tilde{b}} + \epsilon'_2 \tilde{W}_3 \tilde{J}_{\tilde{b}}$$




# Neutralino Decay

The bottom of the SUSY cascade is no longer stable ([Arkani-Hamed and Weiner](#)). It will decay into the dark sector. A clean channel is electroweak-ino production ([Cheung et al.](#))




# Neutralino Decay

The bottom of the SUSY cascade is no longer stable ([Arkani-Hamed and Weiner](#)). It will decay into the dark sector. A clean channel is electroweak-ino production ([Cheung et al.](#))




# Neutralino Decay

The bottom of the SUSY cascade is no longer stable ([Arkani-Hamed and Weiner](#)). It will decay into the dark sector. A clean channel is electroweak-ino production ([Cheung et al.](#))



# Neutralino Decay


The bottom of the SUSY cascade is no longer stable ([Arkani-Hamed and Weiner](#)). It will decay into the dark sector. A clean channel is electroweak-ino production ([Cheung et al.](#))







# Neutralino Decay

The bottom of the SUSY cascade is no longer stable (Arkani-Hamed and Weiner). It will decay into the dark sector. A clean channel is electroweak-ino production (Cheung et al.)



# LHC/Tevatron Reach





\* This is for a squark mass of 750 GeV.

These are large cross-sections.

Some of the parameter space can already be excluded by Tevatron searches...





# Dark Photon + W (work in progress. . .)

Similar to the prompt dark photon production we can consider the associate production of a dark photon together with a W boson. You lose on the cross-section, but you gain from the W mass peak.



# Dark Photon + W (work in progress. . .)

Similar to the prompt dark photon production we can consider the associate production of a dark photon together with a W boson. You lose on the cross-section, but you gain from the W mass peak.



Lepton Jets


# Origin of a Species

Production:

# Origin of a Species

Production:


Prompt Dark Photon




# Origin of a Species

Production:

Prompt Dark Photon




Rare  $Z^0$  decays




# Origin of a Species

Production:


Prompt Dark Photon



Rare  $Z^0$  decays



Gaugino Pair Prod







# Origin of a Species

Production:


Prompt Dark Photon




Rare  $Z^0$  decays



Dark Photon+W




Gaugino Pair Prod




# Origin of a Species

Production:


Prompt Dark Photon




Rare  $Z^0$  decays



Dark Photon+W



Gaugino Pair Prod



Evolution:


Dark Radiation




# Origin of a Species

Production:


Prompt Dark Photon




Rare  $Z^0$  decays



Dark Photon+W



Gaugino Pair Prod




Evolution:

Dark Radiation



Dark Cascades and Lepton Jets




Lepton Jets

GGI

Itay Yavin

# Part III

## Lepton Jets




# Dark Higgs Sector

Something has to spontaneously break the dark gauge-symmetry and I will assume it is some fundamental scalar that gets a vacuum expectation value.

# Dark Higgs Sector

Something has to spontaneously break the dark gauge-symmetry and I will assume it is some fundamental scalar that gets a vacuum expectation value.

$$|D_\mu H_d|^2 \supset g_d^2 v h_d b_\mu b^\mu$$




The dark higgs is unstable,  
but its decay width depends  
on its mass relative to the  
dark photon!

# Dark Higgs Sector

Something has to spontaneously break the dark gauge-symmetry and I will assume it is some fundamental scalar that gets a vacuum expectation value.

$$|D_\mu H_d|^2 \supset g_d^2 v h_d b_\mu b^\mu$$

$$m_h > 2m_b$$




The dark higgs is unstable, but its decay width depends on its mass relative to the dark photon!

# Dark Higgs Sector

Something has to spontaneously break the dark gauge-symmetry and I will assume it is some fundamental scalar that gets a vacuum expectation value.

$$|D_\mu H_d|^2 \supset g_d^2 v h_d b_\mu b^\mu$$



$$m_h > 2m_b$$

Prompt decay  
(Many leptons)


The dark higgs is unstable,  
but its decay width depends  
on its mass relative to the  
dark photon!




# Dark Higgs Sector

Something has to spontaneously break the dark gauge-symmetry and I will assume it is some fundamental scalar that gets a vacuum expectation value.

$$|D_\mu H_d|^2 \supset g_d^2 v h_d b_\mu b^\mu$$




The dark higgs is unstable, but its decay width depends on its mass relative to the dark photon!




# Dark Higgs Sector

Something has to spontaneously break the dark gauge-symmetry and I will assume it is some fundamental scalar that gets a vacuum expectation value.

$$|D_\mu H_d|^2 \supset g_d^2 v h_d b_\mu b^\mu$$




The dark higgs is unstable, but its decay width depends on its mass relative to the dark photon!



# Dark Radiation

Since the dark state are extremely boosted, they will radiate dark gauge-bosons,


$$N_{\gamma'} \sim \frac{\alpha_d}{2\pi} \log \left( \frac{M_{\text{EW}}^2}{M_{\text{dark}}^2} \right)^2 \simeq 1.4 \left( \frac{\alpha_d}{0.1} \right)$$



# Dark Radiation

Since the dark state are extremely boosted, they will radiate dark gauge-bosons,

$$N_{\gamma'} \sim \frac{\alpha_d}{2\pi} \log \left( \frac{M_{\text{EW}}^2}{M_{\text{dark}}^2} \right)^2 \simeq 1.4 \left( \frac{\alpha_d}{0.1} \right)$$




# Cascades in the Dark

After showering finishes, the dark higgses will cascade down to the standard model. If we consider a simple model with 2 dark higgses, then there are several possibilities:


# Cascades in the Dark

After showering finishes, the dark higgses will cascade down to the standard model. If we consider a simple model with 2 dark higgses, then there are several possibilities:



# Cascades in the Dark

After showering finishes, the dark higgses will cascade down to the standard model. If we consider a simple model with 2 dark higgses, then there are several possibilities:



# Lepton Jets

## Lepton Jets -

*Two or more leptons with  $p_T > 10$  GeV inside a cone of  $\Delta R < 0.1$  with hadronic/leptonic isolation cut of  $p_T < 3$  GeV in an annulus of  $0.1 < \Delta R < 0.4$  around the leptons.*




# Lepton Jets

## Lepton Jets -

*Two or more leptons with  $p_T > 10$  GeV inside a cone of  $\Delta R < 0.1$  with hadronic/leptonic isolation cut of  $p_T < 3$  GeV in an annulus of  $0.1 < \Delta R < 0.4$  around the leptons.*

$p_T > 10$  GeV




# Lepton Jets

## Lepton Jets -

*Two or more leptons with  $p_T > 10$  GeV inside a cone of  $\Delta R < 0.1$  with hadronic/leptonic isolation cut of  $p_T < 3$  GeV in an annulus of  $0.1 < \Delta R < 0.4$  around the leptons.*

$0.1 < \Delta R < 0.4$




# Lepton Jets

## Lepton Jets -


*Two or more leptons with  $p_T > 10$  GeV inside a cone of  $\Delta R < 0.1$  with hadronic/leptonic isolation cut of  $p_T < 3$  GeV in an annulus of  $0.1 < \Delta R < 0.4$  around the leptons.*

$$0.1 < \Delta R < 0.4$$



$$\sum p_T < 3 \text{ GeV}$$


# Full Evolution



For simulations: <http://astro.physics.nyu.edu/~iyavin/LeptonJets>

# Lego Plots

For a 500 GeV LSP pair production, the event looks like:



# Experimental Discovery

By defining lepton jets as a searchable object one can look for:

- 1) Lepton-jets + ME
- 2) Lepton-jets + QCD-jets
- 3) Lepton-jets + isolated leptons

While a resonance structure is probably present, since we don't know the mass, it may not very useful to implement mass-window cuts and etc.

# Experimental Efforts

Several experimental groups are working on (designing) searches for lepton-jets


- 1) A. Haas and Y. Gershtein for D0 - Phys. Rev. Lett. 103, 061801 (2009), arXiv:0905:3381
- 2) B. Demirkoz and R. Moore for ATLAS - designing proper triggers for lepton jets.
- 3) K. Cranmer and the NYU group - lepton jet gun.
- 4) H. Lubatti and the Washington group - triggering on long lived neutral particles.
- 5) V. Halyo for CMS - searches for lepton jets.
- 6) Searches at BaBar - See all the local experts.

# Sources of Lepton Jets



# Sources of Lepton Jets



There are different possibilities for obtaining lepton-jets:



# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:


- 1) A non-abelian structure in the dark sector




# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:

- 1) A non-abelian structure in the dark sector



- 2) Dark higgs(es) decay




# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:


- 1) A non-abelian structure in the dark sector



- 2) Dark higgs(es) decay




- 3) Dark radiation




# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:


- 1) A non-abelian structure in the dark sector



- 2) Dark higgs(es) decay



- 3) Dark radiation




There are different ways of producing dark states:


# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:


- 1) A non-abelian structure in the dark sector



- 2) Dark higgs(es) decay



- 3) Dark radiation




There are different ways of producing dark states:

- 1) Prompt dark photon


# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:


- 1) A non-abelian structure in the dark sector



- 2) Dark higgs(es) decay



- 3) Dark radiation




There are different ways of producing dark states:

- 1) Prompt dark photon
- 2) Rare  $Z$  decays


# Sources of Lepton Jets

There are different possibilities for obtaining lepton-jets:


- 1) A non-abelian structure in the dark sector



- 2) Dark higgs(es) decay



- 3) Dark radiation




There are different ways of producing dark states:

- 1) Prompt dark photon

- 2) Rare Z decays

- 3) Susy cascades






# Future Directions

- Searches at LEP and Tevatron.
- Searches at BaBar/Belle (see [Essig, Schuster, and Toro](#)) .
- Tune and modify triggers (see [Demirkoz](#) and [Moore](#)).
- Lepton-Jet observables?
- Other scenarios with similar signatures? (see [Strassler](#) and [Zurek](#)).

Molte Grazie!!!

# Standard Model Background

The SM can give 2 muon pairs recoiling against a jet and that is an irreducible background. Simulation with Madgraph suggest that this is not going to be a serious obstacle:




... and many more ...

$$\sigma = 37 \text{ fb}$$



This can be reduced to  $< 1 \text{ fb}$  by requiring:

$$\Delta R < 0.1$$
$$p_T > 3 \text{ GeV}$$

For all 4 muons.



# Detector Performance



**Bilge Demirköz** and **Roger Moore** investigated ATLAS performance using the prompt dark photon production as a benchmark.

**Bilge Demirköz** also implemented new triggers to help improve the efficiency associated with such events.

# Lepton Jet Efficiency

| Lepton Jet Efficiencies            |              |             |             |              |             |             |
|------------------------------------|--------------|-------------|-------------|--------------|-------------|-------------|
|                                    | 1 Lepton-Jet |             |             | 2 Lepton-Jet |             |             |
| $\text{Br}_{b \rightarrow \pi\pi}$ | 1/7          | 1/3         | 3/5         | 1/7          | 1/3         | 3/5         |
| $\alpha_d$                         |              |             |             |              |             |             |
| 0                                  | 0.49 (0.49)  | 0.47 (0.47) | 0.31 (0.31) | 0.28 (0.28)  | 0.14 (0.15) | 0.05 (0.05) |
| 0.01                               | 0.47 (0.47)  | 0.44 (0.45) | 0.31 (0.32) | 0.3 (0.31)   | 0.16 (0.16) | 0.04 (0.04) |
| 0.03                               | 0.43 (0.41)  | 0.47 (0.48) | 0.3 (0.3)   | 0.27 (0.3)   | 0.14 (0.16) | 0.04 (0.05) |
| 0.1                                | 0.43 (0.39)  | 0.41 (0.44) | 0.29 (0.32) | 0.23 (0.3)   | 0.13 (0.18) | 0.05 (0.07) |
| 0.3                                | 0.38 (0.32)  | 0.34 (0.36) | 0.25 (0.34) | 0.16 (0.3)   | 0.11 (0.22) | 0.05 (0.09) |

**Table 1:** Clean lepton jet efficiencies for different values of the dark gauge-coupling and  $\text{Br}(b \rightarrow \pi^+ \pi^-)$ . The neutralino mass was set to  $\tilde{M} = 300$  GeV. For  $\alpha_d = 0$  dark radiation was switched off. The number of lepton jets increases with  $\alpha_d$  as radiation becomes more likely. The requirement for "clean" lepton jets, as described in the text, results in a decrease in efficiency with the growth of the branching ratio into pion. In brackets are efficiencies for the case where only hadronic isolation is required in the  $0.1 < \Delta R < 0.4$  annulus.