



Results of the **DAMA/LIBRA** experiment

The Dark Matter connection: Theory and Experiment

GGI - Arcetri, Firenze


May 17-21, 2010

P. Belli
INFN-Roma Tor Vergata

Some direct detection processes:

- Scatterings on nuclei

→ detection of nuclear recoil energy



- Inelastic Dark Matter: $W + N \rightarrow W^* + N$

→ W has Two mass states χ_+ , χ_- with δ mass splitting

→ Kinematical constraint for the inelastic scattering of χ_- on a nucleus


$$\frac{1}{2} \mu v^2 \geq \delta \Leftrightarrow v \geq v_{thr} = \sqrt{\frac{2\delta}{\mu}}$$

- Excitation of bound electrons in scatterings on nuclei

→ detection of recoil nuclei + e.m. radiation


- Conversion of particle into e.m. radiation

→ detection of γ , X-rays, e^-



- Interaction only on atomic electrons


→ detection of e.m. radiation



... even WIMPs

- Interaction of light DMp (LDM) on e^- or nucleus with production of a lighter particle

→ detection of electron/nucleus recoil energy



e.g. sterile ν

... also other ideas ...


e.g. signals from these candidates are **completely lost** in experiments based on "rejection procedures" of the e.m. component of their rate

• ... and more

The annual modulation: a model independent signature for the investigation of Dark Matter particles component in the galactic halo

With the present technology, the annual modulation is the main model independent signature for the DM signal. Although the modulation effect is expected to be relatively small **a suitable large-mass, low-radioactive set-up with an efficient control of the running conditions would point out its presence.**

Drukier, Freese, Spergel PRD86
Freese et al. PRD88



- $v_{\text{sun}} \sim 232$ km/s (Sun velocity in the halo)
- $v_{\text{orb}} = 30$ km/s (Earth velocity around the Sun)
- $\gamma = \pi/3$
- $\omega = 2\pi/T$ $T = 1$ year
- $t_0 = 2^{\text{nd}}$ June (when v_{\oplus} is maximum)

$$v_{\oplus}(t) = v_{\text{sun}} + v_{\text{orb}} \cos\gamma \cos[\omega(t-t_0)]$$

$$S_k[\eta(t)] = \int_{\Delta E_k} \frac{dR}{dE_R} dE_R \cong S_{0,k} + S_{m,k} \cos[\omega(t-t_0)]$$

Expected rate in given energy bin changes because the annual motion of the Earth around the Sun moving in the Galaxy

Requirements of the annual modulation


- 1) Modulated rate according cosine
- 2) In a definite low energy range
- 3) With a proper period (1 year)
- 4) With proper phase (about 2 June)
- 5) Just for single hit events in a multi-detector set-up
- 6) With modulation amplitude in the region of maximal sensitivity must be <7% for usually adopted halo distributions, but it can be larger in case of some possible scenarios

To mimic this signature, spurious effects and side reactions must not only - obviously - be able to account for the whole observed modulation amplitude, but also to satisfy contemporaneously all the requirements

The DM annual modulation signature has a different origin and, thus, different peculiarities (e.g. the phase) with respect to those effects connected with the seasons instead



DAMA: an observatory for rare processes @LNGS



DAMA membership

Overall membership in the DAMA activities

Spokesperson: R. Bernabei

P. Belli, R. Bernabei, A. Bussolotti*, F. Montecchia, F. Nozzoli
Dip. di Fisica, Univ. Roma "Tor Vergata" and INFN, sez. Roma Tor Vergata, Italy

F. Cappella, A. d'Angelo, A. Incicchitti, A. Mattei*, D. Prosperi
Dip. di Fisica, Università di Roma "La Sapienza" and INFN, sez. Roma, Italy

R. Cerulli, V. Caracciolo, A. di Marco
INFN - Laboratori Nazionali del Gran Sasso, Italy

C.J. Dai, H.L. He, X.H. Ma, X.D. Sheng, R.G. Wang, Z.P. Ye**
IHEP, Chinese Academy, China;



Università di Roma
Tor Vergata



Institute of High Energy Physics
Chinese Academy of Sciences

+ in some by-product results and small scale experiments:

**F. Danevich, B.V. Grinyov, V.V. Kobychiev,
V.M. Kudovbenko, S.S. Nagorny,
L.L. Nagornaya, D.V. Poda, R.B. Podviyanuk,
O.G. Polischuk, V.I. Tretyak, I. M. Vyshnevskiy,
S.S. Yurchenko and coll.**
Institute for Nuclear Research of Kiev, Ukraine



M. Laubenstein, S. Nisi
INFN - Laboratori Nazionali del Gran Sasso, Italy

S. d'Angelo
Dip. di Fisica and INFN, Università di Roma "Tor Vergata", Italy

**+ in some studies on $\beta^+\beta^+$, EC/ β^+ , EC/EC
decay modes (under the joint Indo-Italian
DST-MAE project):**

**P.K. Raina, A.K. Singh,
P.K. Rath, A. Shukla**
*Indian Institute of Technology,
Kharagpur, India.*



+ in neutron measurements:

M. Angelone, P. Batistoni, M. Pillon
ENEA - C. R. Frascati, Italy

* Technical staff; ** also University of Jing Gangshan, Jiangxi, China.

DAMA/NaI: ≈ 100 kg NaI(Tl)

Performances: N.Cim.A112(1999)545-575, EPJC18(2000)283,
Riv.N.Cim.26 n. 1(2003)1-73, IJMPD13(2004)2127

Results on rare processes:

- Possible Pauli exclusion principle violation PLB408(1997)439
- CNC processes PRC60(1999)065501
- Electron stability and non-paulian transitions in Iodine atoms (by L-shell) PLB460(1999)235
- Search for solar axions PLB515(2001)6
- Exotic Matter search EPJdirect C14(2002)1
- Search for superdense nuclear matter EPJA23(2005)7
- Search for heavy clusters decays EPJA24(2005)51


Results on DM particles:

- PSD PLB389(1996)757
- Investigation on diurnal effect N.Cim.A112(1999)1541
- Exotic Dark Matter search PRL83(1999)4918
- Annual Modulation Signature

PLB424(1998)195, PLB450(1999)448, PRD61(1999)023512, PLB480(2000)23, EPJC18(2000)283,
PLB509(2001)197, EPJC23(2002)61, PRD66(2002)043503, Riv.N.Cim.26 n.1 (2003)1,
IJMPD13(2004)2127, IJMPA21(2006)1445, EPJC47(2006)263, IJMPA22(2007)3155, EPJC53(2008)205,
PRD77(2008)023506, MPLA23(2008)2125.

model independent evidence of a particle DM component in the galactic halo at 6.3σ C.L.

total exposure (7 annual cycles) 0.29 ton x yr



The new DAMA/LIBRA set-up ~250 kg NaI(Tl) (Large sodium Iodide Bulk for Rare processes)

As a result of a second generation R&D for more radiopure NaI(Tl)
by exploiting new chemical/physical radiopurification techniques
(all operations involving crystals and PMTs - including photos - in HP Nitrogen atmosphere)



installing DAMA/LIBRA detectors



assembling a DAMA/ LIBRA detector




filling the inner Cu box with
further shield



detectors during installation; in the
central and right up detectors the new
shaped Cu shield surrounding light
guides (acting also as optical windows)
and PMTs was not yet applied

- *Radiopurity, performances, procedures, etc.:* NIMA592(2008)297
- *Results on DM particles: Annual Modulation Signature:* EPJC56(2008)333, EPJC67(2010)39
- *Results on rare processes: PEP violation in Na and I:* EPJC62(2009)327



closing the Cu box
housing the detectors




view at end of detectors'
installation in the Cu box

The DAMA/LIBRA set-up

For details, radiopurity, performances, procedures, etc.
NIMA592(2008)297


Polyethylene/
paraffin

- 25 x 9.7 kg NaI(Tl) in a 5x5 matrix
- two Suprasil-B light guides directly coupled to each bare crystal
- two PMTs working in coincidence at the single ph. el. threshold



5.5-7.5 phe/keV

Installation




- OFHC low radioactive copper
- Low radioactive lead
- Cadmium foils
- Polyethylene/Paraffin
- Concrete from GS rock




~ 1m concrete from GS rock

- Dismounting/Installing protocol (with "Scuba" system)
- All the materials selected for low radioactivity
- Multicomponent passive shield (>10 cm of Cu, 15 cm of Pb + Cd foils, 10/40 cm Polyethylene/paraffin, about 1 m concrete mostly outside the installation)
- Three-level system to exclude Radon from the detectors
- Calibrations in the same running conditions as production runs
- Installation in air conditioning + huge heat capacity of shield
- Monitoring/alarm system; many parameters acquired with the production data
- Pulse shape recorded by Waveform Analyzer Acqiris DC270 (2chs per detector), 1 Gsample/s, 8 bit, bandwidth 250 MHz
- Data collected from low energy up to MeV region, despite the hardware optimization was done for the low energy



Some on residual contaminants in new ULB NaI(Tl) detectors




α/e pulse shape discrimination has practically 100% effectiveness in the MeV range

The measured α yield in the new DAMA/LIBRA detectors ranges from 7 to some tens α /kg/day

Second generation R&D for new DAMA/LIBRA crystals: new selected powders, physical/chemical radiopurification, new selection of overall materials, new protocol for growing and handling

^{232}Th residual contamination From time-amplitude method. If ^{232}Th chain at equilibrium: it ranges from 0.5 ppt to 7.5 ppt



^{238}U residual contamination First estimate: considering the measured α and ^{232}Th activity, if ^{238}U chain at equilibrium \Rightarrow ^{238}U contents in new detectors typically range from 0.7 to 10 ppt



^{238}U chain splitted into 5 subchains: $^{238}\text{U} \rightarrow ^{234}\text{U} \rightarrow ^{230}\text{Th} \rightarrow ^{226}\text{Ra} \rightarrow ^{210}\text{Pb} \rightarrow ^{206}\text{Pb}$

Thus, in this case: (2.1 ± 0.1) ppt of ^{232}Th ; (0.35 ± 0.06) ppt for ^{238}U
 and: (15.8 ± 1.6) $\mu\text{Bq/kg}$ for $^{234}\text{U} + ^{230}\text{Th}$; (21.7 ± 1.1) $\mu\text{Bq/kg}$ for ^{226}Ra ; (24.2 ± 1.6) $\mu\text{Bq/kg}$ for ^{210}Pb .

$^{\text{nat}}\text{K}$ residual contamination
 The analysis has given for the $^{\text{nat}}\text{K}$ content in the crystals values not exceeding about 20 ppb




^{129}I and ^{210}Pb
 $^{129}\text{I}/^{\text{nat}}\text{I} \approx 1.7 \times 10^{-13}$ for all the new detectors
 ^{210}Pb in the new detectors: $(5 - 30)$ $\mu\text{Bq/kg}$.

No sizable surface pollution by Radon daughters, thanks to the new handling protocols

... more on NIMA592(2008)297


DAMA/LIBRA calibrations

Low energy: various external gamma sources (^{241}Am , ^{133}Ba) and internal X-rays or gamma's (^{40}K , ^{125}I , ^{129}I), routine calibrations with ^{241}Am




$$\frac{\sigma_{LE}}{E} = \frac{(0.448 \pm 0.035)}{\sqrt{E(\text{keV})}} + (9.1 \pm 5.1) \cdot 10^{-3}$$

High energy: external sources of gamma rays (e.g. ^{137}Cs , ^{60}Co and ^{133}Ba) and gamma rays of 1461 keV due to ^{40}K decays in an adjacent detector, tagged by the 3.2 keV X-rays




$$\frac{\sigma_{HE}}{E} = \frac{(1.12 \pm 0.06)}{\sqrt{E(\text{keV})}} + (17 \pm 23) \cdot 10^{-4}$$

Thus, here and hereafter keV means keV electron equivalent



The curves superimposed to the experimental data have been obtained by simulations



Infos about DAMA/LIBRA data taking

Period		Mass (kg)	Exposure (kg × day)	α - β^2
DAMA/LIBRA-1	Sep. 9, 2003 – July 21, 2004	232.8	51405	0.562
DAMA/LIBRA-2	July 21, 2004 – Oct. 28, 2005	232.8	52597	0.467
DAMA/LIBRA-3	Oct. 28, 2005 – July 18, 2006	232.8	39445	0.591
DAMA/LIBRA-4	July 19, 2006 – July 17, 2007	232.8	49377	0.541
DAMA/LIBRA-5	July 17, 2007 – Aug. 29, 2008	232.8	66105	0.468
DAMA/LIBRA-6	Nov. 12, 2008 – Sep. 1, 2009	242.5	58768	0.519
DAMA/LIBRA-1 to -6	Sep. 9, 2003 – Sep. 1, 2009		317697 = 0.87 ton×yr	0.519

- **calibrations: ≈ 72 M events from sources**
- **acceptance window eff: 82 M events (≈ 3 M events/keV)**
- EPJC56(2008)333
- arXiv:1002.1028 (in press on EPJC)

DAMA/NaI (7 years) + DAMA/LIBRA (6 years)

total exposure: 425428 kg×day = 1.17 ton×yr



• First upgrade on Sept 2008:

- replacement of some PMTs in HP N₂ atmosphere
- restore 1 detector to operation
- new Digitizers installed (U1063A Acqiris 1GS/s 8-bit High-Speed cPCI)
- new DAQ system with optical read-out installed

• New upgrade foreseen on fall 2010




... continuously running

Cumulative low-energy distribution of the *single-hit* scintillation events

Single-hit events = each detector has all the others as anticoincidence


(Obviously differences among detectors are present depending e.g. on each specific level and location of residual contaminants, on the detector's location in the 5x5 matrix, etc.)



Efficiencies already accounted for

About the energy threshold:

- The DAMA/LIBRA detectors have been calibrated down to the keV region. This assures a clear knowledge of the "physical" energy threshold of the experiment.
- It obviously profits of the relatively high number of available photoelectrons/keV (from 5.5 to 7.5).
- The two PMTs of each detector in DAMA/LIBRA work in coincidence with hardware threshold at single photoelectron level.
- Effective near-threshold-noise full rejection.
- The software energy threshold used by the experiment is 2 keV.




DAMA/LIBRA-1 to 6 Model Independent Annual Modulation Result

experimental single-hit residuals rate vs time and energy

$\text{Acos}[\omega(t-t_0)]$; continuous lines: $t_0 = 152.5$ d, $T = 1.00$ y

DAMA/LIBRA-1,2,3,4,5,6 (0.87 ton \times yr)

The fit has been done on the DAMA/NaI & DAMA/LIBRA data (1.17 ton \times yr)




2-4 keV

$A = (0.0183 \pm 0.0022)$ cpd/kg/keV

$\chi^2/\text{dof} = 75.7/79$ **8.3 σ C.L.**

Absence of modulation? No

$\chi^2/\text{dof} = 147/80 \Rightarrow P(A=0) = 7 \times 10^{-6}$




2-5 keV

$A = (0.0144 \pm 0.0016)$ cpd/kg/keV

$\chi^2/\text{dof} = 56.6/79$ **9.0 σ C.L.**

Absence of modulation? No

$\chi^2/\text{dof} = 135/80 \Rightarrow P(A=0) = 1.1 \times 10^{-4}$



2-6 keV

$A = (0.0114 \pm 0.0013)$ cpd/kg/keV

$\chi^2/\text{dof} = 64.7/79$ **8.8 σ C.L.**

Absence of modulation? No

$\chi^2/\text{dof} = 140/80 \Rightarrow P(A=0) = 4.3 \times 10^{-5}$

The data favor the presence of a modulated behavior with proper features at 8.8 σ C.L.

Modulation amplitudes measured in each one of the 13 one-year experiments (DAMA/NaI and DAMA/LIBRA)


	A (cpd/kg/keV)	T= 2 π / ω (yr)	t ₀ (day)	C.L.
DAMA/NaI (7 years)				
(2÷4) keV	0.0252 ± 0.0050	1.01 ± 0.02	125 ± 30	5.0 σ
(2÷5) keV	0.0215 ± 0.0039	1.01 ± 0.02	140 ± 30	5.5 σ
(2÷6) keV	0.0200 ± 0.0032	1.00 ± 0.01	140 ± 22	6.3 σ
DAMA/LIBRA (6 years)				
(2÷4) keV	0.0180 ± 0.0025	0.996 ± 0.002	135 ± 8	7.2 σ
(2÷5) keV	0.0134 ± 0.0018	0.997 ± 0.002	140 ± 8	7.4 σ
(2÷6) keV	0.0098 ± 0.0015	0.999 ± 0.002	146 ± 9	6.5 σ
DAMA/NaI + DAMA/LIBRA				
(2÷4) keV	0.0194 ± 0.0022	0.996 ± 0.002	136 ± 7	8.8 σ
(2÷5) keV	0.0149 ± 0.0016	0.997 ± 0.002	142 ± 7	9.3 σ
(2÷6) keV	0.0116 ± 0.0013	0.999 ± 0.002	146 ± 7	8.9 σ

DAMA/NaI (7 annual cycles: 0.29 ton x yr) +
 DAMA/LIBRA (6 annual cycles: 0.87 ton x yr)
 total exposure: 425428 kg×day = 1.17 ton×yr

A, T, t₀ obtained by fitting the single-hit data with $A\cos[\omega(t-t_0)]$

- The modulation amplitudes for the (2 – 6) keV energy interval, obtained when fixing the period at 1 yr and the phase at 152.5 days, are: (0.019±0.003) cpd/kg/keV for DAMA/NaI and (0.010±0.002) cpd/kg/keV for DAMA/LIBRA.
- Thus, their difference: (0.009±0.004) cpd/kg/keV is $\approx 2\sigma$ which corresponds to a modest, but non negligible probability.

The χ^2 test ($\chi^2 = 9.3, 12.2$ and 10.1 over 12 d.o.f. for the three energy intervals, respectively) and the *run test* (lower tail probabilities of 57%, 47% and 35% for the three energy intervals, respectively) accept at 90% C.L. the hypothesis that the modulation amplitudes are normally fluctuating around their best fit values.



Compatibility among the annual cycles

Power spectrum of single-hit residuals

(according to Ap.J.263(1982)835; Ap.J.338(1989)277)




Treatment of the experimental errors and time binning included here

2-6 keV vs 6-14 keV

DAMA/NaI (7 years)
total exposure: 0.29 ton×yr

DAMA/LIBRA (6 years)
total exposure: 0.87 ton×yr

DAMA/NaI (7 years) +
DAMA/LIBRA (6 years)
total exposure: 1.17 ton×yr



Principal mode in the 2-6 keV region:

DAMA/NaI
 $2.737 \cdot 10^{-3} \text{ d}^{-1} \approx 1 \text{ y}^{-1}$

DAMA/LIBRA
 $2.697 \times 10^{-3} \text{ d}^{-1} \approx 1 \text{ yr}^{-1}$

DAMA/NaI+LIBRA
 $2.735 \times 10^{-3} \text{ d}^{-1} \approx 1 \text{ yr}^{-1}$


+

Not present in the 6-14 keV region (only aliasing peaks)

Clear annual modulation is evident in (2-6) keV while it is absent just above 6 keV


Rate behaviour above 6 keV

• No Modulation above 6 keV



Mod. Ampl. (6-10 keV): cpd/kg/keV
 (0.0016 ± 0.0031) DAMA/LIBRA-1
 $-(0.0010 \pm 0.0034)$ DAMA/LIBRA-2
 $-(0.0001 \pm 0.0031)$ DAMA/LIBRA-3
 $-(0.0006 \pm 0.0029)$ DAMA/LIBRA-4
 $-(0.0021 \pm 0.0026)$ DAMA/LIBRA-5
 (0.0029 ± 0.0025) DAMA/LIBRA-6
 → statistically consistent with zero

DAMALIBRA-1 to -6



$\sigma \approx 1\%$, fully accounted by statistical considerations

• No modulation in the whole energy spectrum:

studying integral rate at higher energy, R_{90}

- R_{90} percentage variations with respect to their mean values for single crystal in the DAMA/LIBRA running periods

- Fitting the behaviour with time, adding a term modulated with period and phase as expected for DM particles:

consistent with zero

Period	Mod. Ampl.
DAMA/LIBRA-1	$-(0.05 \pm 0.19)$ cpd/kg
DAMA/LIBRA-2	$-(0.12 \pm 0.19)$ cpd/kg
DAMA/LIBRA-3	$-(0.13 \pm 0.18)$ cpd/kg
DAMA/LIBRA-4	(0.15 ± 0.17) cpd/kg
DAMA/LIBRA-5	(0.20 ± 0.18) cpd/kg
DAMA/LIBRA-6	$-(0.20 \pm 0.16)$ cpd/kg

+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region → $R_{90} \sim$ tens cpd/kg → $\sim 100 \sigma$ far away

No modulation above 6 keV
 This accounts for all sources of bckg and is consistent with studies on the various components


Multiple-hits events in the region of the signal, DAMA/LIBRA 1-6

- Each detector has its own TDs read-out → pulse profiles of multiple-hits events (multiplicity > 1) acquired (exposure: 0.87 ton×yr).
- The same hardware and software procedures as the ones followed for single-hit events

signals by Dark Matter particles do not belong to multiple-hits events, that is:

multiple-hits events = Dark Matter particles events "switched off"

Evidence of annual modulation with proper features as required by the DM annual modulation signature is present in the *single-hit* residuals, while it is absent in the *multiple-hits* residual rate.



This result offers an additional strong support for the presence of Dark Matter particles in the galactic halo, further excluding any side effect either from hardware or from software procedures or from background


Energy distribution of the modulation amplitudes

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)]$$

here $T=2\pi/\omega=1$ yr and $t_0=152.5$ day

DAMA/NaI (7 years) + DAMA/LIBRA (6 years)

total exposure: 425428 kg×day \approx 1.17 ton×yr



A clear modulation is present in the (2-6) keV energy interval, while S_m values compatible with zero are present just above

The S_m values in the (6-20) keV energy interval have random fluctuations around zero with χ^2 equal to 27.5 for 28 degrees of freedom


Statistical distributions of the modulation amplitudes (S_m)

a) S_m for each detector, each annual cycle and each considered energy bin (here 0.25 keV)

b) $\langle S_m \rangle$ = mean values over the detectors and the annual cycles for each energy bin; σ = error associated to the S_m


DAMA/LIBRA (6 years)
total exposure: 0.87 ton \times yr

Each panel refers to each detector separately; 96 entries = 16 energy bins in 2-6 keV energy interval \times 6 DAMA/LIBRA annual cycles (for crys 16, 1 annual cycle, 16 entries)



2-6 keV

Standard deviations of the variable
 $(S_m - \langle S_m \rangle) / \sigma$
for the DAMA/LIBRA detectors



$$x = (S_m - \langle S_m \rangle) / \sigma,$$

$$\chi^2 = \sum x^2$$

Individual S_m values follow a normal distribution since $(S_m - \langle S_m \rangle) / \sigma$ is distributed as a Gaussian with a unitary standard deviation (r.m.s.)



S_m statistically well distributed in all the detectors and annual cycles

Statistical analyses about modulation amplitudes (S_m)


$$x = (S_m - \langle S_m \rangle) / \sigma,$$

$$\chi^2 = \sum x^2$$

$\chi^2/d.o.f.$ values of S_m distributions for each DAMA/LIBRA detector in the (2–6) keV energy interval for the six annual cycles.

DAMA/LIBRA (6 years)

total exposure: 0.87 ton×yr



The $\chi^2/d.o.f.$ values range from 0.7 to 1.22 (96 *d.o.f.* = 16 energy bins \times 6 annual cycles) for 24 detectors \Rightarrow at 95% C.L. the observed annual modulation effect is well distributed in all these detectors.

The remaining detector has $\chi^2/d.o.f. = 1.28$ exceeding the value corresponding to that C.L.; this also is statistically consistent, considering that the expected number of detectors exceeding this value over 25 is 1.25.


- The mean value of the twenty-five points is 1.066, slightly larger than 1. Although this can be still ascribed to statistical fluctuations, let us ascribe it to a possible systematics.
- In this case, one would have an additional error of $\leq 4 \times 10^{-4}$ cpd/kg/keV, if quadratically combined, or $\leq 5 \times 10^{-5}$ cpd/kg/keV, if linearly combined, to the modulation amplitude measured in the (2 – 6) keV energy interval.
- This possible additional error ($\leq 4\%$ or $\leq 0.5\%$, respectively, of the DAMA/LIBRA modulation amplitude) can be considered as an upper limit of possible systematic effects

Energy distributions of cosine (S_m) and sine (Z_m) modulation amplitudes

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)] + Z_m \sin[\omega(t - t_0)]$$


DAMA/NaI (7 years) + DAMA/LIBRA (6 years)

total exposure: 425428 kg×day = 1.17 ton×yr



$t_0 = 152.5 \text{ day (2° June)}$

*maximum at 2° June
as for DM particles*



*maximum at 1° September
T/4 days after 2° June*

The χ^2 test in the (2-14) keV and (2-20) keV energy regions ($\chi^2/\text{dof} = 21.6/24$ and $47.1/36$, probabilities of 60% and 10%, respectively) supports the hypothesis that the $Z_{m,k}$ values are simply fluctuating around zero.

Is there a sinusoidal contribution in the signal? Phase $\neq 152.5$ day?

DAMA/NaI (7 years) + DAMA/LIBRA (6 years)



total exposure: 425428 kg \times day = 1.17 ton \times yr

$$R(t) = S_0 + S_m \cos[\omega(t - t_0)] + Z_m \sin[\omega(t - t_0)] = S_0 + Y_m \cos[\omega(t - t^*)]$$

For Dark Matter signals:

- $|Z_m| \ll |S_m| \approx |Y_m|$
- $\omega = 2\pi/T$
- $t^* \approx t_0 = 152.5d$
- $T = 1 \text{ year}$

Slight differences from 2nd June are expected in case of contributions from non thermalized DM components (as e.g. the SagDEG stream)



E (keV)	S_m (cpd/kg/keV)	Z_m (cpd/kg/keV)	Y_m (cpd/kg/keV)	t^* (day)
2-6	0.0111 ± 0.0013	-0.0004 ± 0.0014	0.0111 ± 0.0013	150.5 ± 7.0
6-14	-0.0001 ± 0.0008	0.0002 ± 0.0005	-0.0001 ± 0.0008	--

Phase as function of energy

$$R(t) = S_0 + Y_m \cos[\omega(t - t^*)]$$

DAMA/NaI (7 years) + DAMA/LIBRA (6 years)

total exposure: 425428 kg×day = 1.17 ton×yr


For DM signals:

$$|Y_m| \approx |S_m|$$

$$t^* \approx t_0 = 152.5d$$

$$\omega = 2\pi/T; \quad T = 1 \text{ year}$$

Slight differences from 2nd June are expected in case of contributions from non thermalized DM components (as the SagDEG stream)



The analysis at energies above 6 keV, the analysis of the multiple-hits events and the statistical considerations about S_m already exclude any sizable presence of systematical effects

Additional investigations on the stability parameters

Modulation amplitudes obtained by fitting the time behaviours of main running parameters, acquired with the production data, when including a DM-like modulation

Running conditions stable at a level better than 1% also in the two new running periods


	DAMA/LIBRA-1	DAMA/LIBRA-2	DAMA/LIBRA-3	DAMA/LIBRA-4	DAMA/LIBRA-5	DAMA/LIBRA-6
Temperature	$-(0.0001 \pm 0.0061) \text{ }^\circ\text{C}$	$(0.0026 \pm 0.0086) \text{ }^\circ\text{C}$	$(0.001 \pm 0.015) \text{ }^\circ\text{C}$	$(0.0004 \pm 0.0047) \text{ }^\circ\text{C}$	$(0.0001 \pm 0.0036) \text{ }^\circ\text{C}$	$(0.0007 \pm 0.0059) \text{ }^\circ\text{C}$
Flux N_2	$(0.13 \pm 0.22) \text{ l/h}$	$(0.10 \pm 0.25) \text{ l/h}$	$-(0.07 \pm 0.18) \text{ l/h}$	$-(0.05 \pm 0.24) \text{ l/h}$	$-(0.01 \pm 0.21) \text{ l/h}$	$-(0.01 \pm 0.15) \text{ l/h}$
Pressure	$(0.015 \pm 0.030) \text{ mbar}$	$-(0.013 \pm 0.025) \text{ mbar}$	$(0.022 \pm 0.027) \text{ mbar}$	$(0.0018 \pm 0.0074) \text{ mbar}$	$-(0.08 \pm 0.12) \times 10^{-2} \text{ mbar}$	$(0.07 \pm 0.13) \times 10^{-2} \text{ mbar}$
Radon	$-(0.029 \pm 0.029) \text{ Bq/m}^3$	$-(0.030 \pm 0.027) \text{ Bq/m}^3$	$(0.015 \pm 0.029) \text{ Bq/m}^3$	$-(0.052 \pm 0.039) \text{ Bq/m}^3$	$(0.021 \pm 0.037) \text{ Bq/m}^3$	$-(0.028 \pm 0.036) \text{ Bq/m}^3$
Hardware rate above single photoelectron	$-(0.20 \pm 0.18) \times 10^{-2} \text{ Hz}$	$(0.09 \pm 0.17) \times 10^{-2} \text{ Hz}$	$-(0.03 \pm 0.20) \times 10^{-2} \text{ Hz}$	$(0.15 \pm 0.15) \times 10^{-2} \text{ Hz}$	$(0.03 \pm 0.14) \times 10^{-2} \text{ Hz}$	$(0.08 \pm 0.11) \times 10^{-2} \text{ Hz}$

All the measured amplitudes well compatible with zero
 + none can account for the observed effect


(to mimic such signature, spurious effects and side reactions must not only be able to account for the whole observed modulation amplitude, but also simultaneously satisfy all the 6 requirements)

Summarizing on a hypothetical background modulation in DAMA/LIBRA 1-6

- No Modulation above 6 keV
- No modulation in the whole energy spectrum




+ if a modulation present in the whole energy spectrum at the level found in the lowest energy region $\rightarrow R_{90} \sim$ tens cpd/kg $\rightarrow \sim 100 \sigma$ far away



- No modulation in the 2-6 keV *multiple-hits* residual rate

No background modulation (and cannot mimic the signature): all this accounts for the all possible sources of bckg



Nevertheless, additional investigations performed ...



- Three examples for specific cases in the following:
1. The muon case
 2. The ^{40}K case
 3. The neutron case

The μ case

MonteCarlo simulation

- muon intensity distribution
- Gran Sasso rock overburden map

events where just one detector fires



Case of fast neutrons produced by μ

Φ_μ @ LNGS $\approx 20 \mu \text{ m}^{-2} \text{ d}^{-1}$ ($\pm 2\%$ modulated)
 Measured neutron Yield @ LNGS: $Y=1\div 7 \cdot 10^{-4} \text{ n}/\mu/(\text{g}/\text{cm}^2)$
 $R_n = (\text{fast n by } \mu)/(\text{time unit}) = \Phi_\mu Y M_{\text{eff}}$

Hyp.: $M_{\text{eff}} = 15 \text{ tons}$; $g \approx \varepsilon \approx f_{\Delta E} \approx f_{\text{single}} \approx 0.5$ (cautiously)
 Knowing that: $M_{\text{setup}} \approx 250 \text{ kg}$ and $\Delta E=4\text{keV}$

Annual modulation amplitude at low energy due to μ modulation:

$$S_m^{(\mu)} = R_n g \varepsilon f_{\Delta E} f_{\text{single}} 2\% / (M_{\text{setup}} \Delta E)$$

g = geometrical factor; ε = detection effic. by elastic scattering
 $f_{\Delta E}$ = energy window ($E>2\text{keV}$) effic.; f_{single} = single hit effic.

$$S_m^{(\mu)} < (0.4\div 3) \times 10^{-5} \text{ cpd}/\text{kg}/\text{keV}$$

Moreover, this modulation also induces a variation in other parts of the energy spectrum and in the *multi-hits* events

It cannot mimic the signature: already excluded also by R_{90} , by *multi-hits* analysis + different phase, etc.

Can (whatever) hypothetical cosmogenic products be considered

as side effects, assuming that they might produce:

- only events at low energy,
- only *single-hit* events,
- no sizable effect in the *multiple-hit* counting rate

?

But, its phase should be (much) larger than μ phase, t_μ :

- if $\tau \ll T/2\pi$: $t_{\text{side}} = t_\mu + \tau$
- if $\tau \gg T/2\pi$: $t_{\text{side}} = t_\mu + T/4$

It cannot mimic the signature: different phase

The phase of the muon flux at LNGS is roughly around middle of July and largely variable from year to year. Last meas. by LVD partially overlapped with DAMA/NaI and fully with DAMA/LIBRA: 1.5% modulation and phase=July 5th $\pm 15 \text{ d}$.

DAMA/NaI + DAMA/LIBRA
 measured a stable phase: May, 26th $\pm 7 \text{ days}$

This phase is 7.3σ far from July 15th and is 5.9σ far from July 5th

R_{90} , multi-hits, phase, and other analyses

NO


No role for ^{40}K in the S_m spectrum

see arXiv:0912.0660

not evidence for any 3 keV peak in the S_m distribution. Behaviour compatible with a **monotonic behaviour** and with a “**structure**” as expected for many Dark Matter candidates (also for WIMP)

The annual modulation signal is equally distributed over all the detectors

Stability of efficiency



^{40}K decay cannot give any modulation, unless evoking new exotic physics.

No modulation has been observed in other energy regions where ^{40}K decay contributes

No modulation in multiple-hit events in 2-6 keV (^{40}K can give double events in two adjacent detectors and multi-site events due to Compton scatterings)

No modulation of the **double coincidence events**, 1461 keV-3 keV

The annual modulation signal is present in the outer and in the inner detectors.

(no dependence on the veto capability, that is different – for geometrical reasons – among the detectors)

The analysis of the double coincidences rules out at more than 10σ any modulation around 3 keV in the *single-hit* events from the hypothetical cases of : i) ^{40}K “exotic” modulation decay; ii) spill-out from double to single events and viceversa.

Even assuming the arXiv:0808.3283 scenario:





- the expected single hit modulation amplitude would be much below the measured modulation amplitude
- the phase (3 jan) would be well different from the measured phase (26 may \pm 7 day).

No role for ^{40}K in the experimental S_m

also see arXiv:0912.0660

The experimental S_m cannot be due to ^{40}K for many reasons.

No modulation of the double coincidence events (1461 keV-3 keV).



The ^{40}K double coincidence events are not modulated

Any modulation contribution around 3 keV in the single-hit events from the hypothetical cases of: i) ^{40}K "exotic" modulated decay; ii) spill-out effects from double to single events and viceversa, is ruled out at more than 10σ

Can a possible thermal neutron modulation account for the observed effect?

NO

• Thermal neutrons flux measured at LNGS :


$$\Phi_n = 1.08 \cdot 10^{-6} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (N.Cim.A101(1989)959)}$$

• Experimental upper limit on the thermal neutrons flux “surviving” the neutron shield in DAMA/LIBRA:

➤ studying triple coincidences able to give evidence for the possible presence of ^{24}Na from neutron activation:

$$\Phi_n < 1.2 \times 10^{-7} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (90\% C.L.)}$$

• Two consistent upper limits on thermal neutron flux have been obtained with DAMA/NaI considering the same capture reactions and using different approaches.



Evaluation of the expected effect:

▶ Capture rate = $\Phi_n \sigma_n N_T < 0.022$ captures/day/kg

HYPOTHESIS: assuming very cautiously a 10% thermal neutron modulation:

➔ $S_m^{(\text{thermal n})} < 0.8 \times 10^{-6}$ cpd/kg/keV ($< 0.01\%$ S_m^{observed})

In all the cases of neutron captures (^{24}Na , ^{128}I , ...) a possible thermal n modulation induces a variation in all the energy spectrum


Already excluded also by R_{90} analysis

MC simulation of the process

When $\Phi_n = 10^{-6} \text{ n cm}^{-2} \text{ s}^{-1}$:

$7 \cdot 10^{-5}$ cpd/kg/keV

$1.4 \cdot 10^{-3}$ cpd/kg/keV



E (MeV)

Can a possible fast neutron modulation account for the observed effect?

NO

In the estimate of the possible effect of the neutron background cautiously not included the 1m concrete moderator, which almost completely surrounds (mostly outside the barrack) the passive shield

Measured fast neutron flux @ LNGS:

$$\Phi_n = 0.9 \cdot 10^{-7} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (Astropart.Phys.4 (1995)23)}$$

By MC: differential counting rate above 2 keV $\approx 10^{-3}$ cpd/kg/keV

HYPOTHESIS: assuming - very cautiously - a 10% neutron modulation: $\Rightarrow S_m^{(\text{fast n})} < 10^{-4}$ cpd/kg/keV ($< 0.5\% S_m^{\text{observed}}$)

• Experimental upper limit on the fast neutrons flux “surviving” the neutron shield in DAMA/LIBRA:

➤ through the study of the inelastic reaction $^{23}\text{Na}(n,n')^{23}\text{Na}^*(2076 \text{ keV})$ which produces two γ 's in coincidence (1636 keV and 440 keV):

$$\Phi_n < 2.2 \times 10^{-7} \text{ n cm}^{-2} \text{ s}^{-1} \text{ (90\%C.L.)}$$

➤ well compatible with the measured values at LNGS. This further excludes any presence of a fast neutron flux in DAMA/LIBRA significantly larger than the measured ones.

Moreover, a possible fast n modulation would induce:

▶ a variation in all the energy spectrum (steady environmental fast neutrons always accompanied by thermalized component)

already excluded also by R_{90}

▶ a modulation amplitude for multiple-hit events different from zero


already excluded by the multiple-hit events

Thus, a possible 5% neutron modulation (ICARUS TM03-01) cannot quantitatively contribute to the DAMA/NaI observed signal, even if the neutron flux would be assumed 100 times larger than measured by various authors over more than 15 years @ LNGS


Summary of the results obtained in the additional investigations of possible systematics or side reactions: DAMA/LIBRA-1 to 6

(previous exposure and details see: NIMA592(2008)297, EPJC56(2008)333, arXiv:0912.4200)

<i>Source</i>	<i>Main comment</i>	<i>Cautious upper limit (90%C.L.)</i>
RADON	Sealed Cu box in HP Nitrogen atmosphere, 3-level of sealing, etc.	$<2.5 \times 10^{-6}$ cpd/kg/keV
TEMPERATURE	Installation is air conditioned+ detectors in Cu housings directly in contact with multi-ton shield→ huge heat capacity + T continuously recorded	$<10^{-4}$ cpd/kg/keV
NOISE	Effective full noise rejection near threshold	$<10^{-4}$ cpd/kg/keV
ENERGY SCALE	Routine + intrinsic calibrations	$<1-2 \times 10^{-4}$ cpd/kg/keV
EFFICIENCIES	Regularly measured by dedicated calibrations	$<10^{-4}$ cpd/kg/keV
BACKGROUND	No modulation above 6 keV; no modulation in the (2-6) keV <i>multiple-hits</i> events; this limit includes all possible sources of background	$<10^{-4}$ cpd/kg/keV
SIDE REACTIONS	Muon flux variation measured at LNGS	$<3 \times 10^{-5}$ cpd/kg/keV



+ they cannot satisfy all the requirements of annual modulation signature



Thus, they can not mimic the observed annual modulation effect

Summarizing

- Presence of modulation for 13 annual cycles at 8.9σ C.L. with the proper distinctive features of the DM signature; all the features satisfied by the data over 13 independent experiments of 1 year each one
- The total exposure by former DAMA/NaI and present DAMA/LIBRA is **1.17 ton × yr (13 annual cycles)**
- In fact, as required by the DM annual modulation signature:

1)

The *single-hit* events show a clear cosine-like modulation, as expected for the DM signal

2)

Measured period is equal to (0.999 ± 0.002) yr, well compatible with the 1 yr period, as expected for the DM signal

3)

Measured phase (146 ± 7) days is well compatible with the roughly about 152.5 days as expected for the DM signal

4)

The modulation is present only in the low energy (2–6) keV energy interval and not in other higher energy regions, consistently with expectation for the DM signal

5)

The modulation is present only in the *single-hit* events, while it is absent in the *multiple-hit* ones as expected for the DM signal

6)

The measured modulation amplitude in NaI(Tl) of the *single-hit* events in the (2–6) keV energy interval is: (0.0116 ± 0.0013) cpd/kg/keV (8.9σ C.L.).

No systematic or side process able to simultaneously satisfy all the many peculiarities of the signature and to account for the whole measured modulation amplitude is available

Model-independent evidence by DAMA/NaI and DAMA/LIBRA

well compatible with several candidates in many astrophysical, nuclear and particle physics scenarios

No other experiment whose result can be directly compared in model independent way with those of DAMA/NaI and DAMA/LIBRA available


Available results from direct searches using different target materials and approaches do not give any robust conflict

Possible model dependent positive hints from indirect searches not in conflict with DAMA; but interpretation and the evidence itself in indirect searches depend e.g. on bckg modeling (also including pulsars, supernovae remnants, ...), on DM spatial velocity distribution, either on forced boost factor or on unnatural clumpiness, etc.

Moreover, whatever hints from other direct searches must be interpreted; in any case large room of compatibility with DAMA is present

Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$

WIMP DM candidate (as in [4])
 considering elastic scattering on nuclei
 SI dominant coupling
 $v_0 = 170 \text{ km/s}$




About the same C.L.

...scaling from NaI




Curve label	Halo model (see ref. [4, 34])	Local density (GeV/cm ³)	Set as in [4]	DM particle mass	$\xi\sigma_{SI}$ (pb)
<i>a</i>	A5 (NFW)	0.2	A	15 GeV	3.1×10^{-4}
<i>b</i>	A5 (NFW)	0.2	A	15 GeV	1.3×10^{-5}
<i>c</i>	A5 (NFW)	0.2	B	60 GeV	5.5×10^{-6}
<i>d</i>	B3 (Evans power law)	0.17	B	100 GeV	6.5×10^{-6}
<i>e</i>	B3 (Evans power law)	0.17	A	120 GeV	1.3×10^{-5}

[4] RNC 26 (2003) 1; [34] PRD66 (2002) 043503




channeling contribution as in EPJC53(2008)205 considered for curve *b*

Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



$\theta = 2.435$

WIMP DM candidate as in [4]
Elastic scattering on nuclei
SI & SD mixed coupling
 $v_0 = 170$ km/s




- Not best fit
- About the same C.L.



...scaling from NaI

Curve label	Halo model (see ref. [4, 34])	Local density (GeV/cm^3)	Set as in [4]	DM particle mass	$\xi\sigma_{SI}$ (pb)	$\xi\sigma_{SD}$ (pb)
<i>f</i>	A5 (NFW)	0.2	A	15 GeV	10^{-7}	2.6
<i>g</i>	A5 (NFW)	0.2	A	15 GeV	1.4×10^{-4}	1.4
<i>h</i>	A5 (NFW)	0.2	B	60 GeV	10^{-7}	1.4
<i>i</i>	A5 (NFW)	0.2	B	60 GeV	8.7×10^{-6}	8.7×10^{-2}
<i>j</i>	B3 (Evans power law)	0.17	A	100 GeV	10^{-7}	1.7
<i>k</i>	B3 (Evans power law)	0.17	A	100 GeV	1.1×10^{-5}	0.11



Examples for few of the many possible scenarios superimposed to the measured modulation amplitudes $S_{m,k}$



LDM candidate
 (as in MPLA23(2008)2125):
 inelastic interaction
 with electron or nucleus
 targets



Light bosonic candidate
 (as in IJMPA21(2006)1445):
 axion-like particles totally
 absorbed by target material



- Not best fit
- About the same C.L.

curve r: also pseudoscalar axion-like candidates (e.g. majoron)
 $m_a = 3.2 \text{ keV } g_{aee} = 3.9 \cdot 10^{-11}$

(NFW) halo model as in [4, 34], local density = 0.17 GeV/cm³, local velocity = 170 km/s

Curve label	DM particle	Interaction	Set as in [4]	m_H	Δ	Cross section (pb)
<i>l</i>	LDM	coherent on nuclei	A	30 MeV	18 MeV	$\xi \sigma_m^{coh} = 1.8 \times 10^{-6}$
<i>m</i>	LDM	coherent on nuclei	A	100 MeV	55 MeV	$\xi \sigma_m^{coh} = 2.8 \times 10^{-6}$
<i>n</i>	LDM	incoherent on nuclei	A	30 MeV	3 MeV	$\xi \sigma_m^{inc} = 2.2 \times 10^{-2}$
<i>o</i>	LDM	incoherent on nuclei	A	100 MeV	55 MeV	$\xi \sigma_m^{inc} = 4.6 \times 10^{-2}$
<i>p</i>	LDM	coherent on nuclei	A	28 MeV	28 MeV	$\xi \sigma_m^{coh} = 1.6 \times 10^{-6}$
<i>q</i>	LDM	incoherent on nuclei	A	88 MeV	88 MeV	$\xi \sigma_m^{inc} = 4.1 \times 10^{-2}$
<i>r</i>	LDM	on electrons	-	60 keV	60 keV	$\xi \sigma_m^e = 0.3 \times 10^{-6}$

About interpretation

- ✓ Not a unique reference model for Dark Matter particles
- ✓ Not a single set of assumptions for parameters in the astrophysical, nuclear and particle physics related arguments
- ✓ Often comparisons are made in inconsistent way


model-dependent analysis: selecting just one model framework by fixing many parameters and by adopting several (astrophysical, nuclear and particle physics) assumptions

- which particle?
- which interaction couplings?
- which Form Factors for each target-material?
- which Spin Factors?
- which nuclear model framework?
- which scaling laws?
- which halo model, profile and parameters?
- is there a presence of non-thermalized components in the halo parameters?
- which velocity distribution?
- which parameters for velocity distribution?
- which instrumental quantities?
- ...

Exclusion plots have no “universal validity” (they depend on the recipe)


For example, which L_{eff} in liquid Xenon experiments?
arXiv:0909.1063, arXiv:1005.0838

No experiment can be directly compared in model independent way with DAMA



... and experimental aspects ...

- Marginal and “selected” exposures. Threshold, small detector response (few phe/keV), energy scale and energy resolution; calibrations in other energy region. Stability of all the operating conditions. Selections of detectors and of data. Handling of (many) “subtraction” procedures and stability in time of all the selection windows and related quantities, etc. Efficiencies. Fiducial volume vs disuniformity of detector, response in liquids? Used values in the calculation (q.f., etc.). Used approximations. ...



Perspectives of DAMA/LIBRA

- Continuously running


- Next upgrade: replacement of all the PMTs with higher Quantum Efficiency (Q.E.) PMTs.

- New PMTs with higher Q.E. in production: 16 prototypes already tested; five of them have been accepted; 4 new prototypes at hand now

- Continuing data taking for many years in the new configuration.


- Special data taking for other rare processes.

- Update corollary analyses with the new data to disentangle among the many possible scenarios for DM candidates, interactions, halo models, nuclear/atomic properties, etc..



•Goals:

- lowering the energy threshold (presently, at 2 keV)
- improvement of the acceptance efficiency
- increase the sensitivity in the *model independent* analysis (amplitude, phase, second order effects, ...)
- improvement of the sensitivity in the *model dependent* analyses, allowing to better disentangle several astrophysical, particle physics and nuclear physics scenarios



Conclusions



- Positive evidence for the presence of DM particles in the galactic halo now supported at 8.9σ C.L. by the cumulative $1.17 \text{ ton} \times \text{yr}$ exposure over 13 annual cycles by the former DAMA/NaI and the present DAMA/LIBRA
- The modulation parameters determined with better precision
- Full sensitivity to many kinds of DM candidates and interactions both inducing recoils and/or e.m. radiation
- Updated/new model dependent corollary investigations on the nature of the DM particle in progress also in the light of some recent strongly model dependent claims
- Investigations other than DM

What next?

- Upgrade in fall 2010 substituting all the PMTs with new ones having higher Q.E. to lower the experimental energy threshold, improve general features and disentangle among at least some of the possible scenarios
- Collect a suitable exposure in the new running conditions
- Investigate second order effects
- R&D toward a 1 ton ULB NaI(Tl) set-up experiment proposed in 1996 as a further step for an ultimate multi-ton & multi-purpose NaI(Tl) experiment



... and ...