

Neutralino Dark Matter in the BMSSM

Nicolás Bernal



CFTP - IST, Lisbon

June 3rd 2010

JCAP 03(2010)007

NB, A. Goudelis

JHEP 08(2009)053

NB, K. Blum, M. Losada, Y. Nir

Outline

- 1 Motivation
- 2 The BMSSM
- 3 Dark Matter
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 Dark Matter Direct Detection
- 5 Dark Matter Indirect Detection
 - γ -rays
 - Positrons
 - Antiprotons
- 6 Summary

Outline

- 1 Motivation**
- 2 The BMSSM**
- 3 Dark Matter**
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 Dark Matter Direct Detection**
- 5 Dark Matter Indirect Detection**
 - γ -rays
 - Positrons
 - Antiprotons
- 6 Summary**

MSSM Higgs potential

The MSSM contains 2 doublets of complex scalar fields of opposite hypercharge:

$$H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}$$

MSSM Higgs potential

The MSSM contains 2 doublets of complex scalar fields of opposite hypercharge:

$$H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}$$

Full tree-level scalar Higgs potential:

$$V_H = (|\mu|^2) |H_u|^2 + (|\mu|^2) |H_d|^2$$

- Quadratic terms comes from F terms in the superpotential

μ : higgsino mass parameter

MSSM Higgs potential

The MSSM contains 2 doublets of complex scalar fields of opposite hypercharge:

$$H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}$$

Full tree-level scalar Higgs potential:

$$V_H = (|\mu|^2 + m_{H_u}^2) |H_u|^2 + (|\mu|^2 + m_{H_d}^2) |H_d|^2 - \mu B (H_u H_d + \text{h.c.})$$

- Quadratic terms comes from F terms in the superpotential and SUSY-breaking terms

μ : higgsino mass parameter

m_H and B : SUSY-breaking mass parameters

MSSM Higgs potential

The MSSM contains 2 doublets of complex scalar fields of opposite hypercharge:

$$H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}$$

Full tree-level scalar Higgs potential:

$$V_H = (|\mu|^2 + m_{H_u}^2) |H_u|^2 + (|\mu|^2 + m_{H_d}^2) |H_d|^2 - \mu B (H_u H_d + \text{h.c.}) \\ + \frac{g_1^2 + g_2^2}{8} (|H_u|^2 - |H_d|^2)^2 + \frac{1}{2} g_2^2 |H_d^\dagger H_u|^2$$

- Quadratic terms comes from F terms in the superpotential and SUSY-breaking terms
 - μ : higgsino mass parameter
 - m_H and B : SUSY-breaking mass parameters
- Quartic terms comes from D terms \rightarrow pure gauge couplings!

MSSM Higgs potential

The MSSM contains 2 doublets of complex scalar fields of opposite hypercharge:

$$H_u = \begin{pmatrix} H_u^+ \\ H_u^0 \end{pmatrix} \quad H_d = \begin{pmatrix} H_d^0 \\ H_d^- \end{pmatrix}$$

Full tree-level scalar Higgs potential:

$$V_H = (|\mu|^2 + m_{H_u}^2) |H_u|^2 + (|\mu|^2 + m_{H_d}^2) |H_d|^2 - \mu B (H_u H_d + \text{h.c.}) \\ + \frac{g_1^2 + g_2^2}{8} (|H_u|^2 - |H_d|^2)^2 + \frac{1}{2} g_2^2 |H_d^\dagger H_u|^2$$

- Quadratic terms comes from F terms in the superpotential and SUSY-breaking terms
 - μ : higgsino mass parameter
 - m_H and B : SUSY-breaking mass parameters
- Quartic terms comes from D terms \rightarrow pure gauge couplings!
- $\rightarrow V_H$ is CP conserving (even though the full L violates CP)

MSSM Higgs potential

The neutral components of the 2 Higgs fields develop vevs:

$$\langle H_u \rangle = v_u = v \sin\beta \qquad \langle H_d \rangle = v_d = v \cos\beta \qquad v \sim 174\text{GeV}$$

EW symmetry breaking: $SU(2)_L \times U(1)_Y \rightarrow U(1)_{\text{EW}}$

The spectrum contains:

- h and H : 2 CP even Higgs bosons
- A : 1 CP odd Higgs boson
- H^+ and H^- : 2 charged Higgs bosons

Tree level Higgs spectrum

In terms of M_A and $\tan\beta$ the tree level Higgs spectrum is

$$m_h^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 - \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$

$$m_H^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 + \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$

$$m_{H^\pm}^2 = m_A^2 + m_W^2$$

Tree level Higgs spectrum

In terms of M_A and $\tan\beta$ the tree level Higgs spectrum is

$$m_h^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 - \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$
$$m_H^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 + \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$
$$m_{H^\pm}^2 = m_A^2 + m_W^2$$

Important constraint: $m_h \leq \text{Min}(m_A, m_Z) |\cos 2\beta| \leq m_Z$

Tree level Higgs spectrum

In terms of M_A and $\tan\beta$ the tree level Higgs spectrum is

$$m_h^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 - \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$
$$m_H^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 + \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$
$$m_{H^\pm}^2 = m_A^2 + m_W^2$$

Important constraint: $m_h \leq \text{Min}(m_A, m_Z) |\cos 2\beta| \leq m_Z$

The LEP II bound $m_h \gtrsim 114 \text{ GeV}$ is already violated!

Tree level Higgs spectrum

In terms of M_A and $\tan\beta$ the tree level Higgs spectrum is

$$m_h^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 - \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$
$$m_H^2 = \frac{1}{2} \left[m_Z^2 + m_A^2 + \sqrt{(m_A^2 - m_Z^2)^2 + 4 m_A^2 m_Z^2 \sin^2 2\beta} \right]$$
$$m_{H^\pm}^2 = m_A^2 + m_W^2$$

Important constraint: $m_h \leq \text{Min}(m_A, m_Z) |\cos 2\beta| \leq m_Z$

The LEP II bound $m_h \gtrsim 114 \text{ GeV}$ is already violated!

→ To avoid a contradiction we need both
large $\tan\beta$ and large radiative corrections

Radiative corrections

Most important RC comes from loops of tops and stops:

$$\delta_{1\text{-loop}} m_h^2 \sim \frac{12}{16\pi} \left[\ln \frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} + \frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right. \\ \left. + \frac{1}{2} \left(\frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \right)^2 \left(2 - \frac{m_{\tilde{t}_1}^2 + m_{\tilde{t}_2}^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right) \right]$$

$$X_t \equiv A_t - \mu \cot \beta$$

Radiative corrections

Most important RC comes from loops of tops and stops:

$$\delta_{1\text{-loop}} m_h^2 \sim \frac{12}{16\pi} \left[\ln \frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} + \frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right. \\ \left. + \frac{1}{2} \left(\frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \right)^2 \left(2 - \frac{m_{\tilde{t}_1}^2 + m_{\tilde{t}_2}^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right) \right]$$

$$X_t \equiv A_t - \mu \cot \beta$$

Consistency with LEP II achieved with

- **Heavy stops** $m_{\tilde{t}} \sim 600$ GeV to few TeV
- ✗ However, the superpartners make the theory natural and they should not be too heavy

Radiative corrections

Most important RC comes from loops of tops and stops:

$$\delta_{1\text{-loop}} m_h^2 \sim \frac{12}{16\pi} \left[\ln \frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} + \frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} + \frac{1}{2} \left(\frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \right)^2 \left(2 - \frac{m_{\tilde{t}_1}^2 + m_{\tilde{t}_2}^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right) \right]$$

$$X_t \equiv A_t - \mu \cot \beta$$

Consistency with LEP II achieved with

- Heavy stops $m_{\tilde{t}} \sim 600$ GeV to few TeV
- Large stop mixing
- ✗ However, large A_t -terms are hard to achieve in specific models of SUSY breaking

Radiative corrections

Most important RC comes from loops of tops and stops:

$$\delta_{1\text{-loop}} m_h^2 \sim \frac{12}{16\pi} \left[\ln \frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} + \frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right. \\ \left. + \frac{1}{2} \left(\frac{|X_t|^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \right)^2 \left(2 - \frac{m_{\tilde{t}_1}^2 + m_{\tilde{t}_2}^2}{m_{\tilde{t}_1}^2 - m_{\tilde{t}_2}^2} \ln \frac{m_{\tilde{t}_1}^2}{m_{\tilde{t}_2}^2} \right) \right]$$

$$X_t \equiv A_t - \mu \cot \beta$$

Consistency with LEP II achieved with

- Heavy stops $m_{\tilde{t}} \sim 600$ GeV to few TeV
- Large stop mixing

✗ SUSY Little Hierarchy Problem

Outline

- 1 Motivation
- 2 **The BMSSM**
- 3 **Dark Matter**
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 **Dark Matter Direct Detection**
- 5 **Dark Matter Indirect Detection**
 - γ -rays
 - Positrons
 - Antiprotons
- 6 **Summary**

Corrections to the MSSM

Assume that there is New Physics beyond the MSSM at a scale M , much above the electroweak scale m_Z and the scale of the SUSY breaking terms m_{susy} .

$$\epsilon \sim \frac{m_{\text{susy}}}{M} \sim \frac{m_Z}{M} \ll 1$$

The corrections to the MSSM can be parametrized by operators suppressed by inverse powers of M ; i.e. by powers of ϵ .

Corrections to the MSSM

Assume that there is New Physics beyond the MSSM at a scale M , much above the electroweak scale m_Z and the scale of the SUSY breaking terms m_{susy} .

$$\epsilon \sim \frac{m_{\text{susy}}}{M} \sim \frac{m_Z}{M} \ll 1$$

The corrections to the MSSM can be parametrized by operators suppressed by inverse powers of M ; i.e. by powers of ϵ .

→ There can be significant effects from non-renormalizable terms on the same order as the one-loop terms.

We focus on an effective action analysis to the Higgs sector as an approach to consider the effects of **New Physics Beyond the MSSM**.

Non-renormalizable operators

Remember the ordinary MSSM superpotential:

$$W_{\text{MSSM}} \supset \int d^2\theta \mu H_u H_d$$

Non-renormalizable operators

Remember the ordinary MSSM superpotential:

$$W_{\text{MSSM}} \supset \int d^2\theta \mu H_u H_d$$

There are only 2 operators at order $\frac{1}{M}$:

$$O_1 = \frac{1}{M} \int d^2\theta (H_u H_d)^2$$

$$O_2 = \frac{1}{M} \int d^2\theta Z (H_u H_d)^2$$

$Z \equiv \theta^2 m_{\text{susy}}$: spurion field

O_1 : is a dimension 5 SUSY operator

O_2 : parametrizes SUSY breaking

→ Both operators can lead to CP violation

BMSSM Higgs potential

Corrections to the MSSM Higgs potential

$$\begin{aligned} \delta L = & 2 \epsilon_1 H_u H_d \left(H_u^\dagger H_u + H_d^\dagger H_d \right) + \epsilon_2 (H_u H_d)^2 + \text{h.c.} \\ & + \frac{\epsilon_1}{\mu^*} \left[2(H_u H_d)(\tilde{H}_u \tilde{H}_d) + 2(\tilde{H}_u H_d)(H_u \tilde{H}_d) \right. \\ & \left. + (H_u \tilde{H}_d)(H_u \tilde{H}_d) + (\tilde{H}_u H_d)(\tilde{H}_u H_d) \right] + \text{h.c.} \end{aligned}$$

where

$$\epsilon_1 \equiv \frac{\mu^* \lambda_1}{M} \quad \epsilon_2 \equiv -\frac{m_{\text{susy}} \lambda_2}{M}$$

BMSSM Higgs potential

Corrections to the MSSM Higgs potential

$$\begin{aligned}\delta L = & 2 \epsilon_1 H_u H_d \left(H_u^\dagger H_u + H_d^\dagger H_d \right) + \epsilon_2 (H_u H_d)^2 + \text{h.c.} \\ & + \frac{\epsilon_1}{\mu^*} \left[2(H_u H_d)(\tilde{H}_u \tilde{H}_d) + 2(\tilde{H}_u H_d)(H_u \tilde{H}_d) \right. \\ & \left. + (H_u \tilde{H}_d)(H_u \tilde{H}_d) + (\tilde{H}_u H_d)(\tilde{H}_u H_d) \right] + \text{h.c.}\end{aligned}$$

where

$$\epsilon_1 \equiv \frac{\mu^* \lambda_1}{M} \quad \epsilon_2 \equiv -\frac{m_{\text{susy}} \lambda_2}{M}$$

- New contributions for Higgs boson masses

BMSSM Higgs potential

Corrections to the MSSM Higgs potential

$$\begin{aligned}\delta L = & 2 \epsilon_1 H_u H_d \left(H_u^\dagger H_u + H_d^\dagger H_d \right) + \epsilon_2 (H_u H_d)^2 + \text{h.c.} \\ & + \frac{\epsilon_1}{\mu^*} \left[2(H_u H_d)(\tilde{H}_u \tilde{H}_d) + 2(\tilde{H}_u H_d)(H_u \tilde{H}_d) \right. \\ & \left. + (H_u \tilde{H}_d)(H_u \tilde{H}_d) + (\tilde{H}_u H_d)(\tilde{H}_u H_d) \right] + \text{h.c.}\end{aligned}$$

where

$$\epsilon_1 \equiv \frac{\mu^* \lambda_1}{M} \quad \epsilon_2 \equiv -\frac{m_{\text{susy}} \lambda_2}{M}$$

- New contributions for Higgs boson masses
- New contributions for higgsino (χ^0 and χ^\pm) masses

BMSSM Higgs potential

Corrections to the MSSM Higgs potential

$$\begin{aligned} \delta L = & 2 \epsilon_1 H_u H_d \left(H_u^\dagger H_u + H_d^\dagger H_d \right) + \epsilon_2 (H_u H_d)^2 + \text{h.c.} \\ & + \frac{\epsilon_1}{\mu^*} \left[2(H_u H_d)(\tilde{H}_u \tilde{H}_d) + 2(\tilde{H}_u H_d)(H_u \tilde{H}_d) \right. \\ & \left. + (H_u \tilde{H}_d)(H_u \tilde{H}_d) + (\tilde{H}_u H_d)(\tilde{H}_u H_d) \right] + \text{h.c.} \end{aligned}$$

where

$$\epsilon_1 \equiv \frac{\mu^* \lambda_1}{M} \quad \epsilon_2 \equiv -\frac{m_{\text{susy}} \lambda_2}{M}$$

- New contributions for Higgs boson masses
- New contributions for higgsino (χ^0 and χ^\pm) masses
- New contributions for Higgs-higgsino couplings

BMSSM Higgs potential

Corrections to the MSSM Higgs potential

$$\begin{aligned} \delta L = & 2 \epsilon_1 H_u H_d \left(H_u^\dagger H_u + H_d^\dagger H_d \right) + \epsilon_2 (H_u H_d)^2 + \text{h.c.} \\ & + \frac{\epsilon_1}{\mu^*} \left[2(H_u H_d)(\tilde{H}_u \tilde{H}_d) + 2(\tilde{H}_u H_d)(H_u \tilde{H}_d) \right. \\ & \left. + (H_u \tilde{H}_d)(H_u \tilde{H}_d) + (\tilde{H}_u H_d)(\tilde{H}_u H_d) \right] + \text{h.c.} \end{aligned}$$

where

$$\epsilon_1 \equiv \frac{\mu^* \lambda_1}{M} \quad \epsilon_2 \equiv -\frac{m_{\text{susy}} \lambda_2}{M}$$

- New contributions for Higgs boson masses
- New contributions for higgsino (χ^0 and χ^\pm) masses
- New contributions for Higgs-higgsino couplings

Vacuum stability: $|\epsilon_1| \lesssim 0.1$, $|\epsilon_2| \lesssim 0.05$ see Blum, Delaunay, Hochberg, 09

Higgs spectrum

We consider the case where the NR operators can still be treated as **perturbations**:

$$M_h^2 \simeq \left(m_h^{\text{tree}}\right)^2 + \delta_{\tilde{t}} m_h^2 + \delta_{\epsilon} m_h^2 \gtrsim (114 \text{ GeV})^2$$

$$\delta_{\epsilon} m_h^2 = 2v^2 \left(\epsilon_2 - 2\epsilon_1 s_{2\beta} - \frac{2\epsilon_1(m_A^2 + m_Z^2)s_{2\beta} + \epsilon_2(m_A^2 - m_Z^2)c_{2\beta}^2}{\sqrt{(m_A^2 - m_Z^2)^2 + 4m_A^2 m_Z^2 s_{2\beta}^2}} \right)$$

$$\delta_{\epsilon} m_h^2 \sim \text{few dozens of GeVs!}$$

The $\delta_{\epsilon} m_h^2$ relaxes the constraint in a significant way:
for $\epsilon_1 \lesssim -0.1$ and $\tan\beta \lesssim 5$, **light and unmixed stops** allowed!

Higgs spectrum

We consider the case where the NR operators can still be treated as **perturbations**:

$$M_h^2 \simeq \left(m_h^{\text{tree}}\right)^2 + \delta_{\tilde{t}} m_h^2 + \delta_{\epsilon} m_h^2 \gtrsim (114 \text{ GeV})^2$$

$$\delta_{\epsilon} m_h^2 = 2v^2 \left(\epsilon_2 - 2\epsilon_1 s_{2\beta} - \frac{2\epsilon_1(m_A^2 + m_Z^2)s_{2\beta} + \epsilon_2(m_A^2 - m_Z^2)c_{2\beta}^2}{\sqrt{(m_A^2 - m_Z^2)^2 + 4m_A^2 m_Z^2 s_{2\beta}^2}} \right)$$

$$\delta_{\epsilon} m_h^2 \sim \text{few dozens of GeVs!}$$

The $\delta_{\epsilon} m_h^2$ relaxes the constraint in a significant way:
for $\epsilon_1 \lesssim -0.1$ and $\tan\beta \lesssim 5$, **light and unmixed stops** allowed!

→ The SUSY little hierarchy problem can be avoided

Higgs spectrum

We consider the case where the NR operators can still be treated as **perturbations**:

$$M_h^2 \simeq \left(m_h^{\text{tree}}\right)^2 + \delta_{\tilde{t}} m_h^2 + \delta_{\epsilon} m_h^2 \gtrsim (114 \text{ GeV})^2$$

$$\delta_{\epsilon} m_h^2 = 2v^2 \left(\epsilon_2 - 2\epsilon_1 s_{2\beta} - \frac{2\epsilon_1(m_A^2 + m_Z^2)s_{2\beta} + \epsilon_2(m_A^2 - m_Z^2)c_{2\beta}^2}{\sqrt{(m_A^2 - m_Z^2)^2 + 4m_A^2 m_Z^2 s_{2\beta}^2}} \right)$$



$$\delta_{\epsilon} m_h^2 \sim \text{few dozens of GeVs!}$$

The $\delta_{\epsilon} m_h^2$ relaxes the constraint in a significant way:
for $\epsilon_1 \lesssim -0.1$ and $\tan\beta \lesssim 5$, **light and unmixed stops** allowed!

→ The SUSY little hierarchy problem can be avoided

Other Higgs masses also receive corrections...

Higgs spectrum



By Berg, Edsjö, Gondolo, Lundström and Sjörs, 09'

The $\delta_\epsilon m_h^2$ relaxes the constraint in a significant way:
for $\epsilon_1 \lesssim -0.1$ and $\tan\beta \lesssim 5$, **light and unmixed stops** allowed!

➔ The SUSY little hierarchy problem can be avoided

Other Higgs masses also receive corrections...

Higgsinos

$$M_{\chi^0} = \begin{pmatrix} M_1 & 0 & -m_Z s_W c_\beta & m_Z s_W s_\beta \\ 0 & M_2 & m_Z c_W c_\beta & -m_Z c_W s_\beta \\ -m_Z s_W c_\beta & m_Z c_W c_\beta & 0 & -\mu \\ m_Z s_W s_\beta & -m_Z c_W s_\beta & -\mu & 0 \end{pmatrix} + \frac{4\epsilon_1 m_W^2}{\mu^* g^2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & s_\beta^2 & s_{2\beta} \\ 0 & 0 & s_{2\beta} & c_\beta^2 \end{pmatrix}$$

The lightest neutralino χ_1^0 is a natural candidate for **cold dark matter!**

The NR operators also modify

- the chargino mass matrix
- Higgs-higgsino-higgsino & Higgs-Higgs-higgsino-higgsino couplings (DM annihilation cross sections)

Berg, Edsjö, Gondolo, Lundström, Sjörs, '09; NB, Blum, Losada, Nir, '09

Higgsinos

$$M_{\chi^0} = \begin{pmatrix} M_1 & 0 & -m_Z s_W c_\beta & m_Z s_W s_\beta \\ 0 & M_2 & m_Z c_W c_\beta & -m_Z c_W s_\beta \\ -m_Z s_W c_\beta & m_Z c_W c_\beta & 0 & -\mu \\ m_Z s_W s_\beta & -m_Z c_W s_\beta & -\mu & 0 \end{pmatrix} + \frac{4\epsilon_1 m_W^2}{\mu^* g^2} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & s_\beta^2 & s_{2\beta} \\ 0 & 0 & s_{2\beta} & c_\beta^2 \end{pmatrix}$$

The lightest neutralino χ_1^0 is a natural candidate for **cold dark matter!**

The NR operators also modify

- the chargino mass matrix
- Higgs-higgsino-higgsino & Higgs-Higgs-higgsino-higgsino couplings (DM annihilation cross sections)

Berg, Edsjö, Gondolo, Lundström, Sjörs, '09; NB, Blum, Losada, Nir, '09

➔ Spectrum, dark matter relic density and DM detection rates are calculated using modified versions of SuSpect and micrOMEGAS

Outline

- 1 Motivation
- 2 The BMSSM
- 3 Dark Matter**
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 Dark Matter Direct Detection
- 5 Dark Matter Indirect Detection
 - γ -rays
 - Positrons
 - Antiprotons
- 6 Summary

Correlated stop-slepton masses: mSUGRA-like

The mSUGRA model is specified by 5 parameters:


- $\tan\beta$: ratio of the Higgs vevs
- $m_{1/2}$: common mass for the gauginos (bino, wino and gluino)
- m_0 : universal scalar mass (sfermions and Higgs bosons)
- A_0 : universal trilinear coupling
- $\text{sign } \mu$: sign of the μ parameter

In mSUGRA scenarios usually the lightest neutralino is the LSP

Because of the LEP constraint over the Higgs mass, the *bulk region* (i.e. low m_0 and low $m_{1/2}$) is ruled out.


Correlated stop-slepton masses

Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



Correlated stop-slepton masses


Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



- Regions excluded: \tilde{t} LSP

Correlated stop-slepton masses


Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



- Regions excluded: \tilde{t} LSP and χ^\pm searches at LEP

Correlated stop-slepton masses


Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



- Regions excluded: $\tilde{\tau}$ LSP and χ^\pm searches at LEP
- Bulk region: LSP is mainly bino-like. DM relic density too high

Correlated stop-slepton masses


Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



- Regions excluded: $\tilde{\tau}$ LSP and χ^\pm searches at LEP
- Bulk region: LSP is mainly bino-like. DM relic density too high
- Regions fulfilling WMAP measurements:
 - ✓ Coannihilation with $\tilde{\tau}$

Correlated stop-slepton masses


Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



- Regions excluded: $\tilde{\tau}$ LSP and χ^\pm searches at LEP
- Bulk region: LSP is mainly bino-like. DM relic density too high
- Regions fulfilling WMAP measurements:
 - ✓ Coannihilation with $\tilde{\tau}$
 - ✓ Higgs- and Z-poles: $m_h \sim m_Z \sim 2m_\chi$ s-channel exchange

Correlated stop-slepton masses



Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$



- Regions excluded: $\tilde{\tau}$ LSP and χ^\pm searches at LEP
- Bulk region: LSP is mainly bino-like. DM relic density too high
- Regions fulfilling WMAP measurements:
 - ✓ Coannihilation with $\tilde{\tau}$
 - ✓ Higgs- and Z-poles: $m_h \sim m_Z \sim 2m_\chi$ *s*-channel exchange
- ✗ However $m_h \lesssim 105$ GeV: The whole region is excluded!

Correlated stop-slepton masses



Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$ $\epsilon_1 = -0.1, \epsilon_2 = 0$



It should not be taken as an extended mSUGRA,
but **just** as a framework specified at low energy.

Correlated stop-slepton masses

Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$ $\epsilon_1 = -0.1, \epsilon_2 = 0$





It should not be taken as an extended mSUGRA,
but **just** as a framework specified at low energy.

- ✓ Important uplift of the Higgs mass → ‘bulk region’ re-opened

Correlated stop-slepton masses

Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$ $\epsilon_1 = -0.1, \epsilon_2 = 0$





It should not be taken as an extended mSUGRA,
but **just** as a framework specified at low energy.

- ✓ Important uplift of the Higgs mass → ‘bulk region’ re-opened
- New region fulfilling DM constraint: Higgs-funnel

Correlated stop-slepton masses

Let's take: $A_0 = 0$ GeV, $\mu > 0$ and $\tan\beta = 3$ $\epsilon_1 = -0.1, \epsilon_2 = 0$



It should not be taken as an extended mSUGRA,
but **just** as a framework specified at low energy.

- ✓ Important uplift of the Higgs mass \rightarrow 'bulk region' re-opened
 - New region fulfilling DM constraint: Higgs-funnel
 - χ_1^0 bino-like: marginal impact on m_χ and ann. cross section

Light stops, heavy sleptons

Now we consider a low-energy scenario giving rise to light stops

- $\tan\beta$: ratio of the Higgs vevs
- μ : higgsino mass parameter
- m_A : pseudoscalar Higgs mass parameter
- X_t : trilinear coupling for stops, $X_t = A_t - \mu/\tan\beta$
- M_2 : wino mass parameter, $M_1 \sim \frac{1}{2}M_2$
- m_U : stop right mass parameter
- m_Q : 3rd generation squarks left mass parameter
- $m_{\tilde{f}}$: mass for sleptons, 1st and 2nd gen. squarks and \tilde{b}_R
 $m_U = 210 \text{ GeV}$, $X_t = 0 \text{ GeV}$, $m_Q = m_{\tilde{f}} = m_A = 500 \text{ GeV}$

Light stops, heavy sleptons


Now we consider a low-energy scenario giving rise to light stops

- $\tan\beta$: ratio of the Higgs vevs
- μ : higgsino mass parameter
- m_A : pseudoscalar Higgs mass parameter
- X_t : trilinear coupling for stops, $X_t = A_t - \mu/\tan\beta$
- M_2 : wino mass parameter, $M_1 \sim \frac{1}{2}M_2$
- m_U : stop right mass parameter
- m_Q : 3rd generation squarks left mass parameter
- $m_{\tilde{f}}$: mass for sleptons, 1st and 2nd gen. squarks and \tilde{b}_R
 $m_U = 210 \text{ GeV}, \quad X_t = 0 \text{ GeV}, \quad m_Q = m_{\tilde{f}} = m_A = 500 \text{ GeV}$


$$m_{\tilde{t}_1} \lesssim 150 \text{ GeV}, \quad 370 \text{ GeV} \lesssim m_{\tilde{t}_2} \lesssim 400 \text{ GeV}$$

A scenario with light unmixed stops is ruled out in the MSSM

Light stops, heavy sleptons




Light stops, heavy sleptons




- Regions excluded: \tilde{t} LSP

Light stops, heavy sleptons




- Regions excluded: \tilde{t} LSP and χ^\pm searches at LEP

Light stops, heavy sleptons




- Regions excluded: \tilde{t} LSP and χ^\pm searches at LEP
- Regions fulfilling WMAP measurements:
 - ✓ Coannihilation with \tilde{t} : $\chi\tilde{t} \rightarrow Wb, tg$ $\tilde{t}\tilde{t} \rightarrow gg$

Light stops, heavy sleptons




- Regions excluded: \tilde{t} LSP and χ^\pm searches at LEP
- Regions fulfilling WMAP measurements:
 - ✓ Coannihilation with \tilde{t} : $\chi\tilde{t} \rightarrow Wb, tg$ $\tilde{t}\tilde{t} \rightarrow gg$
 - ✓ Higgs- and Z-poles: $m_h \sim m_Z \sim 2m_\chi$ s -channel exchange

Light stops, heavy sleptons




- Regions excluded: \tilde{t} LSP and χ^\pm searches at LEP
- Regions fulfilling WMAP measurements:
 - ✓ Coannihilation with \tilde{t} : $\chi\tilde{t} \rightarrow Wb, tg$ $\tilde{t}\tilde{t} \rightarrow gg$
 - ✓ Higgs- and Z-poles: $m_h \sim m_Z \sim 2m_\chi$ s -channel exchange
- ✗ However $m_h \lesssim 85 \text{ GeV}$: The whole region is excluded!

Light stops, heavy sleptons




✓ important uplift of the Higgs mass: $m_h \sim 122$ GeV

Light stops, heavy sleptons




- ✓ important uplift of the Higgs mass: $m_h \sim 122$ GeV
- ✗ NR operators destabilize scalar potential: vacuum metastable

Light stops, heavy sleptons



- ✓ important uplift of the Higgs mass: $m_h \sim 122$ GeV
- ✗ NR operators destabilize scalar potential: vacuum metastable
- new region fulfilling DM constraint: Higgs-funnel

Light stops, heavy sleptons



- ✓ important uplift of the Higgs mass: $m_h \sim 122$ GeV
- ✗ NR operators destabilize scalar potential: vacuum metastable
- new region fulfilling DM constraint: Higgs-funnel
- sizable impact on m_{χ} and ann. cross section when χ_1^0 is higgsino-like


Outline

- 1 Motivation
- 2 The BMSSM
- 3 Dark Matter
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 Dark Matter Direct Detection**
- 5 Dark Matter Indirect Detection
 - γ -rays
 - Positrons
 - Antiprotons
- 6 Summary

Dark matter direct detection

Direct detection experiments are designed to detect **dark matter particles** by their **elastic collision with target nuclei**, placed in a detector on the Earth.

XENON




Exposures: $\varepsilon = 30, 300, 3000 \text{ kg} \cdot \text{year}$
 Xenon1T and 11 days, 4 months or 3 years

Dark matter direct detection

Direct detection experiments are designed to detect **dark matter particles** by their **elastic collision with target nuclei**, placed in a detector on the Earth.

XENON



Exposures: $\varepsilon = 30, 300, 3000 \text{ kg} \cdot \text{year}$
Xenon1T and 11 days, 4 months or 3 years

Xenon discriminates signal from background by simultaneous measurements of:

- scintillation
- ionization

The collaboration expects to have a negligible background.

→ 7 energy bins between [4, 30] keV


Detectability definition:

$$\chi^2 = \sum_{i=1}^7 \frac{(N_i^{\text{tot}} - N_i^{\text{bkg}})^2}{N_i^{\text{tot}}}$$

Dark matter direct detection

Direct detection experiments are designed to detect **dark matter particles** by their **elastic collision with target nuclei**, placed in a detector on the Earth.

XENON



Exposures: $\varepsilon = 30, 300, 3000 \text{ kg} \cdot \text{year}$
Xenon1T and 11 days, 4 months or 3 years

Recoil rates

$$\frac{dN}{dE_r} = \frac{\sigma_{\chi-p} \cdot \rho_0}{2 M_r^2 m_\chi} F(E_r)^2 \int_{v_{\min}(E_r)}^{v_{\text{esc}}} \frac{f(v)}{v} dv$$

$$\text{Reduced mass } M_r = \frac{m_\chi m_N}{m_\chi + m_N}$$

N : number of scatterings ($\text{s}^{-1} \text{kg}^{-1}$)

E_r : nuclear recoil energy $\sim \text{few keV}$

m_χ : WIMP mass

$\sigma_{\chi-p}$: WIMP-proton scattering cross-section

→ Assume pure **spin-independent** coupling


ρ_0 : local WIMP density 0.38 GeV cm^{-3}

F : nuclear form factor Woods-Saxon

$f(v)$: WIMP local vel. distribution M.B.


$$f(v) = \frac{1}{\sqrt{\pi}} \frac{v}{1.05 v_0^2} \left[e^{-(v-1.05 v_0)^2/v_0^2} - e^{-(v+1.05 v_0)^2/v_0^2} \right]$$

Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$ values
 ($m_0 \rightarrow$ increase squark masses, $m_{1/2} \rightarrow$ increase LSP mass)


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$ values
- For low $m_{1/2}$, LSP tends to be a higgsino-bino mixed state ($C_{\chi\chi h}$)


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$ values
- For low $m_{1/2}$, LSP tends to be a higgsino-bino mixed state ($C_{\chi\chi h}$)
- Detection maximised for low $\tan\beta$, $C_{\chi\chi h} \propto \sin 2\beta$ ($|\mu| \gg M_1$)


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$ values
- For low $m_{1/2}$, LSP tends to be a higgsino-bino mixed state ($C_{\chi\chi h}$)
- Detection maximised for low $\tan\beta$, $C_{\chi\chi h} \propto \sin 2\beta$ ($|\mu| \gg M_1$)
- ✓ Sizable amount of the parameter space can be probed


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL


- Detection prospects maximised for low m_0 and $m_{1/2}$ values
- For low $m_{1/2}$, LSP tends to be a higgsino-bino mixed state ($C_{\chi\chi h}$)
- Detection maximised for low $\tan\beta$, $C_{\chi\chi h} \propto \sin 2\beta$ ($|\mu| \gg M_1$)
- ✓ Sizable amount of the parameter space can be probed
- ➔ NR operators \rightarrow deterioration of the detection: m_h
- ✓ But without NR operators, the parameter space was excluded!

Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

✘ Partially ruled out by first results from Xenon100!


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- ✘ Partially ruled out by first results from Xenon100!
- Detection prospects maximised for low μ and/or M_1 : light LSP


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- ✘ Partially ruled out by first results from Xenon100!
 - Detection prospects maximised for low μ and/or M_1 : light LSP
 - Scattering cross section enhanced near $\mu \sim M_1$ (C_{XXh}, C_{XXH})


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- ✘ Partially ruled out by first results from Xenon100!
 - Detection prospects maximised for low μ and/or M_1 : light LSP
 - Scattering cross section enhanced near $\mu \sim M_1$ (C_{XXh}, C_{XXH})
 - Neither Z- nor h -funnel enhance SI direct detection
 - Spin-dependent detection sensible to the Z-peak (non-universality)


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- ✘ Partially ruled out by first results from Xenon100!
- Detection prospects maximised for low μ and/or M_1 : light LSP
- Scattering cross section enhanced near $\mu \sim M_1$ (C_{XXh}, C_{XXH})
- Neither Z- nor h -funnel enhance SI direct detection
- ➔ NR operators deteriorates DD: increase m_h and suppression C_{XXh}

Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- ✘ Partially ruled out by first results from Xenon100!
 - Detection prospects maximised for low μ and/or M_1 : light LSP
 - Scattering cross section enhanced near $\mu \sim M_1$ (C_{XXh} , C_{XXH})
 - Neither Z- nor h -funnel enhance SI direct detection
- ➔ NR operators deteriorates DD: increase m_h and suppression C_{XXh}
- ✓ BMSSM satisfies all DD measurements!

Outline

- 1 Motivation
- 2 The BMSSM
- 3 Dark Matter
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 Dark Matter Direct Detection
- 5 Dark Matter Indirect Detection**
 - γ -rays
 - Positrons
 - Antiprotons
- 6 Summary

Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify

Gamma-rays generated in

DM annihilation in the galactic center

$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$




Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify

Gamma-rays generated in

DM annihilation in the galactic center

$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$




Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify

Gamma-rays generated in

DM annihilation in the galactic center



$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$



Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify
Gamma-rays generated in
DM annihilation in the galactic center

$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$





Fermi telescope (Launched '08)

Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify **Gamma-rays** generated in **DM annihilation** in the galactic center

$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$



Fermi telescope (Launched '08)

Differential event rate

$$\Phi_{\gamma}(E_{\gamma}, \psi) = \sum_i \frac{dN_{\gamma}^i}{dE_{\gamma}} \langle \sigma_i v \rangle \frac{1}{8\pi m_{\chi}^2} \int_{los} \rho(r)^2 dl$$

$\frac{dN}{dE}$: spectrum of secondary particles

E_{γ} : gamma energy

$\langle \sigma v \rangle$: averaged annihilation cross-section by velocity

$\rho(r)$: dark matter halo profile

5-years data acquisition, $\Delta\Omega = 3 \cdot 10^{-5}$ sr



Background: HESS measurements

(Diffuse Galactic emission and Sagittarius A*)

Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify **Gamma-rays** generated in **DM annihilation** in the galactic center

$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$



Fermi telescope (Launched '08)

Differential event rate


$$\Phi_\gamma(E_\gamma, \psi) = \sum_i \frac{dN_\gamma^i}{dE_\gamma} \langle \sigma_i v \rangle \frac{1}{8\pi m_\chi^2} \int_{los} \rho(r)^2 dl$$

$\frac{dN}{dE}$: spectrum of secondary particles

E_γ : gamma energy

$\langle \sigma v \rangle$: averaged annihilation cross-section by velocity


$\rho(r)$: dark matter halo profile



Dark matter indirect detection (γ -rays)

We study the ability of **Fermi** to identify **Gamma-rays** generated in **DM annihilation** in the galactic center

$$\chi\bar{\chi} \rightarrow b\bar{b}, WW \dots \rightarrow \gamma + \dots$$



Fermi telescope (Launched '08)

Differential event rate


$$\Phi_{\gamma}(E_{\gamma}, \psi) = \sum_i \frac{dN_{\gamma}^i}{dE_{\gamma}} \langle \sigma_i v \rangle \frac{1}{8\pi m_{\chi}^2} \int_{los} \rho(r)^2 dl$$

$\frac{dN}{dE}$: spectrum of secondary particles

E_{γ} : gamma energy


$\langle \sigma v \rangle$: averaged annihilation cross-section by velocity

$\rho(r)$: dark matter halo profile




3 halo profiles: Einasto, NFW and NFW_c (adiabatic compression due to baryons)

Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$
- Thresholds: $\chi\chi \rightarrow W^+W^-$, $\chi\chi \rightarrow t\bar{t}$


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$
- Thresholds: $\chi\chi \rightarrow W^+W^-$, $\chi\chi \rightarrow t\bar{t}$
- Detection maximised for high $\tan\beta$ $\chi\chi \rightarrow b\bar{b}$ and $\tau\tau \propto \tan\beta$ and $1/\cos\beta$


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$
- Thresholds: $\chi\chi \rightarrow W^+W^-$, $\chi\chi \rightarrow t\bar{t}$
- Detection maximised for high $\tan\beta$ $\chi\chi \rightarrow b\bar{b}$ and $\tau\tau \propto \tan\beta$ and $1/\cos\beta$
- For large $\tan\beta$ thresholds weaken


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL

- Detection prospects maximised for low m_0 and $m_{1/2}$
- Thresholds: $\chi\chi \rightarrow W^+W^-$, $\chi\chi \rightarrow t\bar{t}$
- Detection maximised for high $\tan\beta$ $\chi\chi \rightarrow b\bar{b}$ and $\tau\tau \propto \tan\beta$ and $1/\cos\beta$
- For large $\tan\beta$ thresholds weaken
- Only scenarios with highly cusped inner regions could be probed


Correlated stop-slepton masses



Exclusion lines: ability to test and exclude at 95% CL


- Detection prospects maximised for low m_0 and $m_{1/2}$
- Thresholds: $\chi\chi \rightarrow W^+W^-$, $\chi\chi \rightarrow t\bar{t}$
- Detection maximised for high $\tan \beta$ $\chi\chi \rightarrow b\bar{b}$ and $\tau\tau \propto \tan \beta$ and $1/\cos \beta$
- For large $\tan \beta$ thresholds weaken
- Only scenarios with highly cusped inner regions could be probed
- NR operators: Higgs pole ‘invisible’ ($v \rightarrow 0$)

Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- Detection enhanced for $M_1 \gg \mu$ ($\chi\chi Z$ and $\chi\chi^\pm W^\mp$ couplings)


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- Detection enhanced for $M_1 \gg \mu$ ($\chi\chi Z$ and $\chi\chi^\pm W^\mp$ couplings)
- $\langle\sigma v\rangle$ enhanced for high $\tan\beta$ ($\chi\chi \rightarrow b\bar{b}, WW$)


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- Detection enhanced for $M_1 \gg \mu$ ($\chi\chi Z$ and $\chi\chi^\pm W^\mp$ couplings)
- $\langle\sigma v\rangle$ enhanced for high $\tan\beta$ ($\chi\chi \rightarrow b\bar{b}, WW$)
- h -funnel could not be tested (no s -wave contribution)


Light stops, heavy sleptons



Exclusion lines: ability to test and exclude at 95% CL

- Detection enhanced for $M_1 \gg \mu$ ($\chi\chi Z$ and $\chi\chi^\pm W^\mp$ couplings)
- $\langle\sigma v\rangle$ enhanced for high $\tan\beta$ ($\chi\chi \rightarrow b\bar{b}, WW$)
- h -funnel could not be tested (no s -wave contribution)
- NFW and Einasto could test some regions, but not relevant

Antimatter (e^+ and \bar{p}) propagation



picture provided by M. Cirelli

→ Diffusion equation solved in the Diffusive zone

Baltz & Edsjö '98; Lavalley, Pochon, Salati & Taillet '06

$$\frac{\partial f}{\partial t} = K(E) \nabla^2 f + Q_{\text{inj}} + \frac{\partial}{\partial E} [b(E)f] - 2h \delta(z) \Gamma_{\text{ann}} f - \frac{\partial}{\partial z} [V_c f]$$

diffusion



source

energy loss



spallation

convective wind

Light stops, heavy sleptons - Positrons




Light stops, heavy sleptons - Positrons




- Perspectives for the oncoming AMS-02 satellite background: Fermi & PAMELA measurements. PAMELA's 'heritage': A quite large background that is difficult to overcome.

Light stops, heavy sleptons - Positrons





- Perspectives for the oncoming AMS-02 satellite background: Fermi & PAMELA measurements. PAMELA's 'heritage': A quite large background that is difficult to overcome.
- ✗ PAMELA excess buries all signals

Light stops, heavy sleptons - Positrons





- Perspectives for the oncoming AMS-02 satellite background: Fermi & PAMELA measurements. PAMELA's 'heritage': A quite large background that is difficult to overcome.
- ✗ PAMELA excess buries all signals
 - Some small hope in the region where the LSP carries a significant higgsino component, due to the rise in the coupling with Z's

Light stops, heavy sleptons - Antiprotons





Light stops, heavy sleptons - Antiprotons





- Perspectives for the oncoming AMS-02 satellite
 background: PAMELA measurements (It seem to confirm the background predicted)

Light stops, heavy sleptons - Antiprotons



- Perspectives for the oncoming AMS-02 satellite background: PAMELA measurements (It seem to confirm the background predicted)
 - The background is not very high, but the signal is quite low!

Light stops, heavy sleptons - Antiprotons



- Perspectives for the oncoming AMS-02 satellite background: PAMELA measurements (It seem to confirm the background predicted)
- The background is not very high, but the signal is quite low!
 - Much better than positrons!

Outline

- 1 Motivation
- 2 The BMSSM
- 3 Dark Matter
 - Correlated stop-slepton masses
 - Light stops, heavy sleptons
- 4 Dark Matter Direct Detection
- 5 Dark Matter Indirect Detection
 - γ -rays
 - Positrons
 - Antiprotons
- 6 Summary

Conclusions and prospects

- NR operators in the Higgs sector introduced for reducing fine-tuning (Little hierarchy)
- Bulk region re-opened
- Possible to have light unmixed stops
- New regions fulfilling the DM constraint:
 - Higgs-pole
 - Higgs-stop coannihilation
- EW baryogenesis opens up
- Both scenarios could be tested by present machines!
- Complementarity with different detection modes

Antimatter propagation

$$\frac{\partial f}{\partial t} = K(E) \nabla^2 f$$

→ Diffusion equation

$$K(E) = K_0 E_{\text{GeV}}^\alpha \quad \text{Diffusion coefficient}$$

Propagation parameters K_0 and α fixed by N-body simulations

Antimatter propagation

$$\frac{\partial f}{\partial t} = K(E) \nabla^2 f + Q_{\text{inj}}$$

→ Source term due to DM DM annihilation

$$Q_{\text{inj}} = \frac{1}{2} \left(\frac{\rho(r)}{m_\chi} \right)^2 \sum_k \langle \sigma v \rangle_k \frac{dN_k}{dE}$$

Antimatter propagation

$$\frac{\partial f}{\partial t} = K(E) \nabla^2 f + Q_{\text{inj}} + \frac{\partial}{\partial E} [b(E)f]$$

→ Energy loss term

$$b(E) = \frac{E_{\text{GeV}}^2}{\tau_E} \quad \text{Energy loss rate}$$

For antiprotons energy losses can be ignored

Antimatter propagation

$$\frac{\partial f}{\partial t} = K(E) \nabla^2 f + Q_{\text{inj}} + \frac{\partial}{\partial E} [b(E)f] - 2h \delta(z) \Gamma_{\text{ann}} f$$

- Annihilation of \bar{p} on interstellar protons in the galactic plane (Spallation)

$$\Gamma_{\text{ann}} = \left(n_H + 4^{2/3} n_{He} \right) \sigma_{\text{ann}}^{p\bar{p}} v_{\bar{p}} \quad \text{Annihilation rate}$$

Annihilation only relevant for antiprotons


Antimatter propagation

$$\frac{\partial f}{\partial t} = K(E) \nabla^2 f + Q_{\text{inj}} + \frac{\partial}{\partial E} [b(E)f] - 2h \delta(z) \Gamma_{\text{ann}} f$$

→ Final Diffusion equation


Semi-analytical 2D diffusion equation

Baltz & Edsjö '98; Lavallo, Pochon, Salati & Taillet '06



picture snatched to M. Cirelli

Positrons from PAMELA



- Steep e^+ excess above 10 GeV
- Very large flux