

Conference “Strong Coupling: from Lattice to AdS/CFT”
Firenze, 3-5 June 2008

*Thermodynamics of the
Einstein-dilaton system
and Improved Holographic QCD*

Elias Kiritsis

Ecole Polytechnique
and
University of Crete

Bibliography

- R. Casero, E. Kiritsis and A. Paredes, “*Chiral symmetry breaking as open string tachyon condensation,*” Nucl. Phys. B **787** (2007) 98; [arXiv:hep-th/0702155].
- U. Gursoy and E. Kiritsis, “*Exploring improved holographic theories for QCD: Part I,*” JHEP **0802** (2008) 032 [ArXiv:0707.1324][hep-th].
- U. Gursoy, E. Kiritsis and F. Nitti, “*Exploring improved holographic theories for QCD: Part II,*” JHEP **0802** (2008) 019 [ArXiv:0707.1349][hep-th].
- U. Gursoy, E. Kiritsis, L. Mazzanti and F. Nitti, “*Deconfinement and Gluon Plasma Dynamics in Improved Holographic QCD,*” [ArXiv:0804.0899][hep-th].
- Ongoing work with:
C. Charmousis, U. Gursoy, L. Mazzanti, G. Michalogiorgakis, F. Nitti

Introduction

- AdS/CFT provides a surprising twist to large- N gauge theories: the existence of extra dimensions, including the radial holographic dimension.
- It has provided a dual description of strongly-coupled gauge theories translating their physics into string/gravitational dynamics.
- The best studied/controlled example is a maximally supersymmetric and conformal gauge theory whose string dual is described in terms of critical string theory (in ten-dimensions).
- There have been further 10d solutions that broke susy and produced gravitational duals to theories that in IR involve only gluon dynamics. They all however have KK modes at the same scale as Λ , which so far we have been unable to decouple.
- Alternative attempts have focused in noncritical string theory, where there are no KK Modes. Here however one has to address the strong curvature problem that is generic. Progress has been sporadic in this direction.

AdS/QCD

♠ A basic phenomenological approach: use a slice of AdS_5 , with a UV cutoff, and an IR cutoff. *Polchinski+Strassler*

♠ It successfully exhibits confinement (trivially via IR cutoff), and power-like behavior in hard scattering amplitudes

♠ It may be equipped with a bifundamental scalar, T , and $U(N_f)_L \times U(N_f)_R$, gauge fields to describe mesons. *Erlich+Katz+Son+Stepanov, DaRold+Pomarol*

Chiral symmetry is broken by hand, via IR boundary conditions. The low-lying meson spectrum looks "reasonable".

♠ Shortcomings:

- The glueball spectrum does not fit very well the lattice calculations. It has the wrong asymptotic behavior $m_n^2 \sim n^2$ at large n .
- Magnetic quarks are confined instead of screened.
- Chiral symmetry breaking is input by hand.
- The meson spectrum has also the wrong UV asymptotics $m_n^2 \sim n^2$.

♠ The asymptotic spectrum can be fixed by introducing a **non-dynamical** dilaton profile $\Phi \sim r^2$ (soft wall) *Karch+Katz+Son+Stephanov*

Improving AdS/QCD

- ♠ We will use input from both string theory and the gauge theory (QCD) in order to provide an improved phenomenological holographic model for real world QCD.
- ♠ This is an exploratory adventure, and we will short-circuit several obstacles on the way.
- ♠ As we will see, we will get an interesting perspective on the physics of pure glue as well as on the quark sector.
- ♠ The model that will be advocated will be a form of dilaton gravity in 5 dimensions supplemented with space filling flavour branes.
- ♠ We will analyze the finite temperature dynamics that will be compared to that of QCD.

A string theory for QCD: basic expectations

- Pure $SU(N_c)$ $d=4$ YM is expected to be dual to a string theory in 5 dimensions only. Essentially a single adjoint field \rightarrow a single extra dimension.
- The theory becomes asymptotically free and conformal at high energy \rightarrow we expect the classical saddle point solution to asymptote to AdS_5 .
- ♠ operators with lowest dimension are expected to be the only (important) non-trivial bulk fields in the large- N_c saddle-point
- scalar YM operators with $\Delta_{UV} > 4 \rightarrow m^2 > 0$ fields near the AdS_5 boundary \rightarrow vanish fast in the UV regime and do not affect correlators of low-dimension operators.
- Their dimension typically grows large in the IR. Large 't Hooft coupling is expected to suppress the growth in the IR
- This is compatible with the success of low-energy SVZ sum rules as compared to data.
- It is prohibitively difficult otherwise
- ♠ Therefore we will consider $T_{\mu\nu} \leftrightarrow g_{\mu\nu}$, $tr[F^2] \leftrightarrow \phi$, $tr[F \wedge F] \leftrightarrow a$

The nature of the string

- Large- N arguments about the axion (dual to the gauge theory θ -angle) indicate that it must be a RR field.
- The string theory must have no on-shell fermionic states at all because there are no gauge invariant fermionic operators in pure YM.
- Therefore the string theory must be a 5d-superstring theory resembling the II-0 class.
- ♠ Another RR field we expect to have is the RR 4-form, as it is necessary to “seed” the D_3 branes responsible for the gauge group.
- It is non-propagating in 5D
- It seems to be however responsible for the non-trivial IR structure of the gauge theory vacuum.

The effective action, I

- as $N_c \rightarrow \infty$, only string tree-level is dominant.
- Relevant field for the vacuum solution: $g_{\mu\nu}, a, \phi, F_5$.
- The vev of $F_5 \sim N_c \epsilon_5$. It appears always in the combination $e^{2\phi} F_5^2 \sim \lambda^2$, with $\lambda \sim N_c e^\phi$. All higher derivative corrections $(e^{2\phi} F_5^2)^n$ are $\mathcal{O}(1)$ at large N_c .
- This is not the case for all other RR fields: in particular for the axion as $a \sim \mathcal{O}(1)$

$$(\partial a)^2 \sim \mathcal{O}(1) \quad , \quad e^{2\phi} (\partial a)^4 = \frac{\lambda^2}{N_c^2} (\partial a)^4 \sim \mathcal{O}(N_c^{-2})$$

Therefore to leading order $\mathcal{O}(N_c^2)$ we can neglect the axion.

The UV regime

- In the far UV, the space should asymptote to AdS_5 .
- The 't Hooft coupling should behave as ($r \rightarrow 0$)

$$\lambda \sim \frac{1}{\log(r\Lambda)} + \dots \rightarrow 0$$

- Therefore, as $r \rightarrow 0$

$$\text{Curvature} \rightarrow \text{finite} \quad , \quad (\partial\phi)^2 \sim \frac{(\partial\lambda)^2}{\lambda^2} \sim \frac{1}{\log^2(r\Lambda)} \rightarrow 0 \quad , \quad \lambda^2 \rightarrow 0$$

- For $\lambda \rightarrow 0$ the potential in the Einstein frame starts as $V(\lambda) \sim \lambda^{\frac{4}{3}}$ and cannot support the asymptotic AdS_5 solution.
- Therefore asymptotic AdS_5 must arise from curvature corrections

$$S_{eff} \sim \int d^5x \frac{1}{\lambda^2} Z(R, 0, 0)$$

- Setting $\lambda = 0$ at leading order we can generically get an AdS_5 solution coming from balancing the higher curvature corrections.

- There is a "good" (but hard to derive the coefficients) perturbative expansion around this asymptotic AdS_5 solution by perturbing around it:

$$e^A = \frac{\ell}{r} [1 + \delta A] \quad , \quad \lambda = \frac{1}{b_0 \log(r\Lambda)} + \dots$$

- This turns out to be a regular expansion of the solutions in powers of

$$\frac{P_n(\log \log(r\Lambda))}{(\log(r\Lambda))^{-n}}$$

- Effectively this can be rearranged as a "perturbative" expansion in $\lambda(r)$.

- Using λ as a radial coordinate the solution for the metric can be written

$$E \equiv e^A = \frac{\ell}{r(\lambda)} [1 + c_1 \lambda + c_2 \lambda^2 + \dots] = \ell (e^{-\frac{b_0}{\lambda}}) [1 + c'_1 \lambda + c'_2 \lambda^2 + \dots] \quad , \quad \lambda \rightarrow 0$$

The IR regime

- Here the situation is a bit more obscure. The constraints/input are: confinement and mass gap.
- We do expect that $\lambda \rightarrow \infty$ at the IR bottom.
- This is a "singularity" in the conventional sense: it must be "repulsive", ie the string theory, and even better some effective field theory should not break down there.
- (Very) naive intuition from $N=4$ and other 10d strongly coupled theories suggests that in this regime there should be a two derivative description of the physics.
- Similar intuition is coming from the linear dilaton solution that suggests that the (string frame) curvature vanishes at the IR bottom.
- At the IR bottom the space must end (singularity) where the scale factor vanishes.
- ♠ If it happens very slowly, we loose confinement
- ♠ if it happens very fast, the singularity is strong and the theory is incomplete (boundary conditions are needed at the singularity).

Improved Holographic QCD: a model

The simplification in this model relies on writing down a two-derivative action

$$S_{\text{Einstein}} = M^3 N_c^2 \int d^5x \sqrt{g} \left[R - \frac{4(\partial\lambda)^2}{3\lambda^2} + V(\lambda) \right]$$

with

$$\lim_{\lambda \rightarrow 0} V(\lambda) = \frac{12}{\ell^2} \left(1 + \sum_{n=1}^{\infty} c_n \lambda^n \right), \quad \lim_{\lambda \rightarrow \infty} V(\lambda) = \lambda^{\frac{4}{3}} \sqrt{\log \lambda} + \text{subleading}$$

The small λ asymptotics “simulate” the UV expansion around AdS_5 .

- There is a 1-1 correspondence between the YM β -function, $\beta(\lambda)$ and W :

$$\left(\frac{3}{4}\right)^3 V(\lambda) = W^2 - \left(\frac{3}{4}\right)^2 \left(\frac{\partial W}{\partial \log \lambda}\right)^2, \quad \beta(\lambda) = -\frac{9}{4} \lambda^2 \frac{d \log W(\lambda)}{d\lambda}$$

once a choice of energy is made (here $E = A_E$). Renormalization and other choices modify $\beta(\lambda)$ beyond two-loop level

There are some shortcomings localized at the UV

- Conformal anomaly is incorrect.
- Shear viscosity ratio is constant and equal to that of N=1 sYM

both of the above need Riemann curvature corrections.

Many other observables are coming out very well both at T=0 and finite T

♠ The axion contribution

$$\delta S = M_p^3 \int d^5x \sqrt{g} Z(\lambda) (\partial a)^2$$

with

$$\lim_{\lambda \rightarrow 0} Z(\lambda) = c_0 + c_1 \lambda + c_2 \lambda^2 + \dots, \quad \lim_{\lambda \rightarrow \infty} Z(\lambda) = C_\infty \lambda^4 + \dots$$

$$a(r) = \theta_{UV} \frac{\int_r^\infty \frac{dr}{e^{3AZ}}}{\int_0^\infty \frac{dr}{e^{3AZ}}}$$

Quarks ($N_f \ll N_c$) and mesons

- Flavor is introduced by N_f $D_4 + \bar{D}_4$ branes pairs inside the bulk background. Their back-reaction on the bulk geometry is suppressed by N_f/N_c .
- The important world-volume fields are

$$T_{ij} \leftrightarrow \bar{q}_a^i \frac{1 + \gamma^5}{2} q_a^j \quad , \quad A_{\mu}^{ijL,R} \leftrightarrow \bar{q}_a^i \frac{1 \pm \gamma^5}{2} \gamma^{\mu} q_a^j$$

Generating the $U(N_f)_L \times U(N_f)_R$ chiral symmetry.

- The UV mass matrix m_{ij} corresponds to the source term of the Tachyon field. It breaks the chiral (gauge) symmetry. The normalizable mode corresponds to the vev $\langle \bar{q}_a^i \frac{1 + \gamma^5}{2} q_a^j \rangle$.
- We show that the expectation value of the tachyon is non-zero and $T \sim 1$, breaking chiral symmetry $SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)_V$. The anomaly plays an important role in this (holographic Coleman-Witten)

- The fact that the tachyon diverges in the IR (fusing D with \bar{D}) constraints the UV asymptotics and determines the quark condensate $\langle \bar{q}q \rangle$ in terms of m_q . A GOR relation is satisfied (for an asymptotic AdS_5 space)

$$m_\pi^2 = -2 \frac{m_q}{f_\pi^2} \langle \bar{q}q \rangle \quad , \quad m_q \rightarrow 0$$

- We can derive formulae for the anomalous divergences of flavor currents, when they are coupled to an external source.
- When $m_q = 0$, the meson spectrum contains N_f^2 massless pseudoscalars, the $U(N_f)_A$ Goldstone bosons.
- The WZ part of the flavor brane action gives the Adler-Bell-Jackiw $U(1)_A$ axial anomaly and an associated Stueckelberg mechanism gives an $O\left(\frac{N_f}{N_c}\right)$ mass to the would-be Goldstone boson η' , in accordance with the Veneziano-Witten formula.
- Studying the spectrum of highly excited mesons, we find the expected property of linear confinement: $m_n^2 \sim n$.
- The detailed spectrum of mesons remains to be worked out

Concrete potential

- The superpotential chosen is

$$W = (3 + 2b_0\lambda)^{2/3} \left[18 + (2b_0^2 + 3b_1) \log(1 + \lambda^2) \right]^{4/3},$$


with corresponding potential

$$\beta(\lambda) = -\frac{3b_0\lambda^2}{3 + 2b_0\lambda} - \frac{6(2b_0^2 + 3b_1)\lambda^3}{(1 + \lambda^2) \left(18 + (2b_0^2 + 3b_1) \log(1 + \lambda^2) \right)}$$


which is everywhere regular and has the correct UV and IR asymptotics.

- b_0 is a free parameter and b_1/b_0^2 is taken from the QCD β -function

Linearity of the glueball spectrum



(a)




(b)

(a) Linear pattern in the spectrum for the first 40 0^{++} glueball states. M^2 is shown units of $0.015\ell^{-2}$.


(b) The first 8 0^{++} (squares) and the 2^{++} (triangles) glueballs. These spectra are obtained in the background I with $b_0 = 4.2, \lambda_0 = 0.05$.

Comparison with lattice data (Meyer)



Comparison of glueball spectra from our model with $b_0 = 4.2$, $\lambda_0 = 0.05$ (boxes), with the lattice QCD data from Ref. I (crosses) and the AdS/QCD computation (diamonds), for (a) 0^{++} glueballs; (b) 2^{++} glueballs. The masses are in MeV, and the scale is normalized to match the lowest 0^{++} state from Ref. I.

$$l_{eff}^2 = 6.88 l^2$$



The string frame scale factor in background I with $b_0 = 4.2$, $\lambda_0 = 0.05$.

We can “measure”

$$\frac{\ell}{\ell_s} \simeq 6.26 \quad , \quad \ell_s^2 R \simeq -0.5 \quad (1)$$

and predict

$$\alpha_s(1.2\text{GeV}) = 0.34,$$

which is within the error of the quoted experimental value $\alpha_s^{(exp)}(1.2\text{GeV}) = 0.35 \pm 0.01$

The fit to Meyer lattice data

J^{PC}	Ref I (MeV)	Our model (MeV)	Mismatch	$N_c \rightarrow \infty$ [?]	Mismatch
0^{++}	1475 (4%)	1475	0	1475	0
2^{++}	2150 (5%)	2055	4%	2153 (10%)	5%
0^{-+}	2250 (4%)	2243	0		
0^{++*}	2755 (4%)	2753	0	2814 (12%)	2%
2^{++*}	2880 (5%)	2991	4%		
0^{-+*}	3370 (4%)	3288	2%		
0^{++**}	3370 (4%)	3561	5%		
0^{++***}	3990 (5%)	4253	6%		

Comparison between the glueball spectra in Ref. I and in our model. The states we use as input in our fit are marked in red. The parenthesis in the lattice data indicate the percent accuracy.

YM at finite temperature

The theory at finite temperature can be described by:



(1) The “thermal vacuum solution”. This is the zero temperature solution we described so far with time periodically identified with period β .

(2) The “black-hole solution”

$$ds^2 = b(r)^2 \left[\frac{dr^2}{f(r)} - f(r)dt^2 + dx^i dx^i \right], \quad \Phi = \Phi(r)$$

We can show the following:

- For $T > T_{\min}$ there are two black-hole solutions with the same temperature but different horizon positions. One is a “large” BH the other is “small”.
- When $T < T_{\min}$ only the “thermal vacuum solution” exists: it describes the confined phase at finite temperature.
- When $T > T_{\min}$ three competing solutions exist. The large BH has the lowest free energy for $T > T_c > T_{\min}$. It describes the deconfined QGP phase.



- All solutions have two parameters: T and Λ .

For the Black hole solution we can calculate the temperature as

$$\frac{1}{4\pi T} = b_T^3(r_h) \int_0^{r_h} \frac{du}{b_T(u)^3}$$

- The free energy is calculated as a boundary term for both the black-holes and the thermal vacuum solution. They are all UV divergent but their differences are finite. We find

$$\mathcal{F} = (M^3 V_3 N_c^2) \left[6b_T(\epsilon) \sqrt{f(\epsilon)} \left[b_T(\epsilon) \dot{b}_T(\epsilon) \sqrt{f(\epsilon)} - b_0(\epsilon) \dot{b}_0(\epsilon) \right] + f(\epsilon) b_T^3(\epsilon) \right]$$

with

$$f(\epsilon) \simeq 1 - \pi T b_T^3(r_h) \frac{\epsilon^4}{\ell^3} \left[1 + \mathcal{O} \left(\frac{1}{\log(\epsilon\Lambda)} \right) \right] + \dots, \quad b_T(\epsilon) - b_0(\epsilon) = \mathcal{C}(T) \epsilon^3 + \dots$$

The rules of AdS/CFT relate $\mathcal{C}(T)$ to the gluon condensate:

$$\mathcal{C}(T) \propto \langle \text{Tr}[F^2] \rangle_T - \langle \text{Tr}[F^2] \rangle_0$$

The free energy difference is therefore given by

$$\frac{\mathcal{F}}{M_p^3 N_c^2 V_3} = 12 \frac{\mathcal{C}(T)}{\ell} - \pi T b^3(r_h) = 12 \frac{\mathcal{C}(T)}{\ell} - \frac{TS}{4M_p^3 N_c^2 V_3},$$

- The existence of the non-trivial deconfinement transition is due to the non-zero condensate $\mathcal{C}(T)$.

- For the YM potential the minimum temperature for the black-holes is $T_{\min} \simeq 210 \text{ MeV}$ with $\lambda_h \simeq 12$. The critical temperature is

$$T_c \simeq 235 \pm 15 \text{ MeV} \quad \text{with} \quad \lambda_h \simeq 8 \quad , \quad \frac{L_h^{\frac{1}{4}}}{T_c} = 0.65\sqrt{N_c}$$

to be compared with $260 \pm 11 \text{ MeV}$ and $0.77\sqrt{N_c}$
Lucini+Teper, Lucini+Teper+Wenger

- The specific heat for the QGP solution is positive as it should. For the small black-hole it is negative.

- In the QGP phase, the $q\bar{q}$ potential is screened.

- Matching the $T \rightarrow \infty$ regime to an ideal gas of free gluons we obtain

$$(M_p \ell)^3 = \frac{1}{45\pi^2} \quad , \quad M_{\text{physical}} = M_p N_c^{\frac{2}{3}} = \left(\frac{8}{45\pi^2 \ell^3} \right)^{\frac{1}{3}} \simeq 4.6 \text{ GeV}$$

- In the QGP phase, the axion is constant and $\langle F \wedge F \rangle$ vanishes.

General phase structure

• For a general potential we can prove the following (under mild assumptions):


i. There exists a phase transition at finite T , if and only if the zero- T theory confines.

ii. This transition is of the first order for all of the confining geometries, with a single exception described in iii:

iii. In the limit confining geometry $b_0(r) \rightarrow \exp(-Cr)$ (as $r \rightarrow \infty$), the phase transition is of the second order and happens at $T = 3C/4\pi$. This is the linear dilaton vacuum solution in the IR.


iv. All of the non-confining geometries at zero T are always in the black hole phase at finite T . They exhibit a second order phase transition at $T = 0^+$.

Temperature versus horizon position




We plot the relation $T(r_h)$ for various potentials parameterized by a . $a = 1$ is the critical value below which there is only one branch of black-hole solutions.

Free energy versus horizon position




We plot the relation $\mathcal{F}(r_h)$ for various potentials parameterized by a . $a = 1$ is the critical value below which there is no first order phase transition .


The transition in the free energy




Equation of state



The speed of sound



The specific heat



The bulk viscosity

It is defined from the Kubo formula

$$\zeta = \frac{1}{9} \lim_{\omega \rightarrow 0} \frac{1}{\omega} \text{Im} G_R(\omega) \quad , \quad G_R(\omega) \equiv \int d^3x \int dt e^{i\omega t} \theta(t) \langle 0 | [T_{ii}(\vec{x}, t), T_{ii}(\vec{0}, 0)] | 0 \rangle$$

Using a parametrization $ds^2 = e^{2A}(f dt^2 + d\vec{x}^2) + \frac{e^{2B}}{f} dr^2$ in a special gauge $\phi = r$ the relevant metric perturbation decouples

Gubser+Nellore+Pufu+Rocha

$$h''_{11} = - \left(-\frac{1}{3A'} - 4A' + 3B' - \frac{f'}{f} \right) h'_{11} + \left(-\frac{e^{2B-2A}}{f^2} \omega^2 + \frac{f'}{6fA'} - \frac{f'}{f} B' \right) h_{11}$$

with


$$h_{11}(0) = 1 \quad , \quad h_{11}(r_h) \simeq C e^{i\omega t} \left| \log \frac{\lambda}{\lambda_h} \right|^{-\frac{i\omega}{4\pi T}}$$

The correlator is given by the conserved number of h-quanta

$$\text{Im} G_R(\omega) = -4M^3 \mathcal{G}(\omega) \quad , \quad \mathcal{G}(\omega) = \frac{e^{4A-B} f}{4A'^2} |\text{Im}[h_{11}^* h'_{11}]|$$

finally giving

$$\frac{\zeta}{s} = \frac{C^2}{4\pi} \frac{V'(\lambda_h)^2}{V(\lambda_h)^2}$$



Open problems

- Tune the dilaton potential
- Re-Calculate quantities relevant for heavy ion collisions: jet quenching, quark energy loss etc.
- Calculate the finite temperature Polyakov loops in various symmetry channels.
- Investigate quantitatively the meson sector
- Investigate the θ dependence of the meson sector.
- Calculate the phase diagram in the presence of baryon number.

The low dimension spectrum

- What are all gauge invariant YM operators of dimension 4 or less?
 - They are given by $Tr[F_{\mu\nu}F_{\rho\sigma}]$.
- Decomposing into U(4) reps:

$$(\square \otimes \square)_{\text{symmetric}} = \square \oplus \square \quad (2)$$

We must remove traces to construct the irreducible representations of O(4):

$$\square = \square \oplus \square \oplus \bullet, \quad \square = \bullet$$

The two singlets are the scalar (dilaton) and pseudoscalar (axion)

$$\phi \leftrightarrow Tr[F^2], \quad a \leftrightarrow Tr[F \wedge F]$$

The traceless symmetric tensor

$$\square \rightarrow T_{\mu\nu} = Tr \left[F_{\mu\nu}^2 - \frac{1}{4} g_{\mu\nu} F^2 \right]$$

is the conserved stress tensor dual to a massless graviton in 5d reflecting the translational symmetry of YM.

$$\square \rightarrow T_{\mu\nu;\rho\sigma}^4 = Tr \left[F_{\mu\nu} F_{\rho\sigma} - \frac{1}{2} (g_{\mu\rho} F_{\nu\sigma}^2 - g_{\nu\rho} F_{\mu\sigma}^2 - g_{\mu\sigma} F_{\nu\rho}^2 + g_{\nu\sigma} F_{\mu\rho}^2) + \frac{1}{6} (g_{\mu\rho} g_{\nu\sigma} - g_{\nu\rho} g_{\mu\sigma}) F^2 \right]$$

It has 10 independent d.o.f, it is not conserved and it should correspond to a similar **massive** tensor in 5d. We do not expect it to play an non-trivial role in the large- N_c , YM vacuum also for reasons of Lorentz invariance.

- Therefore the nontrivial fields are expected to be:

$$g_{\mu\nu}, \phi, a$$

The minimal effective string theory spectrum

- NS-NS $\rightarrow g_{\mu\nu}$, $B_{\mu\nu}$, ϕ
- RR $\rightarrow \text{Spinor}_5 \times \text{Spinor}_5 = F_0 + F_1 + F_2 + (F_3 + F_4 + F_5)$
- ♠ $F_0 \leftrightarrow F_5 \rightarrow C_4$, background flux \rightarrow no propagating degrees of freedom.
- ♠ $F_1 \leftrightarrow F_4 \rightarrow C_3 \leftrightarrow C_0$: C_0 is the axion, C_3 its 5d dual that couples to domain walls separating oblique confinement vacua.
- ♠ $F_2 \leftrightarrow F_3 \rightarrow C_1 \leftrightarrow C_2$: They are associated with baryon number (as we will see later when we add flavor). Dual operators are a mystery (topological currents?).
- In an ISO(3,1) invariant vacuum solution, only $g_{\mu\nu}, \phi, C_0 = a$ can be non-trivial.

$$ds^2 = e^{2A(r)}(dr^2 + dx_4^2) \quad , \quad a(r), \phi(r)$$

The relevant “defects”

- $B_{\mu\nu} \rightarrow$ Fundamental string (F_1). This is the QCD (glue) string: fundamental tension $\ell_s^2 \sim \mathcal{O}(1)$
- Its dual $\tilde{B}_\mu \rightarrow NS_0$: Tension is $\mathcal{O}(N_c^2)$. It is an effective magnetic baryon vertex binding N_c magnetic quarks.
- $C_4 \rightarrow D_3$ branes generating the gauge symmetry.
- $C_3 \rightarrow D_2$ branes : domain walls separating different oblique confinement vacua (where $\theta_{k+1} = \theta_k + 2\pi$). Its tension is $\mathcal{O}(N_c)$
- $C_2 \rightarrow D_1$ branes: These are the magnetic strings (strings attached to magnetic quarks) with tension $\mathcal{O}(N_c)$
- $C_5 \rightarrow D_4$: Space filling flavor branes. They must be introduced in pairs: $D_4 + \bar{D}_4$ for charge neutrality/tadpole cancellation.
- $C_1 \rightarrow D_0$ branes. These are the baryon vertices: they bind N_c quarks, and their tension is $\mathcal{O}(N_c)$.
- $C_0 \rightarrow D_{-1}$ branes: These are the Yang-Mills instantons.

An assessment of IR asymptotics

- As $\lambda \rightarrow \infty$ we assume that the potential terms dominate and we parameterize the effective action in the IR as

$$S_{eff} \sim \int \sqrt{g} \left[R + \frac{4(\partial\lambda)^2}{3\lambda^2} + V(\lambda) \right], \quad V(\lambda) = \frac{4}{3}\lambda^2 \left(\frac{dW}{d\lambda} \right)^2 + \frac{64}{27}W^2$$

Parameterize the IR asymptotics ($\lambda \rightarrow \infty$) as

$$W(\lambda) \sim (\log \lambda)^{\frac{P}{2}} \lambda^Q$$

- $Q > 2/3$ or $Q = 2/3$ and $P > 1$ leads to confinement and a singularity at finite $r = r_0$.

$$e^A(r) \sim \begin{cases} (r_0 - r)^{\frac{4}{9Q^2 - 4}} & Q > \frac{2}{3} \\ \exp \left[-\frac{C}{(r_0 - r)^{1/(P-1)}} \right] & Q = \frac{2}{3} \end{cases}$$

- $Q = 2/3$, and $0 \leq P < 1$ leads to confinement and a singularity at $r = \infty$
The scale factor e^A vanishes there as

$$e^A(r) \sim \exp[-Cr^{1/(1-P)}].$$

The asymptotic spectrum of glueballs is linear only if $P = \frac{1}{2}$

- $Q = 2/3, P = 1$ leads to confinement but the singularity may be at a finite or infinite value of r depending on subleading asymptotics of the superpotential.

♠ If $Q < 2\sqrt{2}/3$, no *ad hoc* boundary conditions are needed to determine the glueball spectrum: the singularity is “good” (repulsive).

♠ when $Q > 2\sqrt{2}/3$, the spectrum is not well defined without extra boundary conditions in the IR because both solutions to the mass eigenvalue equation are IR normalizable.

Selecting the IR asymptotics

Only the $Q = 2/3$, $0 \leq P < 1$ is compatible with

- Confinement (it happens non-trivially: a minimum in the string frame scale factor)
- Mass gap+discrete spectrum (except $P=0$)
- good singularity
- $R \rightarrow 0$ partly justifying the original assumption. More precisely: the string frame metric becomes flat at the IR . But $(\partial\phi)^2 \sim V(\lambda)$.

♠ It is interesting that the lower endpoint: $P=0$ corresponds to linear dilaton and flat space (string frame). It is confining with a mass gap but continuous spectrum.

- For linear asymptotic trajectories for fluctuations (glueballs) we must choose $P = 1/2$

$$V(\lambda) = \lambda^{\frac{4}{3}} [1 + c_1\lambda^2 + c_2\lambda^4 + \dots] \sim \lambda^{\frac{4}{3}} \sqrt{\log \lambda} + \text{subleading} \quad \text{as} \quad \lambda \rightarrow \infty$$

Further α' corrections

There are further dilaton terms generated by the 5-form in:

- The kinetic terms of the graviton and the dilaton $\sim \lambda^{2n}$.
- The kinetic terms on probe D_3 branes that affect the identification of the gauge-coupling constant, $\sim \lambda^{2n+1}$. There is also a multiplicative factor relating g_{YM^2} to e^ϕ , (not known). Can be traded for b_0 .
- Corrections to the identification of the energy. At $r = 0$, $E = 1/r$. There can be log corrections to our identification $E = e^A$, and these are a power series in $\sim \lambda^{2n}$.
- It is a remarkable fact that all such corrections affect the higher that the first two terms in the β -function (or equivalently the potential), that are known to be non-universal!

the metric is also insensitive to the change of b_0 by changing Λ .

Organizing the vacuum solutions

A useful variable is the phase variable

$$X \equiv \frac{\Phi'}{3A'} = \frac{\beta(\lambda)}{3\lambda} \quad , \quad e^\Phi \equiv \lambda$$

and a superpotential

$$W^2 - \left(\frac{3}{4}\right)^2 \left(\frac{\partial W}{\partial \Phi}\right)^2 = \left(\frac{3}{4}\right)^3 V(\Phi).$$

with

$$A' = -\frac{4}{9}W \quad , \quad \Phi' = \frac{dW}{d\Phi}$$

$$X = -\frac{3 d \log W}{4 d \log \lambda} \quad , \quad \beta(\lambda) = -\frac{9}{4} \lambda \frac{d \log W}{d \log \lambda}$$

♠ The equations have three integration constants: (two for Φ and one for A) One corresponds to the “gluon condensate” in the UV. It must be set to zero otherwise the IR behavior is unacceptable. The other is Λ . The third one is a gauge artifact (corresponds to overall translation in the radial coordinate).

The IR regime

For any asymptotically AdS_5 solution ($e^A \sim \frac{\ell}{r}$):

- The scale factor $e^{A(r)}$ is monotonically decreasing

*Girardello+Petrini+Porrati+Zaffaroni
Freedman+Gubser+Pilch+Warner*

- Moreover, there are only three possible, mutually exclusive IR asymptotics:

♠ *there is another asymptotic AdS_5 region, at $r \rightarrow \infty$, where $\exp A(r) \sim \ell'/r$, and $\ell' \leq \ell$ (equality holds if and only if the space is exactly AdS_5 everywhere);*

♠ there is a curvature singularity at some finite value of the radial coordinate, $r = r_0$;

♠ there is a curvature singularity at $r \rightarrow \infty$, where the scale factor vanishes and the space-time shrinks to zero size.

Wilson-Loops and confinement

- Calculation of the static quark potential using the vev of the Wilson loop calculated via an F-string worldsheet.

Rey+Yee, Maldacena

$$T E(L) = S_{\text{minimal}}(X)$$

We calculate

$$L = 2 \int_0^{r_0} dr \frac{1}{\sqrt{e^{4A_S(r)} - 4A_S(r_0)} - 1}.$$

It diverges when e^{A_S} has a minimum (at $r = r_*$). Then

$$E(L) \sim T_f e^{2A_S(r_*)} L$$

- **Confinement** $\rightarrow A_S(r_*)$ is finite. This is a more general condition that considered before as A_S is not monotonic in general.
- Effective string tension

$$T_{\text{string}} = T_f e^{2A_S(r_*)}$$

General criterion for confinement

- the geometric version:

A geometry that shrinks to zero size in the IR is dual to a confining 4D theory if and only if the Einstein metric in conformal coordinates vanishes as (or faster than) e^{-Cr} as $r \rightarrow \infty$, for some $C > 0$.

- It is understood here that a metric vanishing at finite $r = r_0$ also satisfies the above condition.

- ♠ the superpotential

A 5D background is dual to a confining theory if the superpotential grows as (or faster than)

$$W \sim (\log \lambda)^{P/2} \lambda^{2/3} \quad \text{as } \lambda \rightarrow \infty, \quad P \geq 0$$

- ♠ the β -function A 5D background is dual to a confining theory if and only if

$$\lim_{\lambda \rightarrow \infty} \left(\frac{\beta(\lambda)}{3\lambda} + \frac{1}{2} \right) \log \lambda = K, \quad -\infty \leq K \leq 0$$

(No explicit reference to any coordinate system) Linear trajectories correspond to $K = -\frac{3}{16}$

Classification of confining superpotentials

Classification of confining superpotentials $W(\lambda)$ as $\lambda \rightarrow \infty$ in IR:

$$W(\lambda) \sim (\log \lambda)^{\frac{P}{2}} \lambda^Q, \quad \lambda \sim E^{-\frac{9}{4}Q} \left(\log \frac{1}{E} \right)^{\frac{P}{2Q}}, \quad E \rightarrow 0.$$

- $Q > 2/3$ or $Q = 2/3$ and $P > 1$ leads to confinement and a singularity at finite $r = r_0$.

$$e^A(r) \sim \begin{cases} (r_0 - r)^{\frac{4}{9Q^2 - 4}} & Q > \frac{2}{3} \\ \exp \left[-\frac{C}{(r_0 - r)^{1/(P-1)}} \right] & Q = \frac{2}{3} \end{cases}$$

- $Q = 2/3$, and $0 \leq P < 1$ leads to confinement and a singularity at $r = \infty$. The scale factor e^A vanishes there as

$$e^A(r) \sim \exp[-Cr^{1/(1-P)}].$$

- $Q = 2/3, P = 1$ leads to confinement but the singularity may be at a finite or infinite value of r depending on subleading asymptotics of the superpotential.

♠ If $Q < 2\sqrt{2}/3$, no *ad hoc* boundary conditions are needed to determine the glueball spectrum \rightarrow One-to-one correspondence with the β -function. This is unlike standard AdS/QCD and other approaches.

- when $Q > 2\sqrt{2}/3$, the spectrum is not well defined without extra boundary conditions in the IR because both solutions to the mass eigenvalue equation are IR normalizable.

Confining β -functions

A 5D background is dual to a confining theory if and only if

$$\lim_{\lambda \rightarrow \infty} \left(\frac{\beta(\lambda)}{3\lambda} + \frac{1}{2} \right) \log \lambda = K, \quad -\infty \leq K \leq 0$$

(No explicit reference to any coordinate system). Linear trajectories correspond to $K = -\frac{3}{16}$

- We can determine the geometry if we specify K :
- $K = -\infty$: the scale factor goes to zero at some finite r_0 , not faster than a power-law.
- $-\infty < K < -3/8$: the scale factor goes to zero at some finite r_0 faster than any power-law.
- $-3/8 < K < 0$: the scale factor goes to zero as $r \rightarrow \infty$ faster than $e^{-Cr^{1+\epsilon}}$ for some $\epsilon > 0$.
- $K = 0$: the scale factor goes to zero as $r \rightarrow \infty$ as e^{-Cr} (or faster), but slower than $e^{-Cr^{1+\epsilon}}$ for any $\epsilon > 0$.

The borderline case, $K = -3/8$, is certainly confining (by continuity), but whether or not the singularity is at finite r depends on the subleading terms.

Parameters

- All dimensionless coefficients of the potential are a priori parameters. However, a simple form is typically chosen for simplicity. In our example we fit only one parameter.

- We also have M_p , and the AdS length, ℓ . Asking correct $T \rightarrow \infty$ thermodynamics fixes

$$(M_p \ell)^3 = \frac{1}{45\pi^2} \quad , \quad M_{\text{physical}} = M_p N_c^{\frac{2}{3}} = \left(\frac{8}{45\pi^2 \ell^3} \right)^{\frac{1}{3}} \simeq 4.6 \text{ GeV}$$

ℓ is not a parameter but a unit of length.

- We have 3 initial conditions in the system of graviton-dilaton equations:

- ♠ One is fixed by picking the branch that corresponds asymptotically to $\lambda \sim \frac{1}{\log(r\Lambda)}$

- ♠ The other fixes $\Lambda \rightarrow \Lambda_{QCD}$.

- ♠ The third is a gauge artifact as it corresponds to a choice of the origin of the radial coordinate.

Comments on confining backgrounds

- For all confining backgrounds with $r_0 = \infty$, although the space-time is singular in the Einstein frame, the string frame geometry is asymptotically flat for large r . Therefore only λ grows indefinitely.
- String world-sheets do not probe the strong coupling region, at least classically. The string stays away from the strong coupling region.
- Therefore: singular confining backgrounds have generically the property that the singularity is *repulsive*, i.e. only highly excited states can probe it. This will also be reflected in the analysis of the particle spectrum (to be presented later)
- The confining backgrounds must also screen magnetic color charges. This can be checked by calculating 't Hooft loops using D_1 probes:
 - ♠ All confining backgrounds with $r_0 = \infty$ and most at finite r_0 screen properly
 - ♠ In particular “hard-wall” AdS/QCD confines also the magnetic quarks.

Particle Spectra: generalities

- Linearized equation:

$$\ddot{\xi} + 2\dot{B}\dot{\xi} + \square_4\xi = 0 \quad , \quad \xi(r, x) = \xi(r)\xi^{(4)}(x), \quad \square\xi^{(4)}(x) = m^2\xi^{(4)}(x)$$

- Can be mapped to Schrodinger problem

$$-\frac{d^2}{dr^2}\psi + V(r)\psi = m^2\psi \quad , \quad V(r) = \frac{d^2B}{dr^2} + \left(\frac{dB}{dr}\right)^2 \quad , \quad \xi(r) = e^{-B(r)}\psi(r)$$

- Mass gap and discrete spectrum visible from the asymptotics of the potential.

- Large n asymptotics of masses obtained from WKB

$$n\pi = \int_{r_1}^{r_2} \sqrt{m^2 - V(r)} \, dr$$

- Spectrum depends only on initial condition for λ ($\sim \Lambda_{QCD}$) and an overall energy scale (e^A) that must be fixed.

- scalar glueballs

$$B(r) = \frac{3}{2}A(r) + \frac{1}{2} \log \frac{\beta(\lambda)^2}{9\lambda^2}$$

- tensor glueballs

$$B(r) = \frac{3}{2}A(r)$$

- pseudo-scalar glueballs

$$B(r) = \frac{3}{2}A(r) + \frac{1}{2} \log Z(\lambda)$$

- Universality of asymptotics

$$\frac{m_{n \rightarrow \infty}^2(0^{++})}{m_{n \rightarrow \infty}^2(2^{++})} \rightarrow 1 \quad , \quad \frac{m_{n \rightarrow \infty}^2(0^{+-})}{m_{n \rightarrow \infty}^2(0^{++})} = \frac{1}{4}(d-2)^2$$

predicts $d = 4$ via

$$\frac{m^2}{2\pi\sigma_a} = 2n + J + c,$$

The axion background

- The kinetic term of the axion is suppressed by $1/N_c^2$. (it is an angle in the gauge theory, it is RR in string theory)

$$\ddot{a} + \left(3\dot{A} + \frac{\dot{Z}(\lambda)}{Z(\lambda)} \right) \dot{a} = 0 \quad \rightarrow \quad \dot{a} = \frac{C e^{-3A}}{Z(\lambda)}$$

It can be interpreted as the flow equation of the effective θ -angle.

- The full solution is


$$a(r) = \theta_{UV} + 2\pi k + C \int_0^r \frac{e^{-3A}}{Z(\lambda)} \, , \quad C = \langle \text{Tr}[F \wedge F] \rangle$$

- The vacuum energy is


$$E(\theta_{UV}) = \frac{M^3}{2N_c^2} \int d^5x \sqrt{g} Z(\lambda) (\partial a)^2 = \frac{M^3}{2N_c^2} C a(r) \Big|_{r=0}^{r=r_0}$$

- Consistency requires to impose that $a(r_0) = 0$. This determines C and

$$E(\theta_{UV}) = -\frac{M^3}{2} \text{Min}_k \frac{(\theta_{UV} + 2\pi k)^2}{\int_0^{r_0} \frac{dr}{e^{3A} Z(\lambda)}} \, , \quad \frac{a(r)}{\theta_{UV} + 2\pi k} = \frac{\int_r^{r_0} \frac{dr}{e^{3A} Z(\lambda)}}{\int_0^{r_0} \frac{dr}{e^{3A} Z(\lambda)}}$$



(a)



(b)

(a) An example of the axion profile (normalized to one in the UV) as a function of energy, in one of the explicit cases we treat numerically. The energy scale is in MeV, and it is normalized to match the mass of the lowest scalar glueball from lattice data, $m_0 = 1475 \text{ MeV}$. The axion kinetic function is taken as $Z(\lambda) = Z_a(1 + c_a \lambda^4)$, with $c_a = 100$ (the masses do not depend on the value of Z_a). The vertical dashed line corresponds to $\Lambda_p \equiv \frac{1}{\ell} \frac{\exp\left[A(\lambda_0) - \frac{1}{b_0 \lambda_0}\right]}{(b_0 \lambda_0)^{b_1/b_0^2}}$. In this particular case $\Lambda = 290 \text{ MeV}$.

(b) A detail showing the different axion profiles for different values of c_a . The values are $c_a = 0.1$ (dashed line), $c_a = 10$ (dotted line) and $c_a = 100$ (solid line).

Tachyon dynamics

- In the vacuum the gauge fields vanish and $T \sim 1$. Only DBI survives

$$S[\tau] = T_{D_4} \int dr d^4x \frac{e^{4A_s(r)}}{\lambda} V(\tau) \sqrt{e^{2A_s(r)} + \dot{\tau}(r)^2} \quad , \quad V(\tau) = e^{-\frac{\mu^2}{2}\tau^2}$$

- We obtain the nonlinear field equation:

$$\ddot{\tau} + \left(3\dot{A}_S - \frac{\dot{\lambda}}{\lambda}\right) \dot{\tau} + e^{2A_S} \mu^2 \tau + e^{-2A_S} \left[4\dot{A}_S - \frac{\dot{\lambda}}{\lambda}\right] \dot{\tau}^3 + \mu^2 \tau \dot{\tau}^2 = 0.$$

- In the UV we expect

$$\tau = m_q r + \sigma r^3 + \dots \quad , \quad \mu^2 \ell^2 = 3$$

- We expect that the tachyon must diverge before or at $r = r_0$. We find that indeed it does **at the singularity**. For the $r_0 = \infty$ backgrounds

$$\tau \sim \exp\left[\frac{2}{a} \frac{R}{\ell^2} r\right] \quad \text{as} \quad r \rightarrow \infty$$

- Generically the solutions have spurious singularities: $\tau(r_*)$ stays finite but its derivatives diverges as:

$$\tau \sim \tau_* + \gamma \sqrt{r_* - r}.$$


The condition that they are absent determines σ as a function of m_q .

- The easiest spectrum to analyze is that of vector mesons. We find ($r_0 = \infty$)

$$\Lambda_{glueballs} = \frac{1}{R}, \quad \Lambda_{mesons} = \frac{3}{\ell} \left(\frac{\alpha \ell^2}{2R^2} \right)^{(\alpha-1)/2} \propto \frac{1}{R} \left(\frac{\ell}{R} \right)^{\alpha-2}.$$

This suggests that $\alpha = 2$. preferred also from the glue sector.

Fluctuations around the AdS₅ extremum



- In QCD we expect that

$$\frac{1}{\lambda} = \frac{1}{N_c e^{\phi}} \sim \frac{1}{\log r} \quad , \quad ds^2 \sim \frac{1}{r^2} (dr^2 + dx_{\mu} dx^{\mu}) \quad \text{as } r \rightarrow 0$$

- Any potential with $V(\lambda) \sim \lambda^a$ when $\lambda \ll 1$ gives a power different that of AdS₅
- There is an AdS₅ minimum at a finite value λ_* . This cannot be the UV of QCD as dimensions do not match.

Near an AdS extremum

$$V = \frac{12}{l^2} - \frac{16\xi}{3l^2}\phi^2 + \mathcal{O}(\phi^3) \quad , \quad \frac{18}{l}\delta A' = \delta\phi'^2 - \frac{4}{l^2}\phi^2 = \mathcal{O}(\delta\phi^2) \quad , \quad \delta\phi'' - \frac{4}{l}\delta\phi' - \frac{4\xi}{l^2}\delta\phi = 0$$

where $\phi \ll 1$. The general solution of the second equation is

$$\delta\phi = C_+ e^{\frac{(2+2\sqrt{1+\xi})u}{l}} + C_- e^{\frac{(2-2\sqrt{1+\xi})u}{l}}$$

For the potential in question

$$V(\phi) = \frac{e^{\frac{4}{3}\phi}}{l_s^2} \left[5 - \frac{N_c^2}{2} e^{2\phi} - N_f e^\phi \right] \quad , \quad \lambda_0 \equiv N_c e^{\phi_0} = \frac{-7x + \sqrt{49x^2 + 400}}{10} \quad , \quad x \equiv \frac{N_f}{N_c}$$

$$\xi = \frac{5}{4} \left[\frac{400 + 49x^2 - 7x\sqrt{49x^2 + 400}}{100 + 7x^2 - x\sqrt{49x^2 + 400}} \right] \quad , \quad \frac{l_s^2}{l^2} = e^{\frac{4}{3}\phi_0} \left[\frac{100 + 7x^2 - x\sqrt{49x^2 + 400}}{400} \right]$$

The associated dimension is $\Delta = 2 + 2\sqrt{1 + \xi}$ and satisfies

$$2 + 3\sqrt{2} < \Delta < 2 + 2\sqrt{6} \quad \text{or equivalently} \quad 6.24 < \Delta < 6.90$$

It corresponds to an irrelevant operator. It is most probably relevant for the Banks-Zaks fixed points.

Bigazzi+Casero+Cotrone+Kiritsis+Paredes

RETURN


Estimating the importance of logarithmic scaling

We keep the IR asymptotics of background II, but change the UV to power asymptoting AdS₅, with a small λ_* .

$$e^A(r) = \frac{\ell}{r} e^{-(r/R)^2}, \quad \Phi(r) = \Phi_0 + \frac{3r^2}{2R^2} \sqrt{1 + 3\frac{R^2}{r^2}} + \frac{9}{4} \log \frac{2\frac{r}{R} + 2\sqrt{\frac{r^2}{R^2} + \frac{3}{2}}}{\sqrt{6}}.$$



$$W_{conf} = W_0 \left(9 + 4b_0^2(\lambda - \lambda_*)^2 \right)^{1/3} \left(9a + (2b_0^2 + 3b_1) \log [1 + (\lambda - \lambda_*^2)] \right)^{2a/3}.$$

We fix parameters so that the physical QCD scale is the same (as determined from asymptotic slope of Regge trajectories).




The stars correspond to the asymptotically free background I with $b_0 = 4.2$ and $\lambda_0 = 0.05$; the squares correspond to the results obtained in the first background with $R = 11.4\ell$; the triangles denote the spectrum in the second background with $b_0 = 4.2$, $l_i = 0.071$ and $l_* = 0.01$. These values are chosen so that the slopes coincide asymptotically for large n .

Profile of coupling and scale factor




The scale factor and 't Hooft coupling that follow from β . $b_0 = 4.2$, $\lambda_0 = 0.05$, $A_0 = 0$. The units are such that $\ell = 0.5$. The dashed line represents the scale factor for pure AdS .

Dependence of absolute mass scale on λ_0



Dependence on initial condition λ_0 of the absolute scale of the lowest lying glueball (shown in Logarithmic scale)


Dependence of mass ratios on λ_0



The mass ratios R_{20}


$$R_{20} = \frac{m_{2++}}{m_{0++}}.$$

The glueball wavefunctions



Normalized wave-function profiles for the ground states of the 0^{++} (solid line), 0^{-+} (dashed line), and 2^{++} (dotted line) towers, as a function of the radial conformal coordinate. The vertical lines represent the position corresponding to $E = m_{0^{++}}$ and $E = \Lambda_p$.

Comparison of scalar and tensor potential




Effective Schrödinger potentials for scalar (solid line) and tensor (dashed line) glueballs. The units are chosen such that $\ell = 0.5$.

The lattice glueball data

J^{++}	Ref. I ($m/\sqrt{\sigma}$)	Ref. I (MeV)	Ref. II (mr_0)	Ref. II (MeV)	$N_c \rightarrow \infty(m/\sqrt{\sigma})$
0	3.347(68)	1475(30)(65)	4.16(11)(4)	1710(50)(80)	3.37(15)
0*	6.26(16)	2755(70)(120)	6.50(44)(7)	2670(180)(130)	6.43(50)
0**	7.65(23)	3370(100)(150)	NA	NA	NA
0***	9.06(49)	3990(210)(180)	NA	NA	NA
2	4.916(91)	2150(30)(100)	5.83(5)(6)	2390(30)(120)	4.93(30)
2*	6.48(22)	2880(100)(130)	NA	NA	NA
R_{20}	1.46(5)	1.46(5)	1.40(5)	1.40(5)	1.46(11)
R_{00}	1.87(8)	1.87(8)	1.56(15)	1.56(15)	1.90(17)


Available lattice data for the scalar and the tensor glueballs. Ref. I = [H. B. Meyer, \[arXiv:hep-lat/0508002\]](#). and Ref. II = [C. J. Morningstar and M. J. Peardon, \[arXiv:hep-lat/9901004\]](#) + [Y. Chen et al., \[arXiv:hep-lat/0510074\]](#). The first error corresponds to the statistical error from the continuum extrapolation. The second error in Ref.I is due to the uncertainty in the string tension $\sqrt{\sigma}$. (Note that this does not affect the mass ratios). The second error in the Ref. II is the estimated uncertainty from the anisotropy. In the last column we present the available large N_c estimates according to [B. Lucini and M. Teper, \[arXiv:hep-lat/0103027\]](#). The parenthesis in this column shows the total possible error followed by the estimations in the same reference.

Pseudoscalar glueballs




Lowest 0^{-+} glueball mass in MeV as a function of c_a in $Z(\lambda) = Z_a(1 + c_a\lambda^4)$.

α -dependence of scalar spectrum




The 0^{++} spectra for varying values of α that are shown at the right end of the plot. The symbol $*$ denotes the AdS/QCD result.

Comparison with lattice data: Ref II



Comparison of glueball spectra from our model with $b_0 = 2.55, \lambda_0 = 0.05$ (boxes), with the lattice QCD data from Ref. II (crosses) and the AdS/QCD computation (diamonds), for (a) 0^{++} glueballs; (b) 2^{++} glueballs. The masses are in MeV, and the scale is normalized to match the lowest 0^{++} state from Ref. II.

The thermodynamic quantities



Detailed plan of the presentation

- Title page 0 minutes
- Bibliography 1 minutes
- Introduction 3 minutes
- AdS/QCD 5 minutes
- Improving AdS/QCD 6 minutes
- A string theory for QCD: basic expectation 8 minutes
- The nature of the string theory 9 minutes
- Effective action I 10 minutes
- The UV regime 13 minutes
- The IR regime 15 minutes
- Improved Holographic QCD: a model 19 minutes
- Quarks ($N_f \ll N_c$) and mesons 23 minutes

THE DATA

- Concrete models: I 24 minutes
- Linearity of the glueball spectrum 25 minutes
- Comparison with lattice data (Meyer) 26 minutes
- The fit to Meyer Lattice data 27 minutes
- YM at finite temperature 31 minutes
- The general phase structure 32 minutes
- Temperature versus horizon position 33 minutes
- The free energy as a function of r_h 34 minutes
- The transition in the free energy 35 minutes
- Equation of state 36 minutes
- The speed of sound 37 minutes
- The specific heat 38 minutes
- Bulk viscosity 40 minutes
- Open problems 41 minutes

- The low dimension spectrum 43 minutes
- The minimal effective string theory spectrum 45 minutes
- The relevant defects 48 minutes
- An assessment of IR asymptotics 52 minutes
- Selecting the IR asymptotics 55 minutes
- Further α' corrections 57 minutes
- Organizing the vacuum solutions 59 minutes
- The IR regime 61 minutes
- Wilson loops and confinement 63 minutes
- General criterion for confinement 66 minutes
- Classification of confining superpotentials 69 minutes
- Confining β -functions 72 minutes
- Parameters 74 minutes
- Comments on confining backgrounds 76 minutes
- Particle Spectra: generalities 79 minutes

- The axion background 82 minutes
- Tachyon dynamics 86 minutes
- Fluctuations around the AdS_5 extremum 89 minutes
- Estimating the importance of logarithmic scaling 91 minutes
- The scale factor 93 minutes
- Dependence of absolute mass scale on λ_0 94 minutes
- Dependence of mass ratios on λ_0 95 minutes
- The glueball wavefunctions 96 minutes
- Comparison of scalar and tensor potential 97 minutes
- The lattice glueball data 98 minutes
- Pseudoscalar Glueballs 99 minutes
- α -dependence of scalar spectrum 100 minutes
- Comparison with lattice data: Ref II n101 minutes
- The thermodynamic quantities 102 minutes