

Effective form factors for free fermionic models at finite temperature

Oleksandr Gamayun

Warsaw University

SciPost Phys. 10, 070 (2021); SciPost Phys 5, 006 (2022);
PRB **105**, 085145 + 2202.07657:

Nikolai Iorgov and Yurij Zhuravlev & Daniel Chernowitz
Miłosz Panfil, Felipe Taha Sant'Ana

GGI 13/05/2022

Randomness, Integrability and Universality

Motivation: Integrable 1D models

- ▶ Heisenberg Spin-Chain

$$H_{XXZ} = -J \sum_{j=1}^N (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + \Delta S_j^z S_{j+1}^z)$$

Interacting Bose-Gas (Lieb-Liniger model)

- ▶

$$H = \sum_{j=1}^N \frac{p_i^2}{2m} + c \sum_{i < j} \delta(x_i - x_j)$$

Observables

$$\langle \mathbf{q} | S_j^z(t) S_{j'}^z(0) | \mathbf{q} \rangle = ? \quad \quad \langle \mathbf{q} | \rho(x, t) \rho(0, 0) | \mathbf{q} \rangle = ?$$

- ▶ inelastic neutron scattering
- ▶ Bragg spectroscopy
- ▶ Interference experiments

Motivation: Bethe Ansatz Solution

The eigenstate of N-quasiparticle (magnons, atoms etc) can be written as

$$|\mathbf{q}\rangle = |q_1, q_2, \dots, q_N\rangle = \sum_{\sigma} (-1)^{[\sigma]} A_{\sigma}(\{\mathbf{q}\}) e^{i \sum_{j=1}^N q_{\sigma[j]} x_j}$$

Periodic boundary conditions provide quantization

$$\theta_{\text{kin}}(q_j) + \frac{1}{L} \sum_{j \neq i} \theta_{\text{scat}}(q_i - q_j) = \frac{2\pi}{L} n_j$$

Hilbert space is given by the ordered sets of integers



$$\langle \mathbf{q} | \mathcal{O}(x, t) \mathcal{O}(0, 0) | \mathbf{q} \rangle = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + ixP_{\mathbf{k}}}$$


$E_{\mathbf{k}}$, $P_{\mathbf{k}}$ and $\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle$ are known by means of Algebraic Bethe Ansatz

$$|\mathbf{q}\rangle = B(q_1)B(q_2)\dots B(q_N)|0\rangle$$

Motivation: Summation over intermediate states

$$\langle \mathbf{q} | \mathcal{O}(x, t) \mathcal{O}(0, 0) | \mathbf{q} \rangle = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + ixP_{\mathbf{k}}}$$

- ▶ Numerics (ABACUS)
- ▶ Field theory ($k_F x \gg 1$, $k_F^2 t \gg 1$)



Left: Comparison between ABACUS and inelastic neutron scattering for $KCuF_3$. [PRL 111 137205].

Right: The threshold singularities in the Non-Linear Luttinger Liquid.

Both approaches fail at finite temperature !!!

Motivation: Hardcore Lieb-Liniger

$$H = \sum_{j=1}^N \frac{p_j^2}{2m} + c \sum_{i < j} \delta(x_i - x_j) \equiv \int dx \left[\frac{\partial_x \psi^+ \partial_x \psi}{2m} + c [\psi^+ \psi]^2 \right]$$

$$\rho(x, t) = \langle \psi^+(x, t) \psi(0, 0) \rangle = \sum_{\mathbf{k}} |\langle \mathbf{k} | \psi | \mathbf{q} \rangle|^2 e^{-it(E_{\mathbf{k}} - E_{\mathbf{q}}) - ix(P_{\mathbf{k}} - P_{\mathbf{q}})}$$

$$e^{ik_j L} = \prod_{i=1}^N \frac{k_j - k_i + ic}{k_j - k_i - ic}$$

$$e^{iq_j L} = \prod_{i=1}^{N+1} \frac{q_j - q_i + ic}{q_j - q_i - ic}$$

$$c \rightarrow \infty$$

$$k_j = \frac{2\pi}{L}(n_j - 1/2)$$

$$q_j = \frac{2\pi}{L} n_j$$

$$\rho(x, 0) \sim \det_{[-k_F, k_F]} \left(1 - \frac{2}{\pi} \frac{\sin(x(p - q))}{p - q} + e^{-ix(p+q)} \right) - \det_{[-k_F, k_F]} \left(1 - \frac{2}{\pi} \frac{\sin(x(p - q))}{p - q} \right)$$

$$k_F = \pi N/L$$

Fredholm determinants

$$\tau(x) = \det(1 + \hat{V})$$



$$\hat{V}f(q) = \int_{\gamma} dq' V(q, q') f(q')$$

- ▶ Infinite dimensional determinant

$$\det(1 + \hat{V}) = \lim_{N \rightarrow \infty} \det \left(\delta_{ij} + \frac{1}{N} V(x_i, x_k) \right) \Big|_{1 \leq i, k \leq N}$$

- ▶ Effective numerical evaluation (F. Bornemann "*On the Numerical Evaluation of Fredholm Determinants*" [0804.2543])

$$\int_a^b f(q) dq = \lim_{N \rightarrow \infty} \sum_{k=1}^N \omega_k f(x_k), \quad \det(1 + \hat{V}) = \lim_{N \rightarrow \infty} \det(\delta_{ij} + \sqrt{\omega_i} V(x_i, x_k) \sqrt{\omega_k}) \Big|_{1 \leq i, k \leq N}$$

Generalized Sine Kernels

$$\tau(x) = \det \left(1 + \frac{e^{2\pi i \nu} - 1}{\pi} n_F(q) \frac{\sin x(q-p)/2}{q-p} \right)$$

$$\tau_{XX} = \det_{[-\pi, \pi]} \left(1 - \frac{\omega_F(q)}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} - \frac{\omega_F(q)}{\pi} e^{-\frac{ip(x-1)+iq(x+1)}{2}} \right) - \det \left(1 - \frac{\omega_F(q)}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} \right)$$

- ▶ Random matrices
- ▶ Mobile impurity [SciPost Phys. 8, 053 (2020), New J. Phys. 18 (2016), 045005]
- ▶ Return probability from the domain wall initial state $|\text{DW}\rangle = |\uparrow\uparrow\dots\uparrow\downarrow\downarrow\dots\downarrow\rangle$ [J.M. Stephan, J. Stat (2017)]

$$\langle \text{DW} | e^{zH_{XXX}} | \text{DW} \rangle = \det_{\mathbb{R}_+} \left(1 - e^{-p^2/4} \frac{\sin \sqrt{z}(p-q)}{\pi(p-q)} e^{-q^2/4} \right)$$

- ▶ Persistence of spin configurations [I. Dornic (2018)] $n_F(q) = 1/\cosh(q)$
- ▶ Classical integrable systems $n_F(q) = r(q)$

Sine Kernel ($T = 0$)

$$\tau(x) = \tau(x, t=0) = \det_{[-k_F, k_F]} \left(1 + \frac{e^{2\pi i \nu} - 1}{\pi} \frac{\sin x(q-p)/2}{q-p} \right)$$

Form-factor presentation

$$\tau(x, t) = \langle \mathbf{q} | \mathcal{O}(x, t) \mathcal{O}(0, 0) | \mathbf{q} \rangle = \sum_{k_1 < k_2 \dots < k_N} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + iP_{\mathbf{k}}x}$$

$|\mathbf{q}\rangle$: free fermions: $q_i = \frac{2\pi n_i}{L}$; $|\mathbf{k}\rangle$: **shifted** free fermions: $k_i = \frac{2\pi(n_i - \nu)}{L}$

$$P_{\mathbf{k}} = \sum_i k_i, \quad E_{\mathbf{k}} = \sum_i k_i^2 / 2$$

The form-factor (overlap):

$$|\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 = \left(\frac{2}{L} \sin \pi \nu \right)^{2N} \left(\det_{N \times N} \frac{1}{k_i - q_j} \right)^2 \rightarrow |\langle \mathbf{q} | \mathbf{k} \rangle|^2.$$

$$\tau = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathbf{k} \rangle|^2 e^{-itE_{\mathbf{k}} + iP_{\mathbf{k}}x} = \det \left(\frac{2}{L} \oint \frac{dk}{\cot \frac{kL}{2} + \cot \pi \nu} \frac{e^{-itk^2/2 + ikx}}{(k - q_i)(k - q_j)} \right) = \det(1 + \hat{V})$$

Field theory treatment = Microscopic bosonization ($T = 0$)

- ▶ Form-Factor summation


$$\tau(x, t) = \sum_{\mathbf{k}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}} + itE_{\mathbf{k}}} = \det(1 + \hat{V})$$

- ▶ Orthogonality Catastrophe: $|\langle \mathbf{q} | \mathcal{O} | \mathbf{k}_{\text{vac}} \rangle|^2 = \mathcal{A}/N^{2\alpha}$
- ▶ Soft-mode summation

$$\tau(x, t) \sim \sum_{\text{IR}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}} + itE_{\mathbf{k}}} = \langle e^{\sqrt{\alpha}\varphi(x, t)} e^{-\sqrt{\alpha}\varphi(0, 0)} \rangle = \frac{\mathcal{A}}{(x - k_F t)^{\alpha}(x + k_F t)^{\alpha}}$$

- ▶ Nonlinear contributions

$$\tau(x, t) \sim \sum_{Q+\text{IR}} |\langle \mathbf{q} | \mathcal{O} | \mathbf{k} \rangle|^2 e^{-ixP_{\mathbf{k}} + itE_{\mathbf{k}} + ix(Q - k_F) + it(Q^2 - k_F^2)/2} = \frac{\mathcal{B}}{\sqrt{t}(x - k_F t)^{\tilde{\alpha}}(x + k_F t)^{\alpha}}$$



Slavnov (1989); Slavnov and Korepin (1991); A. Shashi, L. I. Glazman, J.-S. Caux, and A. Imambekov (2011); N. Kitanine, K.K. Kozlowski, J.-M. Maillet, N.A. Slavnov, and V. Terras (2009-2012); K.K. Kozlowski, J.-M. Maillet (2015);

Combinatorics of orthogonality catastrophe

Generic overlap

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \left(\frac{2}{L} \sin \pi \nu \right)^{2N} \left(\det_{N \times N} \frac{1}{k_i - q_j} \right)^2 = \left(\frac{2}{L} \sin \pi \nu \right)^{2N} \frac{\prod_{i>j} (k_i - k_j)^2 \prod_{i>j} (q_i - q_j)^2}{\prod_{i,j} (k_i - q_j)^2}.$$

Fermi sea integers

$$k_j = \frac{2\pi}{L}(n_j - \nu), \quad q_j = \frac{2\pi}{L}n_j, \quad n_j = -\frac{N-1}{2} + j - 1, \quad j = 1, 2, \dots, N$$

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \left(\frac{\sin \pi \nu}{\pi \nu} \right)^{2N} \prod_{i \neq j} \left(1 - \frac{\nu}{i-j} \right)^{-2} = \frac{G^2(1-\nu)G^2(1+\nu)G^4(N+1)}{G^2(N-\nu+1)G^2(N+\nu+1)}.$$

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \frac{G^2(1-\nu)G^2(1+\nu)}{N^{2\nu^2}}$$


For $\nu = \nu(k)$, ($\nu_{\pm} = \nu(\pm k_F)$), $k_F = \pi L/N$)

$$|\langle \mathbf{k}_{\text{vac}} | \mathbf{q} \rangle|^2 = \frac{G^2(1-\nu_-)G^2(1+\nu_+)(2\pi)^{\nu_- - \nu_+}}{N^{\nu_-^2 + \nu_+^2}} \exp \left(\int_{[-k_F, k_F]^2} \left(\frac{\nu(\lambda) - \nu(\mu)}{\lambda - \mu} \right)^2 d\lambda d\mu \right)$$

Static + zero temperature



$$\tau(x) = \frac{G^2(1-\nu)G^2(1+\nu)}{(-2ix)^{\nu^2}(2ix)^{\nu^2}} e^{-2i\nu x} + (\nu \rightarrow \nu + \mathbb{Z})$$

$\text{Re}[\tau(x,0)]$



Dynamics $T = 0$

$$\begin{aligned} \tau(x, t) = & \frac{G^2(1-\nu)G^2(1+\nu)}{(2i(t-x))^{\nu^2}(2i(x+t))^{\nu^2}} e^{-2i\nu x} + \\ & \frac{G^2(1-\nu)G^2(\nu)}{\nu^2(2i(t-x))^{(1+\nu)^2}(2i(x+t))^{\nu^2}} \left(\frac{x-t}{x+t}\right)^{2\nu} \frac{e^{-i(t-x)^2/(2t)-2i\nu x}}{(x/t-1)^2} \sqrt{\frac{2\pi}{-it}} \theta(x^2 > t^2) + \\ & + \frac{G^2(-\nu)G^2(1+\nu)}{\nu^2(2i(t-x))^{(1-\nu)^2}(2i(x+t))^{\nu^2}} \left(\frac{x+t}{x-t}\right)^{2\nu} \frac{e^{i(t-x)^2/(2t)-2i\nu x}}{(x/t-1)^2} \sqrt{\frac{2\pi}{it}} \theta(x^2 < t^2) + (\nu \rightarrow \nu + \mathbb{Z}) \end{aligned}$$



Finite temperature (Static)

$$\tau(x) = \det \left(1 + \frac{n_F(q)}{\pi} (e^{2\pi i \nu} - 1) \frac{\sin(x(p-q))}{p-q} \right)$$

- ▶ It is challenging to do microscopic

- ▶ Overlaps are too small $\sim e^{-cN}$
- ▶ Too many soft modes $\sim e^{cN}$



- ▶ Heuristic approach instead!

Dressing: inhomogeneous and complex valued!!!

$$\frac{n_F(q)}{\pi} (e^{2\pi i \nu} - 1) = \frac{e^{2\pi i \nu_T(q)} - 1}{\pi}, \quad \nu \rightarrow \nu_T(q) = \frac{1}{2\pi i} \log(1 + (e^{2\pi i \nu} - 1)n_F(q))$$

- ▶ Effectively **ONE** form factor


$$\boxed{\tau(x) \approx \exp \left(-ix \int_{-\infty}^{\infty} \nu_T(q) dq - \frac{1}{2} \int_{\mathbb{R}^2} \left(\frac{\nu_T(q) - \nu_T(q')}{q' - q} \right) dq dq' \right)}$$



$\text{Tr}(e^{-\beta H} \mathcal{O}(x, t) \dots) / \text{Tr} e^{-\beta H} = \langle \mathcal{O}(z = x+it) \dots \rangle_{S^1 \times \mathbb{R}^1} \stackrel{z \rightarrow z' = e^{2\pi z/\beta}}{=} \sim \langle \mathcal{O}(z') \dots \rangle_{\mathbb{R}^2}$

CFT prediction for correlation length:

$$\tau(x) \Big|_{T=0} = \frac{\mathcal{A}}{x^{\nu^2}} \implies \tau(x) = \frac{\mathcal{A}}{(\sinh(xT)/T)^{\nu^2}} \sim e^{-x/\xi} \implies 1/\xi \sim T ???$$



XY spins chain

$$\mathbf{H}_{XY} = -\frac{1}{2} \sum_{j=1}^L \left[\frac{1+\gamma}{2} \sigma_j^x \sigma_{j+1}^x + \frac{1-\gamma}{2} \sigma_j^y \sigma_{j+1}^y + h \sigma_j^z \right]$$

Spectrum of fermionic (Majorana) excitations

$$E(q) = \sqrt{(h - \cos q)^2 + \gamma^2 \sin^2 q}$$

Bogolyubov rotation angle

$$e^{i\theta(q)} = \frac{h - \cos q - i\gamma \sin q}{\sqrt{(h - \cos q)^2 + \gamma^2 \sin^2 q}}$$

Finite temperature spin-spin correlation function

$$\tau(x) = \tau(x) \equiv \frac{\text{Tr} \sigma_{x+1}^x \sigma_1^x e^{-\beta \mathbf{H}_{XY}}}{\text{Tr} e^{-\beta \mathbf{H}_{XY}}} = \det_{[-\pi, \pi]} (1 + \hat{V} + \hat{W}) - \det_{[-\pi, \pi]} (1 + \hat{V})$$

XY model [A.G. Izergin, V.S. Kapitonov, N.A. Kitanine, solv-int/9710028]

$$V(p, q) = -\frac{\omega_F(q)}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}}, \quad W(p, q) = -\frac{\omega_F(q)}{\pi} e^{-i(p+q)x/2} e^{-\frac{i(p-q)}{2}}$$

$$\omega_F(q) = \frac{1}{2} \left(1 - e^{i\theta(q)} \tanh \frac{\beta E(q)}{2} \right)$$

Form-factors

$$\tilde{\tau}(x) = \sum_{\mathbf{q}} |\langle \mathbf{k} | \mathbf{q} \rangle|^2 e^{-ix \left(\sum_{i=1}^{N+1} k_i - \sum_{i=1}^N q_i \right)}$$

with

$$e^{ikL} = e^{-2\pi i \nu(k)}, \quad e^{iqL} = 1.$$

$$|\langle \mathbf{k} | \mathbf{q} \rangle|^2 = A \left(\prod_{i=1}^{N+1} \frac{\sin \pi \nu(k_i)}{L} \right)^2 \frac{\prod_{i>j}^{N+1} \sin^2 \frac{k_i - k_j}{2} \prod_{i>j}^N \sin^2 \frac{q_j - q_i}{2}}{\prod_{i=1}^{N+1} \prod_{j=1}^N \sin^2 \frac{k_i - q_j}{2}}$$

$$\tilde{\tau}(x) = \det_{[-\pi, \pi]} \left(1 + \hat{V} + \hat{W} \right) - \det_{[-\pi, \pi]} \left(1 + \hat{V} \right)$$



$$V(p, q) = -\frac{e^{2\pi i \nu(q)} - 1}{\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} + O(e^{-\#x}), \quad W(p, q) = -\frac{e^{2\pi i \nu(q)} - 1}{\pi} e^{\frac{-i(p+q)x}{2}} e^{\frac{-i(p-q)}{2}}$$

$$e^{2\pi i \nu(k)} = 1 - 2\omega_F(k) = e^{i\theta(k)} \tanh \frac{\beta E(k)}{2}$$

$$e^{2\pi i \nu(k)} = 1 - 2\omega_F(k) = e^{i\theta(k)} \tanh \frac{\beta E(k)}{2}$$

$$\nu(\pi) - \nu(-\pi) = \delta \in \mathbb{Z}$$

$$e^{ikL} = e^{-2\pi i \nu(k)}, \quad e^{iqL} = 1.$$



Winding of the effective phase


Total number of solutions

$$e^{iqL} = 1, \quad q_j = \frac{2\pi}{L} \left(-\frac{L+1}{2} + j \right), \quad j = 1, 2, \dots L$$

$$e^{ikL} = e^{-2\pi i \nu(k)}, \quad k_j \approx \frac{2\pi}{L} \left(-\frac{L+1}{2} + j - \nu_j \right), \quad j = 1, 2, \dots L+\delta$$

$$\nu(k) \rightarrow \nu_\delta(k) = \nu(k) - \delta \frac{k + \pi}{2\pi} \implies e^{ik(L+\delta)} = (-1)^\delta e^{-2\pi i \nu_\delta(k)}$$

a)




$-q_F$

h

q_F

b)



$-q_F$

h

q_F

For $\delta = 1$ there is only one! form-factor

$$\delta < 0$$

$$\mathbf{q}^{a_1, \dots, a_n} = \{q_1, \dots, \hat{q}_{a_1}, \dots, \hat{q}_{a_n}, \dots q_L\} \quad \delta = 1 - n$$

$$\Delta P_{a_1, \dots, a_n} = \sum_{i=1}^{L-n+1} k_i - \sum_{i=1}^L q_i + \sum_{i=1}^n q_{a_i} \approx \delta\pi - \int_{-\pi}^{\pi} \nu(q) dq + \sum_{i=1}^n q_{a_i}.$$

$$e^{-ix\Delta P_{a_1, \dots, a_n}} |\langle \mathbf{k} | \mathbf{q}^{a_1, \dots, a_n} \rangle|^2 = \mathcal{A}_\delta[\nu] \prod_{i>j}^n \left(2 \sin \frac{q_{a_i} - q_{a_j}}{2} \right)^2 \prod_{i=1}^n \mathcal{Y}_{a_i},$$

$$\tau(x) = \det_{1 \leq j, k \leq n} [Y_\delta(x+j-k)] \exp \left(ix \int_{-\pi}^{\pi} \nu_\delta(q) dq - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu_\delta(q) - \nu_\delta(k)}{2 \sin \frac{q-k}{2}} \right]^2 \right),$$

where $\nu_\delta(q) \equiv \nu(q) - \delta(q + \pi)/(2\pi)$ has zero winding number and $Y_\delta(x)$ stands for

$$Y_\delta(x) = \int_{-\pi}^{\pi} \frac{dq}{2\pi} \left(e^{-2\pi i \nu(q)} - 1 \right) \exp \left(-i(x - \delta)q + i\delta\pi - \int_{-\pi}^{\pi} dk \nu_\delta(k) \cot \frac{q - k + i0}{2} \right).$$

$$e^{2\pi i \nu(k)} = 1 - 2\omega_F(k) = e^{i\theta(k)} \tanh \frac{\beta E(k)}{2} \quad \tau(x) = \mathcal{A}(T, h, \gamma) e^{-x/\xi(T, h, \gamma)}$$

Ferromagnetic $h \leq 1$ ($\delta = 1$)

$$\log \mathcal{A} = -\frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left[\frac{\nu(q) - \nu(k) - (q - k)/2\pi}{2 \sin \frac{q-k}{2}} \right]^2$$





$$\xi^{-1} = i\pi - i \int_{-\pi}^{\pi} \nu(q) dq = -\frac{1}{2\pi} \int_{-\pi}^{\pi} dk \log \tanh \frac{\beta E(k)}{2}$$

Paramagnetic $h > 1$ ($\delta = 0$)

$$\xi^{-1} = -\frac{1}{2\pi} \int_{-\pi}^{\pi} dk \log \tanh \frac{\beta E(k)}{2} + \log y_+, \quad y_{\pm} = \frac{h + \sqrt{h^2 + \gamma^2 - 1}}{1 \pm \gamma}$$

$$\log \mathcal{A} = \log \frac{2}{\beta \sqrt{h^2 + \gamma^2 - 1}} - i \int_{-\pi}^{\pi} dq \nu(q) \frac{e^{iq} + y_+}{e^{iq} - y_+} - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dk \left(\frac{\nu(q) - \nu(k)}{2 \sin \frac{q-k}{2}} \right)^2$$

Correlation functions



Bonus: from Fredholm to Toeplitz and back

$$\det(1 + \hat{S}_\nu) = \det_{0 \leq n, m \leq x-1} c_{n-m}, \quad c_k = \int_{-\pi}^{\pi} \frac{dq}{2\pi} e^{2\pi i \nu(q)} e^{-ikq}$$

$$\hat{S}_\nu(p, q) = \frac{e^{2\pi i \nu(p)} - 1}{2\pi} \frac{\sin \frac{x(p-q)}{2}}{\sin \frac{p-q}{2}} \sim \frac{e^{2\pi i \nu(p)} - 1}{2\pi} \sum_{n=0}^{x-1} e^{in(q-p)} \sim \sum_n \mathcal{A}_{qn} \mathcal{B}_{np}$$

$$\det(1 + \mathcal{AB}) = \det(1 + \mathcal{BA})$$

$$\det(1 + \hat{S}_\nu + \delta \hat{V}_\nu) - \det(1 + \hat{S}_\nu) = \det_{0 \leq n, m \leq x-1} \tilde{c}_{n-m}$$

$$\tilde{c}_k = \int_{-\pi}^{\pi} \frac{dq}{2\pi} e^{2\pi i \nu_1(q)} e^{-ikq}, \quad \nu_1(q) = \nu(q) - \frac{q + \pi}{2\pi}$$

$$\det(1 + \hat{S}_\nu + \delta \hat{V}_\nu) - \det(1 + \hat{S}_\nu) = \det(1 + \hat{S}_{\nu_1})$$

Szegő theorem for Toeplitz determinant [Szegő (1915), Fisher & Hartwig (1969)]

$$\log \det_{0 \leq i, j \leq x-1} c_{i-j} = x k_0 + \sum_{n=1}^{\infty} n k_n k_{-n}$$

$$\nu_\delta(q) = \frac{-1}{2\pi i} \sum_{n=-\infty}^{\infty} k_n e^{iqn} \quad - \frac{1}{2} \int_{-\pi}^{\pi} dq \int_{-\pi}^{\pi} dp \left[\frac{\nu_\delta(q) - \nu_\delta(p)}{2 \sin \frac{q-p}{2}} \right]^2 = \sum_{n=1}^{\infty} n k_n k_{-n},$$

$$\mathcal{A} = XY,$$

where:

$$X = \prod_{l=1}^{\infty} \frac{(1 - \lambda_1^{-1} f_{2l-1}) (1 - \lambda_1^{-1} g_{2l-1}) (1 - \lambda_2^{-1} f_{2l-1}) (1 - \lambda_2^{-1} g_{2l-1})}{(1 - \lambda_1^{-1} f_{2l}) (1 - \lambda_1^{-1} g_{2l}) (1 - \lambda_2^{-1} f_{2l}) (1 - \lambda_2^{-1} g_{2l})}$$

$$Y = \prod_{i,j=1}^{\infty} \frac{(1 - f_{2j} f_{2i-1})(1 - f_{2i} f_{2j-1})(1 - g_{2j} g_{2i-1})(1 - g_{2i} g_{2j-1})}{(1 - f_{2j} f_{2i})(1 - f_{2j-1} f_{2i-1})(1 - g_{2j} g_{2i})(1 - g_{2j-1} g_{2i-1})} \times \\ \times \frac{(1 - f_{2j} g_{2i-1})(1 - f_{2i} g_{2j-1})(1 - g_{2j} f_{2i-1})(1 - g_{2i} f_{2j-1})}{(1 - f_{2j} g_{2i})(1 - g_{2j} f_{2i})(1 - g_{2j-1} f_{2i-1})(1 - f_{2j-1} g_{2i-1})}$$

and $\lambda_1, \lambda_2, f, g$ are defined as

$$\lambda_1 = \left\{ h + [h^2 - (1 - \gamma^2)]^{1/2} \right\} / (1 - \gamma), \quad \lambda_2 = \left\{ h - [h^2 - (1 - \gamma^2)]^{1/2} \right\} / (1 - \gamma)$$

$$f_k = \frac{h + W_k}{1 - \gamma^2} - \left[\left(\frac{h + W_k}{1 - \gamma^2} \right)^2 - 1 \right]^{1/2}, \quad g_k = \frac{h - W_k}{1 - \gamma^2} - \left[\left(\frac{h - W_k}{1 - \gamma^2} \right)^2 - 1 \right]^{1/2}$$

with

$$W_k = \{ \gamma^2 h^2 - (1 - \gamma^2) [\gamma^2 + (k\pi)^2 \beta^{-2}] \}^{1/2}$$


Dynamics

$$\tau(x, t) = \det \left(\mathbb{1} + \rho(q) \frac{e_+(p)e_-(q) - e_-(p)e_+(q)}{p - q} \right)$$

$$e_-(q) = e^{\frac{i}{2}xq - \frac{i}{4}tq^2}, \quad e_+(q) = -\frac{\sin^2(\pi\nu)e^{-\frac{i}{2}xq + \frac{i}{4}tq^2}}{\pi} \left[\cot(\pi\nu) + i\text{Erf}\left(\frac{(i+1)(x-qt)}{2\sqrt{t}}\right) \right]$$

$$\text{Erf}\left(\frac{(x-qt)(1+i)}{2\sqrt{t}}\right) \rightarrow \text{Sign}(x/t - q)$$

$$\nu_T(q) = \frac{1}{2\pi i} \log \left[1 + (e^{2\pi i \nu} - 1)\rho(q) \right] \theta_\epsilon(x/t - q) - \frac{1}{2\pi i} \log \left[1 + (e^{-2\pi i \nu} - 1)\rho(q) \right] \theta_\epsilon(q - x/t).$$



$$\theta_\epsilon(z) = \frac{1 + S(z/\epsilon)}{2}, \quad \epsilon \sim \frac{1}{\sqrt{t}}$$

$$\log \tau \approx C_0 - \frac{1}{2} \Delta^2 \log t + i \int dk \nu_T(k) (x - tk)$$

$$\Delta = \frac{i}{2\pi} \log \left(1 + 2 \left(\cos(2\pi\nu) - 1 \right) \left(\rho - \rho^2 \right) \right)$$

$$\rho = \rho(x/t)$$

Summary and outlook

- ▶ Riemann Hilbert Problem for Fredholm determinant
- ▶ Phase shift dressing
- ▶ Different types of soft mode contributions
- ▶ Universality
- ▶ Asymptotic for classical integrable models?
- ▶ Relation with QTM? Thermal form-factors?