


Lecture notes GGI 2023 - Becattini



Bibliography

- Density operator → 1902.01089
- Entropy current → 1903.05422
- Spin and polarization → 2004.04050 and 2103.10917

Classical approach to relativistic hydrodynamics

Decomposition of currents onto a four-velocity

$$T^{\mu\nu} = \rho' u^\mu u^\nu + p' \Delta^{\mu\nu} + q^\mu u^\nu + q^\nu u^\mu + \Pi^{\mu\nu}$$

$$\rho' \equiv T^{\mu\nu} u_\mu u_\nu$$

$$q^\mu \equiv \Delta_\alpha^\mu T^{\alpha\beta} u_\beta$$

$$\Delta^{\mu\nu} \equiv g^{\mu\nu} - u^\mu u^\nu$$

$$\Delta^{\mu\alpha} \Delta_\alpha^\nu = \Delta^{\mu\nu}$$

$$\Pi^{\mu\nu} \equiv (\Delta_\alpha^\mu \Delta_\beta^\nu - \frac{1}{3} \Delta_{\alpha\beta} \Delta^{\mu\nu}) T^{\alpha\beta}$$

\Rightarrow

$$q \cdot u = 0$$

$$u_\mu \Pi^{\mu\nu} = \Pi^{\mu\nu} u_\nu = 0$$

$$\Delta_{\mu\nu} \Pi^{\mu\nu} = 0$$

$$p' = \frac{1}{3} \Delta_{\mu\nu} T^{\mu\nu}$$

$$j^\mu = n' u^\mu + J^\mu \quad n' \equiv j \cdot u \quad J^\mu \equiv \Delta_\alpha^\mu j^\alpha$$

All of them are functions of x

Equilibrium $U \equiv U_{eq} = \text{constant}$

$$\lim_{U \rightarrow U_{eq}} T^{\mu\nu} = (\rho_{eq} + p_{eq}) u_{eq}^\mu u_{eq}^\nu - p_{eq} g^{\mu\nu} \Rightarrow \lim_{U \rightarrow U_{eq}} q_\mu \nabla^\mu = 0$$

$$\lim_{U \rightarrow U_{eq}} j^\mu = n_{eq} u_{eq}^\mu$$

$\rho_{eq}, p_{eq}, n_{eq}$ are the thermodynamic equilibrium functions

$$\rho_{eq}(T, \mu) \quad p_{eq}(T, \mu) \quad n_{eq}(T, \mu) \quad \rho_{eq} + p_{eq} = TS_{eq} + \mu n_{eq}$$

$$\Delta \rho \equiv \rho'(x) - \rho_{eq}(x) = \rho'(x) - \rho(x)$$

$$\pi \equiv p'(x) - p_{eq}(x) = p'(x) - p(x)$$

$$\Delta n \equiv n'(x) - n_{eq}(x) = n'(x) - n(x) \rightsquigarrow \text{disregarded for the sake of simplicity}$$

Derivation of constitutive equations

$$Tds = d\rho - \mu dn \Rightarrow \frac{\partial s}{\partial \rho} = \frac{1}{T} \quad \frac{\partial s}{\partial n} = -\frac{\mu}{T} \Rightarrow \partial_\rho s = \frac{1}{T} \partial_\rho \rho - \frac{\mu}{T} \partial_\rho n$$

$$u^\mu \partial_\mu s \equiv \dot{s} \Rightarrow T\dot{s} = \dot{\rho} - \mu \dot{n}$$

$$\partial_\mu T^{\mu\nu} = 0 \Rightarrow \partial_\mu [(\rho + p) u^\mu u^\nu - pg^{\mu\nu} + q^\mu u^\nu + q^\nu u^\mu + \Pi^{\mu\nu} + \pi \Delta^{\mu\nu} + \Delta g u^\mu u^\nu] = 0$$

Contract with u_ν

$$u^\mu \partial_\mu (\rho + p) + (\rho + p) \partial \cdot u - u^\nu \partial_\nu p + \partial \cdot q + u_\nu u^\mu \partial_\mu q^\nu + u_\nu \partial_\mu \Pi^{\mu\nu} + \pi u_\nu \partial_\mu \Delta^{\mu\nu} + \partial_\mu (\Delta g u^\mu)$$

$$= \dot{\rho} + \dot{p} + (\rho + p) \partial \cdot u - \dot{p} + \partial \cdot q - q_\nu \underbrace{u^\mu \partial_\mu u^\nu}_{A_\nu} - \Pi^{\mu\nu} \partial_\mu u_\nu - \pi \partial \cdot u + \partial \cdot (\Delta g u)$$

acceleration field

$$= \dot{\rho} + \cancel{\dot{p}} + (\rho + p) \partial \cdot u - \cancel{\dot{p}} + \partial \cdot q - q \cdot A - \Pi^{\mu\nu} \partial_\mu u_\nu - \pi \partial \cdot u + \partial \cdot (\Delta \rho u) = 0$$

$$\Rightarrow \dot{\rho} = -(\rho + p) \partial \cdot u - \partial \cdot q + q \cdot A + \Pi : \partial u + \pi \partial \cdot u - \partial \cdot (\Delta \rho u)$$

Similarly

$$\partial_\mu j^\mu = 0 \Rightarrow \partial_\mu (n u^\mu + J^\mu) = 0 \quad n \partial \cdot u + \dot{n} + \partial \cdot J = 0 \quad \text{and}$$

$$\dot{n} = -n \partial \cdot u - \partial \cdot J$$

Altogether

$$\begin{aligned} T \dot{s} &= -(\rho + p) \partial \cdot u - \partial \cdot q + q \cdot A + \Pi : \partial u + \pi \partial \cdot u + \mu n \partial \cdot u + \mu \partial \cdot J - \partial \cdot (\Delta \rho u) \\ &= (-\rho - p + \mu n) \partial \cdot u - \partial \cdot q + q \cdot A + \Pi : \partial u + \pi \partial \cdot u + \mu \partial \cdot J - \partial \cdot (\Delta \rho u) \\ &= -T s \partial \cdot u - \partial \cdot q + q \cdot A + \Pi : \partial u + \pi \partial \cdot u + \mu \partial \cdot J - \partial \cdot (\Delta \rho u) \end{aligned}$$

$$\dot{s} + s \partial \cdot u = -\frac{\partial \cdot q}{T} + \frac{q \cdot A}{T} + \frac{\nabla \cdot \partial u}{T} + \frac{\pi}{T} \partial \cdot u + \frac{\mu}{T} \partial \cdot J - \frac{\partial \cdot (\Delta g u)}{T}$$

$\brace{ = -\partial \cdot \left(\frac{q}{T} \right) + q \cdot \partial \left(\frac{1}{T} \right) + \frac{q \cdot A}{T} + \frac{\nabla \cdot \partial u}{T} + \frac{\pi}{T} \partial \cdot u + \partial \cdot \left(\frac{\mu}{T} J \right) - J \cdot \partial \left(\frac{\mu}{T} \right) - \frac{\partial \cdot (\Delta g u)}{T}}$

↓

$$\partial_\mu (s u^\mu) = -\partial \cdot \left(\frac{q}{T} \right) + q \cdot \partial \left(\frac{1}{T} \right) + q \cdot A + \nabla \cdot \frac{\partial u}{T} + \frac{\pi}{T} \partial \cdot u + \partial \cdot \left(\frac{\mu}{T} J \right) - J \cdot \partial \left(\frac{\mu}{T} \right)$$

$$- \partial \cdot \left(\frac{\Delta g u}{T} \right) + \Delta g u \cdot \partial \left(\frac{1}{T} \right)$$

$$\partial \cdot \left(s u + \frac{q}{T} - \frac{\mu}{T} J + \frac{\Delta g u}{T} \right) = q \cdot \left(A + \partial \left(\frac{1}{T} \right) \right) + \nabla \cdot \frac{\partial u}{T} + \frac{\pi}{T} \partial \cdot u - J \cdot \partial \left(\frac{\mu}{T} \right)$$

$\begin{matrix} \text{entropy} \\ \text{current} \end{matrix} \quad s^\mu$

$$+ \Delta g u \cdot \partial \left(\frac{1}{T} \right)$$

Constitutive equations requiring $\partial_\mu S^\mu \geq 0$

E.g. $q = -k \left(A + \partial \left(\frac{1}{T} \right) \right)$ $k > 0$ thermal conductivity

Now:

$$\begin{array}{l} T^{\mu\nu}[T, u, \mu] \\ j^\mu[T, u, \mu] \end{array} \rightarrow \begin{array}{l} \partial_\mu T^{\mu\nu} = 0 \\ \partial_\mu j^\mu = 0 \end{array}$$

5 equations for 5 unknowns
 $T, u, \mu \quad (u \cdot u = 1)$

Two fundamental questions

1) Is it true that $\varphi + p = Ts + \mu n$ out of equilibrium (or in different global equilibria)?

Does $Tds = d\varphi - \mu dn$ hold?

2) How to deal with quantum observables? Spin


DENSITY OPERATOR

$$\hat{\rho} = ?$$

$$T^{\mu\nu}(x, t) = \text{Tr}(\hat{\rho}(0)\hat{T}^{\mu\nu}(x, t))$$

Heisenberg $\hat{T}^{\mu\nu}(x, t) = e^{i\hat{H}t} \hat{T}^{\mu\nu}(x, 0) e^{-i\hat{H}t}$ $\hat{\rho}$ fixed


Schrödinger $\hat{T}^{\mu\nu}(x, 0)$ fixed $\hat{\rho}(t) = e^{-i\hat{H}t} \hat{\rho}(0) e^{i\hat{H}t}$



Suppose that a good approximation can be found of

$$\hat{\rho}(t_0) = e^{-i\hat{H}t_0} \hat{\rho}(0) e^{i\hat{H}t_0} \text{ at a suitable time } t_0.$$

→ evolve $\hat{T}^{\mu\nu}$ or any other fields from t_0


$$\Sigma_0 \rightarrow \hat{\rho}(\Sigma_0)$$

“local equilibrium is achieved”

$$S = -\text{Tr}(\hat{\rho} \log \hat{\rho})$$

10 constants P_o^μ $J_o^{\mu\nu}$ mean values + some charges $Q_{o:i}$

Each constant \longleftrightarrow Lagrange multiplier

$$\begin{aligned} F[\hat{\rho}] = & -\text{Tr}(\hat{\rho} \log \hat{\rho}) - b_\mu (\text{Tr}(\hat{\rho} \hat{P}^\mu) - P_o^\mu) + \underline{\omega}_{\mu\nu} \left(\text{Tr}(\hat{\rho} \hat{J}^{\mu\nu}) - J_o^{\mu\nu} \right) \\ & + \sum_i \zeta_i (\text{Tr}(\hat{\rho} \hat{Q}_i) - Q_{o:i}) + \lambda (\text{Tr}(\hat{\rho}) - 1) \end{aligned}$$


$$\frac{\delta F}{\delta \hat{\rho}} = 0 \quad \text{Solution} \quad \hat{\rho} = \frac{e^{-b \cdot \hat{P} + \frac{1}{2} \omega \cdot \hat{J} + \sum_i \zeta_i \hat{Q}_i}}{Z}$$

$$\text{Problem: prove it} \quad Z = \text{Tr} \left(e^{-b \cdot \hat{P} + \frac{1}{2} \omega \cdot \hat{J} + \sum_i \zeta_i \hat{Q}_i} \right)$$

$\hat{\rho}$ approximated by $\hat{\rho}_{LE}(\Sigma_0)$

$$\hat{\rho}_{LE}(\Sigma_0) = \exp\left[-\int d\Sigma_\mu \hat{T}^\mu \beta_\nu - \zeta \hat{j}^\mu\right] / Z$$

Gauss theorem



$$\begin{aligned}
 - \int_{\Sigma_0} d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu + \int_{\Sigma_t} d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu &= \\
 = \int d\Omega \nabla_\mu (\hat{T}^{\mu\nu} \beta_\nu) & \\
 \text{if } \int_{\Sigma_L} d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu &= 0
 \end{aligned}$$

$$-\int_{\Sigma_0} d\Sigma \hat{T}^{\mu\nu} \beta_\nu = -\int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu + \int d\Omega \hat{T}^{\mu\nu} \nabla_\mu \beta_\nu$$

$$\Rightarrow \hat{P}_{LE}(\Sigma_0) = \frac{1}{Z} \exp \left[- \int d\Sigma_r (\hat{T}^{\mu\nu} \beta_\nu - \zeta \hat{j}^\mu) + \int d\Omega (\hat{T}^{\mu\nu} \nabla_\mu \beta_\nu - \nabla_\mu \zeta \cdot \hat{j}^\mu) \right]$$

Linear response

$$\hat{\rho} = \frac{1}{Z} e^{\hat{A} + \hat{B}}$$

$$Z = \text{Tr}(e^{\hat{A} + \hat{B}})$$

Kubo identity

$$e^{\hat{A} + \hat{B}} = [I + \int_0^1 dz e^{z(\hat{A} + \hat{B})} \hat{B} e^{-z\hat{A}}] e^{\hat{A}}$$

can be iterated

$$\Rightarrow e^{\hat{A} + \hat{B}} \cong [I + \int_0^1 dz e^{z\hat{A}} \hat{B} e^{-z\hat{A}}] e^{\hat{A}}$$

linear response

$$\hat{\rho} \cong \hat{\rho}_A (1 - \langle \hat{B} \rangle_A) + \int_0^1 dz e^{z\hat{A}} \hat{B} e^{-z\hat{A}} \hat{\rho}_A$$

$$\hat{\rho}_A = \frac{e^{\hat{A}}}{\text{Tr}(e^{\hat{A}})}$$

Example:

$$\hat{A} = - \int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu - \zeta \hat{j}^\mu$$

$$\hat{\rho}_{LE}(0) \cong \hat{\rho}_{LE} (1 - \langle \hat{B} \rangle_{LE}) + \int_0^1 dz e^{z\hat{A}} \hat{B} e^{-z\hat{A}} \hat{\rho}_{LE}$$

$$\langle \hat{T}^{\mu\nu}(x) \rangle = \text{Tr} (\hat{\rho} \hat{T}^{\mu\nu}(x))$$

$$\hat{B} = \int d\Omega \hat{T}^{\mu\nu} \partial_\mu \beta_\nu - \partial_\mu \zeta \hat{j}^\mu$$

$$\langle \hat{T}^{\mu\nu}(x) \rangle \cong \langle \hat{T}^{\mu\nu}(x) \rangle_{LE} + \int_0^1 dz \langle \hat{T}^{\mu\nu}(x) e^{z\hat{A}} \hat{B} e^{-z\hat{A}} \rangle_{LE} - \langle \hat{T}^{\mu\nu}(x) \rangle_{LE} \langle \hat{B} \rangle_{LE}$$

$$= \langle \hat{T}^{\mu\nu}(x) \rangle_{LE} + \int_0^1 dz \langle \hat{T}^{\mu\nu}(x) e^{z\hat{A}} \hat{B} e^{-z\hat{A}} \rangle_{LE,C} \quad \rangle_C \rightarrow \begin{matrix} \text{subtraction} \\ \text{of mean values} \end{matrix}$$

$$= \langle \hat{T}^{\mu\nu}(x) \rangle_{LE} + \int_0^1 dz \int d\Omega(y) \langle \hat{T}^{\mu\nu}(x) e^{z\hat{A}} \hat{T}^{\rho\sigma}(y) e^{-z\hat{A}} \rangle_{LE,C} \partial_\rho \beta_\sigma(y) + \dots$$

Hydro approx. $\partial_\rho \beta_\sigma(y) \simeq \partial_\rho \beta_\sigma(x)$

$$\Rightarrow \langle \hat{T}^{\mu\nu}(x) \rangle_{LE} + \partial_\rho \beta_\sigma(x) \int_0^1 dz \int d\Omega(y) \langle \hat{T}^{\mu\nu}(x) e^{z\hat{A}} \hat{T}^{\rho\sigma}(y) e^{-z\hat{A}} \rangle_{LE,C} + \dots$$

What about hydro equations?

① $n_p T^{\mu\nu} = n_p T_{LE}^{\mu\nu} \rightarrow$ find $\beta_n(x)$, all we need. Depends on the foliation

Similar to the definition of other frames

$$T^{\mu\nu} u_\mu = \lambda u^\nu \text{ eigenvector}$$

How to determine $T^{\mu\nu}$? $\rightarrow n_p \delta T^{\mu\nu} [\beta_\Sigma, g, \Sigma] = 0$

② Apparently $T^{\mu\nu}(x) = \text{Tr} \left[e^{\frac{\hat{A} + \hat{B}}{Z}} \hat{T}^{\mu\nu}(x) \right] \rightarrow T^{\mu\nu}(x) = T^{\mu\nu}[\beta_\Sigma, g, \Sigma]$

$$\frac{\delta}{\delta \Sigma} T^{\mu\nu}[\beta_\Sigma, g, \Sigma] = 0 \quad \text{eventually}$$

Expand in gradients $\rightarrow T^{\mu\nu}(\beta_\Sigma, \partial\beta_\Sigma, \partial^2\beta_\Sigma, \dots, g, \dots, \Sigma)$

$$T^{\mu\nu} = a \beta_\Sigma^\mu \beta_\Sigma^\nu + \dots$$

Entropy current

1) $S = -\text{Tr}(\hat{\rho} \log \hat{\rho}) \longrightarrow -\text{Tr}(\hat{\rho}_{LE} \log \hat{\rho}_{LE})$ $\frac{d\hat{\rho}_{LE}}{dt} \neq 0$

maximized
↓ constrained ignorance

2) Extensivity

Sia $\hat{\rho}(\lambda) = \frac{1}{Z(\lambda)} \exp \left[-\lambda \int d\Sigma_\mu \hat{T}^{\mu\nu} B_\nu \right]$ ($S=0$ per semplicità) $\hat{\rho} \equiv \hat{\rho}(1)$

Corrisponde a moltiplicare la temperatura conovente $\frac{1}{\sqrt{\beta^2}}$ per $\frac{1}{\lambda}$.

$$Z(\lambda) = \text{Tr} \left(\exp \left[-\lambda \int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu \right] \right) \quad \Rightarrow \quad \frac{\partial \log Z}{\partial \lambda}$$

$$= \frac{1}{Z(\lambda)} \frac{\partial}{\partial \lambda} \text{Tr} \left(\exp \left[-\lambda \int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu \right] \right) = \frac{\text{Tr}}{Z(\lambda)} \left(\exp \left[-\lambda \int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu \right] (-1) \int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu \right) =$$

$$= - \int d\Sigma_r \langle \hat{T}^{\mu\nu} \rangle(\lambda) \beta_\nu \quad \langle \hat{T}^{\mu\nu} \rangle(\lambda) \text{ mean value with } \hat{\rho}(\lambda)$$

Integrating between 1 and λ_0

$$\int_1^{\lambda_0} d\lambda \frac{\partial}{\partial \lambda} \log Z(\lambda) = \log Z(\lambda_0) - \log Z = - \int_1^{\lambda_0} d\lambda \int d\Sigma_r \langle \hat{T}^{\mu\nu} \rangle(\lambda) \beta_\nu = \int d\Sigma_r \left(\int_{\lambda_0}^1 d\lambda \langle \hat{T}^{\mu\nu} \rangle(\lambda) \beta_\nu \right)$$

If λ_0 such that $\log Z(\lambda_0) = 0$ avrō

$$\log Z = \int d\Sigma_r \phi^\mu \quad \text{con} \quad \phi^\mu = \int_1^{\lambda_0} d\lambda \langle \hat{T}^{\mu\nu} \rangle(\lambda) \beta_\nu$$

Entropy current

$$S = \int d\Sigma_\mu (\phi^\mu + T_{LE}^{\mu\nu} \beta_\nu - \zeta j_{LE}^\mu) \quad \text{but } n_p T^{\mu\nu} = n_p T_{LE}^{\mu\nu}$$
$$n_p j^\mu = n_p j_{LE}^\mu$$

$$S = \int d\Sigma_\mu (\phi^\mu + T^{\mu\nu} \beta_\nu - \zeta j^\mu)$$

$$\Rightarrow S^\mu = \phi^\mu + T^{\mu\nu} \beta_\nu - \zeta j^\mu$$

Entropy production

$$\frac{\delta}{\delta \Sigma} \left(\int d\Sigma_r V^\mu \right) = \frac{1}{2} \int d\tilde{S}_{\mu\nu} \xi^\nu V^\mu + \int d\Sigma_r \xi^\mu \nabla \cdot V$$

$$Z_{\phi_c(\Sigma)} - Z_\Sigma = \int_{\phi_c(\Sigma)} d\Sigma_r \phi^\mu - \int d\Sigma_r \phi^\mu \quad \text{ovvero} \quad \mathcal{L}_\xi \int d\Sigma_r \phi^\mu \text{ domain derivative} = \int d\Sigma_\mu \xi^\mu \nabla \cdot \phi \quad \text{e} \quad \int d\tilde{S} = 0$$

$$\begin{aligned} Z_{\phi_c(\varepsilon)} &= \text{Tr} \left(e^{- \int_{\phi_c(\Sigma)} d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu} \right) \cong \text{Tr} \left(e^{- \int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu - \varepsilon \int d\Sigma_r \xi^\mu \nabla \cdot (\hat{T}^{\mu\nu} \beta_\nu)} \right) \text{e} \quad \int d\tilde{S} = 0 \\ &\cong Z - \varepsilon \text{Tr} \left(e^{- \int d\Sigma \cdot \xi \nabla_\mu (\hat{T}^{\mu\nu} \beta_\nu)} \right) \end{aligned}$$

Ora dividendo per Z e ho

$$\lim_{\varepsilon \rightarrow 0} \frac{Z_\varepsilon - Z}{\varepsilon Z} = \mathcal{L}_\xi (\log Z) = - \int d\Sigma \cdot \xi \langle \hat{T}^{\mu\nu} \rangle_{\varepsilon} \nabla_\mu \beta_\nu \quad \text{e pertanto}$$

$$\mathcal{L}_\xi \int d\Sigma_r \phi^\mu = \int d\Sigma \cdot \xi \nabla \cdot \phi = - \int d\Sigma \cdot \xi \langle \hat{T}^{\mu\nu} \rangle_{\varepsilon} \nabla_\mu \beta_\nu$$

$$\text{being } \xi \text{ arbitrary} \Rightarrow \nabla \cdot \phi = - \langle \hat{T}^{\mu\nu} \rangle_{LE} \nabla_\mu \beta_\nu$$

$$\text{and, with current, } \nabla \cdot \phi = - \langle \hat{T}^{\mu\nu} \rangle_{LE} \nabla_\mu \beta_\nu + \langle \hat{j}^\mu \rangle_{LE} \nabla_\mu \xi$$

$$\Rightarrow \nabla \cdot s = \nabla \cdot (\phi + T^{\mu\nu} \beta_\nu - j^\mu) = \nabla \cdot \phi + \cancel{\nabla_\mu T^{\mu\nu}} \beta_\nu + T^{\mu\nu} \nabla_\mu \beta_\nu - \cancel{\nabla_\mu j^\mu} \xi$$

$$-\nabla \xi \cdot j = -T^{\mu\nu}_{LE} \nabla_\mu \beta_\nu + j^\mu_{LE} \nabla_\mu \xi + T^{\mu\nu} \nabla_\mu \beta_\nu - \cancel{\nabla_\mu \xi j^\mu} \Rightarrow$$

$$\nabla \cdot s = (T^{\mu\nu} - T^{\mu\nu}_{LE}) \nabla_\mu \beta_\nu - (j^\mu - j^\mu_{LE}) \nabla_\mu \xi$$

Comments on entropy

$$1) \quad S^r = \phi^r + T^{\mu\nu} \beta_\nu - \zeta j^\mu$$

is it independent of Σ (better: the foliation?)

No, because $\zeta(n)$ and $\beta(n)$. It can be made independent only if the foliation is fixed or, of course, at global equilibrium.

{
example : enforce $n = \hat{\beta}$

2) The relation $T_s + \mu n = \rho + p$ is not frame-independent.

At global equilibrium β is a Killing vector and entropy current is unique

$$\text{Landau frame} \quad S \cdot u_L = \phi \cdot u_L + u_{L\nu} T^{\mu\nu} \beta_\nu - \zeta j \cdot u_L$$

$$u_{L\nu} T^{\mu\nu} \beta_\nu = u_{L\nu} T^{\mu\nu} [(\beta \cdot u_L) u_{\mu\nu} + \beta_{T\nu}] = (\beta \cdot u_L) g_L$$


So we have

$$S \cdot u_L \equiv S_L = \phi \cdot u_L + \rho_L \frac{1}{T_L} - \zeta n_L$$

$$T_L = \frac{1}{\beta \cdot u_L} < \frac{1}{\sqrt{\beta^2}} \quad \beta \cdot u_L = \sqrt{\beta^2} \quad u \cdot u_L > \sqrt{\beta^2}$$

At global equilibrium with rotation we have $\beta = \frac{1}{T_0} (1, \underline{\omega} \times \underline{x})$
a Killing vector.

$$T^2 = \frac{T_0^2}{\sqrt{1 - \omega^2 r^2}} \quad \text{Tolman's law} \quad T_L^2 = \frac{1}{\beta^2} \frac{1}{(\hat{\beta} \cdot \hat{u}_L)^2} = T_0^2 \frac{1}{\sqrt{1 - \omega^2 r^2}} \cdot \frac{1}{(\hat{\beta} \cdot \hat{u}_L)^2}$$



At global equilibrium with rotation we have $\beta = \frac{1}{T_0} (1, \underline{\omega} \times \underline{x})$

a Killing vector.

However

$$u^\mu = \frac{\beta^\mu}{\sqrt{\beta^2}}, \quad \alpha^\mu = \varpi^{\mu\nu} u_\nu, \quad w^\mu = -\frac{1}{2} \epsilon^{\mu\nu\rho\sigma} \varpi_{\nu\rho} u_\sigma, \quad l^\mu = \epsilon^{\mu\nu\rho\sigma} w_\nu \alpha_\rho u_\sigma.$$

$$\begin{aligned} \langle :T^{\mu\nu}(x):\rangle = & \rho u^\mu u^\nu - p \Delta^{\mu\nu} + W w^\mu w^\nu + \mathcal{A} \alpha^\mu \alpha^\nu + G^l l^\mu l^\nu + G(l^\mu u^\nu + l^\nu u^\mu) + \mathbb{A}(\alpha^\mu u^\nu + \alpha^\nu u^\mu) \\ & + G^\alpha(l^\mu \alpha^\nu + l^\nu \alpha^\mu) + \mathbb{W}(w^\mu u^\nu + w^\nu u^\mu) + A^w(\alpha^\mu w^\nu + \alpha^\nu w^\mu) + G^w(l^\mu w^\nu + l^\nu w^\mu), \end{aligned}$$

$$T^{\mu\nu} u_\nu = \underbrace{\rho u^\mu + G l^\mu + \mathbb{A} \alpha^\mu + \mathbb{W} w^\mu}_{\text{heat flux? NO}}$$

$$T^{\mu\nu} u_\nu = \frac{\pi^2}{30} T^4 u^r + \frac{1-70\xi}{240\pi^2} T^4 l^r$$



$$u_L \neq \hat{\beta} !$$

free scalar field $m=0$

$$\rho = \frac{\pi^2}{30\beta^4} + \frac{4\xi-1}{12\beta^4} w^2 + \frac{6\xi-1}{12\beta^4} \alpha^2 + \frac{4\xi-1}{48\pi^2\beta^4} w^4 + \frac{60\xi-11}{480\pi^2\beta^4} \alpha^4 + \frac{270\xi-61}{720\pi^2\beta^4} \alpha^2 w^2,$$

$$p = \frac{\pi^2}{90\beta^4} - \frac{\xi}{6\beta^4} w^2 + \frac{1-6\xi}{18\beta^4} \alpha^2 - \frac{\xi}{24\pi^2\beta^4} w^4 + \frac{19-120\xi}{1440\pi^2\beta^4} \alpha^4,$$

$$W = \frac{2\xi-1}{12\beta^4} + \frac{2\xi-1}{48\pi^2\beta^4} w^2 + \frac{120\xi-29}{360\pi^2\beta^4} \alpha^2,$$

$$\mathcal{A} = \frac{1-6\xi}{12\beta^4} + \frac{1}{360\pi^2\beta^4} w^2 + \frac{1-6\xi}{48\pi^2\beta^4} \alpha^2,$$

$$G^l = \frac{1-70\xi}{240\pi^2\beta^4},$$

$$G = \frac{6\xi+1}{36\beta^4} + \frac{10\xi-1}{240\pi^2\beta^4} w^2 + \frac{30\xi-7}{720\pi^2\beta^4} \alpha^2,$$

$$\mathbb{A} = 0,$$

$$G^\alpha = 0,$$

$$G^w = A^w = \mathbb{W} = 0.$$

PARTICLES AND SPIN

$$[\alpha(\mathbf{p}), \alpha^\dagger(\mathbf{p}')] = 2\epsilon \delta^3(\mathbf{p}-\mathbf{p}')$$

Spectrum $\text{Tr}(\hat{\rho} \alpha^\dagger(\mathbf{p}) \alpha(\mathbf{p})) = 2\epsilon \frac{dN}{d^3 p}$

Spin $\frac{\text{Tr}(\hat{\rho} \alpha_r^\dagger(\mathbf{p}) \alpha_s(\mathbf{p}))}{\sum_r \text{Tr}(\hat{\rho} \alpha_r^\dagger(\mathbf{p}) \alpha_s(\mathbf{p}))} = \Theta_{rs}(\mathbf{p})$ Spin density matrix of a particle

Spin polarization vector

$$S^k(\mathbf{p}) = \sum_{rs}^S D(J_i)_{rs} \Theta_{sr}(\mathbf{p}) n_i^k(\mathbf{p}) = \sum_i \text{tr}(D^s(J_i) \Theta(\mathbf{p})) [p]_i^k$$

$$\hat{\rho} = \hat{\rho}_{LE}(0) = \frac{\exp \left[- \int_{F_0} d\sum_p (\hat{T}^{\mu\nu} \beta_\nu - S \hat{j}^\mu) + \int d\Omega \hat{T}^{\mu\nu} \nabla_\mu \beta_\nu \right]}{Z}$$

↓
 local equilibrium

↓
 dissipative

$\text{Tr}(\hat{\rho}_{\text{LE}}(\text{FO}) a^\dagger(p) a(p))$ how to expand β_v in the exponent?

$$\int d\Sigma_r \hat{T}^{\mu\nu} \beta_v \cong \beta_v(x) \hat{P}^\nu + \frac{1}{2} \varpi : \hat{J}_x + \frac{1}{2} \xi : \hat{Q}_x$$

$$\hat{J}_x^\mu = \int d\Sigma_\lambda \hat{T}^{\lambda\nu} x^\mu - (\mu \leftrightarrow \nu) \quad \hat{Q}_x^\mu = \int d\Sigma_\lambda \hat{T}^{\lambda\nu} x^\mu + (\mu \leftrightarrow \nu)$$

It is necessary to have a local operator!

Covariant Wigner Operator

Scalar $\widehat{W}(x, k) = \frac{2}{(2\pi)^4} \int d^4y : \widehat{\psi}^\dagger(x + y/2) \widehat{\psi}(x - y/2) : e^{-iy \cdot k}$

Dirac
$$\begin{aligned} \widehat{W}(x, k)_{AB} &= -\frac{1}{(2\pi)^4} \int d^4y e^{-ik \cdot y} : \Psi_A(x - y/2) \overline{\Psi}_B(x + y/2) : \\ &= \frac{1}{(2\pi)^4} \int d^4y e^{-ik \cdot y} : \overline{\Psi}_B(x + y/2) \Psi_A(x - y/2) : \end{aligned}$$

General idea: re-express $a^\dagger a$ as a function of $\hat{W}(x, k)$

$$W(x, k) = W(x, k)\theta(k^2)\theta(k^0) + W(x, k)\theta(k^2)\theta(-k^0) + W(x, k)\theta(-k^2) \equiv W_+(x, k) + W_-(x, k) + W_S(x, k)$$

\downarrow particle \downarrow antiparticle \searrow mixing

1) use field expansions in plane waves to expand \hat{W}

2) Show that $k^\mu \partial_\mu \hat{W} = 0 \Rightarrow \int_{\Sigma} d\Sigma_\mu k^\mu \hat{W}(x, k)$ independent of Σ
and k on-shell!

3) Show the relation

$$\frac{dN}{d^3 p} = \int dP_0 \int d\Sigma_\mu p^\mu W_+(x, p) \quad W_+(x, p) = \text{Tr}(\hat{\rho}_{LE}^{(0)} \hat{W}_+(x, p))$$

$$\frac{dN}{d^3 p} = \int dP_0 \int d\Sigma_\mu p^\mu \text{tr} W_+(x, p) \frac{1}{m}$$

Example

$$W(x, k) = \frac{2}{(2\pi)^3} \int \frac{d^3 p}{2\varepsilon} \frac{d^3 p'}{2\varepsilon'} e^{i(p-p') \cdot x} \left[\delta^4(k - (p+p')/2) \langle \hat{a}^\dagger(p) \hat{a}(p') \rangle + \delta^4(k + (p+p')/2) \langle \hat{b}^\dagger(p) \hat{b}(p') \rangle \right] \\ + \delta^4(k - (p-p')/2) \left[e^{i(p+p') \cdot x} \langle \hat{a}^\dagger(p) \hat{b}^\dagger(p') \rangle + e^{-i(p+p') \cdot x} \langle \hat{b}(p') \hat{a}(p) \rangle \right]$$

$$k^r \partial_r W_{+} = 0 \quad \Rightarrow \int d\Sigma_r k^r W_{+}(x, k) = \int d^3 x \ k^o W_{+}(x, k)$$

$$\int d^3 x \rightarrow (2\pi)^3 \delta^3(\underline{p} - \underline{p}') \Rightarrow$$

$$\int d\Sigma \cdot k W_{+}(x, k) = 2 \int \frac{d^3 p}{2\varepsilon} \frac{k^o}{2\varepsilon} \delta^4(k-p) \langle a^\dagger(p) a(p) \rangle = \int \frac{d^3 p}{2\varepsilon} \delta^3(\underline{p} - \underline{k}) \delta(k^o - p^o) \langle a^\dagger, a_p \rangle =$$

$$= \frac{1}{2\varepsilon} \delta(k^o - \sqrt{\underline{k}^2 + m^2}) \langle a_{\underline{k}}^\dagger a_{\underline{k}} \rangle = \delta(k^o - \sqrt{\underline{k}^2 + m^2}) \frac{dN}{d^3 k}$$

$$\Rightarrow \frac{dN}{d^3 k} = \int dk^o \int d\Sigma \cdot k W(x, k)$$

Spin: more elaborate

$$S^\mu(p) = \frac{1}{2} \frac{\int d\Sigma \cdot p \operatorname{tr}_4(\gamma^\mu \gamma^5 W_+(x, p))}{\int d\Sigma \cdot p \operatorname{tr}_4 W_+(x, p)}$$

Derivation:
F.B. Lect. Not. Phys. 987
(2021) 15

$$S^\mu(p) = -\frac{1}{4} \epsilon^{\mu\beta\gamma\delta} p_\delta \frac{\int d\Sigma_\lambda \operatorname{tr}_4(\{\gamma^\lambda, \Sigma_{\beta\gamma}\} W_+(x, p))}{\int d\Sigma_\lambda p^\lambda \operatorname{tr}_4 W_+(x, p)}$$

$$S^\mu(p) = -\frac{1}{2m} \epsilon^{\mu\beta\gamma\delta} p_\delta \frac{\int d\Sigma_\lambda p^\lambda \operatorname{tr}_4(\Sigma_{\beta\gamma} W_+(x, p))}{\int d\Sigma_\lambda p^\lambda \operatorname{tr}_4 W_+(x, p)}$$

Calculation at LE

$$\hat{g}_{LE}(0) \approx \hat{g}_{LE}(\Sigma_{FO}) \rightarrow \text{dissipation neglected}$$

$$W(x, k) = \text{Tr} \left(e^{-\frac{\int d\Sigma_r \hat{T}^{\mu\nu} \beta_\nu - \zeta \hat{j}^\mu}{Z}} \hat{W}_+(x, k) \right) \rightarrow \begin{matrix} \text{Taylor expand } \beta \\ \text{from } x \end{matrix}$$

$$\simeq \frac{\text{Tr}}{Z} \left(e^{-\beta(x) \cdot \hat{P}} + \frac{1}{2} \boldsymbol{\omega} : \hat{J}_x + \frac{1}{2} \boldsymbol{\xi} : \hat{Q}_x \hat{W}_+(x, k) \right)$$

Retaining the main term

$$\text{Tr} \left(e^{-\beta(x) \cdot \hat{P}} \hat{W}_+(x, k) \right) = \frac{2}{(2\pi)^3} \int \frac{d^3 p}{2\varepsilon} \frac{d^3 p'}{2\varepsilon'} e^{i(p-p') \cdot x} \delta^4(k - \frac{p+p'}{2}) \text{Tr} \left(e^{-\beta(x) \cdot \hat{P}} a_p^+ a_{p'}^- \right)$$

$$= \frac{2}{(2\pi)^3} \int \frac{d^3 p}{2\varepsilon} \frac{d^3 p'}{2\varepsilon'} e^{i(p-p') \cdot x} \delta^4(k - \frac{p+p'}{2}) \frac{2\varepsilon \delta^3(p-p')}{e^{\beta(x) \cdot p} - 1} = \frac{1}{\varepsilon_k} \frac{\delta(k^0 - \varepsilon_k)}{(2\pi)^3} n_B(\beta(x))$$

$$\Rightarrow \varepsilon \frac{dN}{d^3 p} = \int d\Sigma \cdot p \frac{1}{e^{\beta(x) \cdot p} - 1}$$

Spin

$$S^\mu(p) = \frac{1}{2} \frac{\int d\Sigma \cdot p \operatorname{tr}_4(\gamma^\mu \gamma^5 W_+(x, p))}{\int d\Sigma \cdot p \operatorname{tr}_4 W_+(x, p)}$$

We cannot stop at the leading order (0^{th} in the gradients) but include 1st order terms

$$\frac{\operatorname{Tr}}{\Sigma} \left(e^{-\beta(x) \cdot \hat{P}} + \frac{1}{2} \omega : \hat{J}_x + \frac{1}{2} \xi : \hat{Q}_x \hat{W}_+(x, k) \right)$$

For the detailed and most up-to-date derivation

see F.B., M. Buzzegoli, A. Palermo PLB 820 (2021) 136519

Two terms arise:

$$S_{\varpi}^{\mu}(p) = -\frac{1}{8m}\epsilon^{\mu\nu\rho\sigma}p_{\sigma}\frac{\int_{\Sigma} d\Sigma_{\lambda} p^{\lambda} n_F(1-n_F)\varpi_{\nu\rho}}{\int_{\Sigma} d\Sigma_{\lambda} p^{\lambda} n_F},$$

$$S_{\xi}^{\mu}(p) = -\frac{1}{4m}\epsilon^{\mu\nu\sigma\tau}\frac{p_{\tau}p^{\rho}}{\varepsilon}\frac{\int_{\Sigma} d\Sigma \cdot p n_F(1-n_F)\hat{t}_{\nu}\xi_{\sigma\rho}}{\int_{\Sigma} d\Sigma \cdot p n_F},$$

Also,