

Duality in interacting Bose gases

Fabian Essler (Oxford)

joint work with **Bruno Bertini** (Nottingham) and **Etienne Granet** (Quantinuum)



SFT 2024 - Lectures on Statistical Field Theories

Feb 05, 2024 - Feb 16, 2024

Apply (deadline: Nov 15, 2023)

Note: you are applying after the deadline



Integrable models are typically not “adiabatically connected” to free theories:

- Hierarchy of multi-particle bound states
- weakly interacting bosons behave like fermions at low E

Is there an integrable model where elementary excitations can be understood as in Fermi liquid by simple (perturbative) dressing of elementary fermions?

N-particle Hamiltonian:

$$H = \sum_{j=1}^N \left(-\frac{\partial^2}{\partial x_j^2} - \mu \right) + 2c \sum_{i < j} \delta(x_i - x_j)$$

2nd quantisation:

$$H = \int dx \, \Phi^\dagger(x) \left[-\partial_x^2 - \mu \right] \Phi(x) + c \int dx \, (\Phi^\dagger(x))^2 (\Phi(x))^2$$

$$[\Phi(x, t), \Phi^\dagger(y, t)] = \delta(x - y)$$

Consider the repulsive case $c > 0$.

Local conservation laws:

$$[Q_n, Q_m] = 0 = [H, Q_n]$$

Impenetrable bosons

“Impenetrable limit” $c \rightarrow \infty$: equivalent to free **fermions**

$$H(\infty) = - \int_0^L dx \Psi^\dagger(x) \partial_x^2 \Psi(x) \qquad \Psi^\dagger(x) = \Phi^\dagger(x) e^{i\pi \int_0^x dz \Phi^\dagger(z) \Phi(z)}$$

$$\{\Psi(x), \Psi^\dagger(y)\} = \delta(x - y)$$

2 sectors: periodic/anti-periodic bc's for odd/even fermion #

What about $c \gg 1$?

There should be a description in terms of **weakly interacting fermions**: adiabatic continuity with $c = \infty$ (dressing elementary fermions like in Fermi liquid)

Cheon-Shigehara model

Cheon&Shigehara '98

interacting bosons



interacting fermions

$$H_{\text{CS}}(a, c) = \sum_{j=1}^N \left[-\frac{\partial^2}{\partial x_j^2} - \mu \right] + 2 \sum_{i < j} \left[\frac{c}{2} - \frac{1}{a} \right] \left[\delta(x_j - x_k - a) + \delta(x_j - x_k + a) \right]$$

a is a **regulator** that needs to be taken to zero.

($a \rightarrow 0$: discontinuous wave-fns, continuous derivatives)

$$\lim_{a \rightarrow 0} \psi_{\text{CS}}(x_1, \dots, x_N) = \psi_{\text{LL}}(x_1, \dots, x_N) \prod_{j < k} \text{sgn}(x_j - x_k)$$

large c : looks like

strongly interacting bosons



strongly interacting fermions

No perturbative calculations possible.

Hm.

A different dual fermion theory

$$H(a, c) = \sum_{j=1}^N \left[-\frac{\partial^2}{\partial x_j^2} - \mu \right] + 2 \sum_{i < j} V_{a,c}(x_i - x_j)$$

Regularised interaction potential:

$$V_{a,c}(x) = \frac{2\sigma''\left(\frac{x}{a}\right)}{cx + 2\sigma\left(\frac{x}{a}\right)}$$

$\sigma(x)$ a smooth function s.t.

$$\sigma(-x) = -\sigma(x) , \quad \sigma'(0) > 0 , \quad \sigma'(x \neq 0) \geq 0$$

$$\lim_{x \rightarrow \infty} \sigma(x) = 1 , \quad \lim_{x \rightarrow \infty} x^2 \sigma''(x) = 0.$$


In the limit $a \rightarrow 0$ this is equivalent to $H_{\text{CS}}(a \rightarrow 0, c)$, but standard many-body perturbative calculations are possible!

Fermion spectral function at finite temperature


from fermion self-energy at order c^{-2}

$$\Sigma(\omega_n, k) = \text{diagram 1} + \text{diagram 2} + \text{diagram 3}$$

The diagrams represent the fermion self-energy at order c^{-2} . Diagram 1 is a tadpole diagram with a loop labeled ω_m, q . Diagram 2 is a bubble diagram with two internal loops labeled ω_{m_1}, q_1 and ω_{m_3}, q_3 , and an external loop labeled ω_{m_2}, q_2 . Diagram 3 is a more complex diagram with multiple loops and external lines labeled with ω_n, k and ω_m, q .



$$\mu = 1, c = 4, \beta = 0.5$$



$$\mu = 1, c = 4, \beta = 1$$

Free fermion acquires finite life-time.

Conserved charges for weakly interacting fermions

Consider $H = \sum_p \epsilon(p) \psi_p^\dagger \psi_p + \frac{\beta}{L} \sum_{\mathbf{p}_4} V(\mathbf{p}_4) \psi_{p_1}^\dagger \psi_{p_2}^\dagger \psi_{p_3} \psi_{p_4}$

(only quartic interactions)

Try to construct conserved charges order-by-order in β

$$\begin{aligned} Q^{(n)} = \int dx Q^{(n)}(x) &= \sum_p p^n \psi_p^\dagger \psi_p + \sum_{\mathbf{p}_4} S_1^{(n)}(\mathbf{p}_4) \psi_{p_1}^\dagger \psi_{p_2}^\dagger \psi_{p_3} \psi_{p_4} \\ &+ \sum_{\mathbf{p}_6} S_2^{(n)}(\mathbf{p}_6) \psi_{p_1}^\dagger \psi_{p_2}^\dagger \psi_{p_3}^\dagger \psi_{p_4} \psi_{p_5} \psi_{p_6} + \dots \end{aligned}$$

where $S_k^{(n)}$ has an expansion in β starting at β^k

Can determine $S_k^{(n)}$ recursively for any interaction potential!

???

Charges are generally **non-local** and become quasi-local only for special **integrable** $V(\mathbf{p}_4)$!

In the spinless fermion case we find at $\mathcal{O}(\beta^2)$:

1. Cheon-Shigehara model
2. Calogero-Sutherland model.

Precisely the known integrable models!

This construction is equivalent to

$$\widetilde{H} = e^{-iS} H e^{iS} , \quad S = \sum_{n \geq 1} \beta^n S_n ,$$

$$\widetilde{H} = \sum_{n \geq 1} \sum_{k_1 < \dots < k_n} h_n(k_1, \dots, k_n) \hat{n}(k_1) \hat{n}(k_2) \dots \hat{n}(k_n)$$

$$h_n(k_1, \dots, k_n) = \begin{cases} \mathcal{O}(\beta^{n-1}) & \text{if } H \text{ integrable} \\ \mathcal{O}(\beta^0) & \text{if } H \text{ not integrable} \end{cases}$$

$$\widetilde{H} \prod_{j=1}^N c^\dagger(k_j) |0\rangle = E(k) \prod_{j=1}^N c^\dagger(k_j) |0\rangle$$

Undo the unitary transformation: $A(I) = e^{iS} c^\dagger(k) e^{-iS} , \quad k = \frac{2\pi I}{L}$

fermionic creation operators for Bethe integers!

$$H \prod_{j=1}^N A^\dagger(I_j) |0\rangle = E(I) \prod_{j=1}^N A^\dagger(I_j) |0\rangle + \mathcal{O}(c^{-n})$$

Summary

1. Duality that maps strongly interacting Bose gas to weakly interacting fermions.
2. Perturbative analysis now feasible.
3. Generalizations to multi-species Yang-Gaudin models possible.
4. Operator equivalences ?
5. "Perturbative" construction of conserved charges to higher orders $\rightarrow$ models with interesting "pre thermal" dynamics