

Weyl anomalies in CFTs with boundaries and defects

Jacopo Sisti

Florence, 02/02/2024



UPPSALA
UNIVERSITET

Based on:

[Chalabi, Herzog, O'Bannon, Robinson, JS; JHEP 2022]

[Bianchi, Chalabi, Prochazka, Robinson, JS; JHEP 2021]

[Chalabi, Herzog, Ray, Robinson, JS, Stergiou; JHEP 2023]

“Physics is like a train which does not stop at the station. One first must run fast to catch that train”

IUTP-88/A054



Applied Conformal Field Theory

Paul Ginsparg[†]

Lyman Laboratory of Physics
Harvard University
Cambridge, MA 02138

Lectures given at Les Houches summer session, June 28 – Aug. 5, 1988.

To appear in Les Houches, Session XLIX, 1988, *Champs, Cordes et Phénomènes Critiques/ Fields, Strings and Critical Phenomena*, ed. by E. Brézin and J. Zinn-Justin, ©Elsevier Science Publishers B.V. (1989).



REVIEWS OF MODERN PHYSICS, VOLUME 83, OCTOBER–DECEMBER 2011

Topological insulators and superconductors

Xiao-Liang Qi

Microsoft Research, Station Q, Elings Hall, University of California,
Santa Barbara, California 93106, USA
and Department of Physics, Stanford University, Stanford, California 94305, USA

Shou-Cheng Zhang

Department of Physics, Stanford University, Stanford, California 94305, USA

(Received 2 August 2010; published 14 October 2011)

Topological insulators are new states of quantum matter which cannot be adiabatically connected to conventional insulators and semiconductors. They are characterized by a full insulating gap in the bulk and gapless edge or surface states which are protected by time-reversal symmetry. These topological materials have been theoretically predicted and experimentally observed in a variety of systems, including HgTe quantum wells, BiSb alloys, and Bi₂Te₃ and Bi₂Se₃ crystals. Theoretical models, material properties, and experimental results on two-dimensional and three-dimensional topological insulators are reviewed, and both the topological band theory and the topological field theory are discussed. Topological superconductors have a full pairing gap in the bulk and gapless surface states consisting of Majorana fermions. The theory of topological superconductors is reviewed, in close analogy to the theory of topological insulators.

DOI: 10.1103/RevModPhys.83.1057

PACS numbers: 73.20.-r, 73.43.-f, 85.75.-d, 74.90.+n

Defects

Local operators are not enough

- To distinguish groups with same algebra but different global structure
- To characterise phase of matter
- Strings and Branes

Some Examples:

- Boundaries (space-time ends on them)
- Interfaces (for instance between different phases)
- Line defects: for example Wilson lines
- Topological Defects (non-invertible symmetries)

Defects


Extended objects that break part of the translational symmetry

- Localised on submanifolds of the spacetime
- p =dimension of defect space-time, $q=d-p$ is the codimension
- The stress energy tensor is no longer conserved:

$$\partial_\mu T^{\mu\nu} = \hat{\mathcal{D}}^\nu \delta^q (\Sigma)$$

$\hat{\mathcal{D}}^\nu$ = displacement operator
protected dimension $p+1$

- We mainly focus on conformal field theories
- Conformal group is broken: $SO(d+1, 1) \longrightarrow SO(d+1-p, 1) \times SO(q)$
- New non-trivial correlators: **one-point functions**
- Localised contribution to anomalies: i.e. **defect Weyl anomalies**



Correlators in defect CFTs

[Billò, Gonçalves, Lauria, Meineri, 2016]

[Herzog, Shrestha, 2020]

Bulk correlators:

$$\langle \mathcal{O}(x) \rangle = \frac{A_{\mathcal{O}}}{|x|^{\Delta_{\mathcal{O}}}}$$


$$\langle T^{ij} \rangle = h \frac{(d - q + 1)\delta^{ij} - d \frac{x_{\perp}^i x_{\perp}^j}{|x_{\perp}|^2}}{|x_{\perp}|^d}$$

$$\langle T^{ab} \rangle = -h \frac{(q - 1)\delta^{ab}}{|x_{\perp}|^d}$$

i, j, k, \dots = orthogonal

a, b, c, \dots = parallel

$$\langle \mathcal{O}_1(x) \mathcal{O}_2(y) \rangle = \frac{1}{|x_{\perp}|^{\Delta_1} |y_{\perp}|^{\Delta_2}} f(\xi_1, \xi_2), \quad \xi_1 = \frac{s^2}{4|x_{\perp}| |y_{\perp}|}, \quad \xi_2 = \frac{x_{\perp} \cdot y_{\perp}}{|x_{\perp}| |y_{\perp}|}, \quad s = x - y$$



Bulk-defect correlators:

$$\left\langle \hat{\mathcal{O}}_{\hat{\Delta}}(\mathbf{x}) \mathcal{O}_{\Delta}(y) \right\rangle = \frac{C_{\hat{\mathcal{O}}\mathcal{O}}}{|y_{\perp}|^{\Delta - \hat{\Delta}} (y_{\perp}^2 + (\mathbf{x} - \mathbf{y})^2)^{\hat{\Delta}}}$$

Defect correlators:

$$\left\langle \hat{\mathcal{O}}_1(\mathbf{x}) \hat{\mathcal{O}}_2(\mathbf{y}) \right\rangle = \frac{\delta_{\Delta_{\hat{\mathcal{O}}_1}, \Delta_{\hat{\mathcal{O}}_2}}}{|\mathbf{x} - \mathbf{y}|^{2\Delta_{\hat{\mathcal{O}}_1}}}$$

$$\left\langle \hat{\mathcal{D}}^i(\mathbf{y}) \hat{\mathcal{D}}^j(\mathbf{0}) \right\rangle = \frac{C_{\hat{\mathcal{D}}\hat{\mathcal{D}}}}{|\mathbf{y}|^{2p+2}} \delta^{ij}$$

$$\left\langle \hat{\mathcal{O}}_1(\mathbf{x}_1) \hat{\mathcal{O}}_2(\mathbf{x}_2) \hat{\mathcal{O}}_3(\mathbf{x}_3) \right\rangle = \frac{C_{123}}{|\mathbf{x}_{12}|^{\Delta_1 + \Delta_2 - \Delta_3} |\mathbf{x}_{13}|^{\Delta_1 + \Delta_3 - \Delta_2} |\mathbf{x}_{23}|^{\Delta_2 + \Delta_3 - \Delta_1}}$$

(Bulk) Weyl Anomaly

- QFT on curved space-time
- Suppose the action invariant under Weyl rescaling (classical) $g_{\mu\nu} \rightarrow e^{2\omega(x)} g_{\mu\nu}$

$$\delta_\omega S = - \int_{\mathcal{M}_d} d^d x \sqrt{g} \delta\omega T^\mu{}_\mu = 0 \quad \longrightarrow \quad T^\mu{}_\mu = 0$$

- Effective action is not invariant: Weyl anomaly (quantum effect) $\langle T^{\mu\nu} \rangle = -\frac{2}{\sqrt{g}} \frac{\delta W}{\delta g_{\mu\nu}}$

$$\delta_\omega W = - \int_{\mathcal{M}_d} d^d x \sqrt{g} \delta\omega \langle T^\mu{}_\mu \rangle \neq 0$$

$$\longrightarrow \quad W = \frac{a_{d-2}}{\epsilon^{d-2}} + \frac{a_{d-4}}{\epsilon^{d-4}} \dots + \left[\int d^d x \sqrt{g} \langle T^\mu{}_\mu \rangle \right] \log \epsilon + \mathcal{O}(1)$$

- Local quantity of dimension d

$$T^\mu{}_\mu = \frac{1}{(4\pi)^{\frac{d}{2}}} \left[(-)^{\frac{d}{2}-1} a_{\mathcal{M}} E_d + \sum_n c_n I_n \right]$$

anomaly coefficients
 “central charges”  curvature invariants

- Bulk anomalies absent in odd dimensions

(Bulk) Weyl Anomaly

2 dimensions:

$$T^\mu{}_\mu = \frac{c}{24\pi} R$$

c = Virasoro central charge

[Zamolodchikov, 1986]

$$c_{\text{UV}} \geq c_{\text{IR}}$$

$$\langle T(x)T(0) \rangle = \frac{c}{|x|^4}$$

4 dimensions:

$$T^\mu{}_\mu = \frac{1}{16\pi^2} \left(-aE_4 + cW_{\mu\nu\rho\sigma}W^{\mu\nu\rho\sigma} \right)$$

$$a_{\text{UV}} \geq a_{\text{IR}}$$

[Cardy, 1988]

[Komargodski, Schwimmer, 2011]

$$\langle T_{\mu\nu}(x)T_{\rho\sigma}(0) \rangle \propto \frac{c}{|x|^4} I_{\mu\nu,\rho\sigma}(x)$$

6 dimensions:

$$T^\mu{}_\mu = \frac{1}{(4\pi)^3} (a E_6 + c_1 I_1 + c_2 I_2 + c_3 I_3)$$

$$a_{\text{UV}} \geq a_{\text{IR}} \quad (\text{conjectured})$$

$$I_1 = W_{\mu\lambda\rho\nu} W^{\lambda\sigma\tau\rho} W_\sigma{}^{\mu\nu}{}_\tau \quad I_2 = W_{\mu\nu}{}^{\lambda\rho} W_{\lambda\rho}{}^{\sigma\tau} W_{\sigma\tau}{}^{\mu\nu}$$

$$\langle T_{\mu\nu}(x)T_{\rho\sigma}(0) \rangle \propto \frac{c_3}{|x|^4} I_{\mu\nu,\rho\sigma}(x)$$

$$I_3 = W_{\mu\nu\lambda\rho} \left(D^2 \delta_\sigma^\nu - \frac{6}{5} R \delta_\sigma^\nu + 4 R_\sigma^\nu \right) W^{\sigma\nu\lambda\rho}$$

$$\langle TTT \rangle \propto c_1 + c_2$$

Defect Weyl Anomaly

- In the presence of a p-dimensional defect:

$$\delta W = -\frac{1}{2} \int_{\mathcal{M}_d} d^d x \sqrt{g} \delta g_{\mu\nu} \langle T^{\mu\nu} \rangle - \frac{1}{2} \int_{\Sigma_p} d^p y \sqrt{\bar{g}} \left(\delta g_{\mu\nu} \langle T^{\mu\nu} |_{\Sigma_p} \rangle + 2\delta X^i(y^a) \langle \mathcal{D}_i \rangle \right)$$


- Defect-localised contribution to Weyl anomaly

$$T^\mu{}_\mu = T^\mu{}_\mu|_{\mathcal{M}_d} + \delta^{(q)}(x_\perp) T^\mu{}_\mu|_{\Sigma_p} \quad D_\mu T^{\mu i} = \delta^{(q)}(x_\perp) \mathcal{D}^i$$

- For one-dimensional boundary is “trivial” $T^\mu{}_\mu|_{\Sigma_1} = \frac{c}{12\pi} K$

Defect Weyl Anomaly

$$\begin{aligned} e_a^\mu &= \partial_a X^\mu & X^\mu(Y_a) &\quad \text{embedding function} \\ \bar{g}_{ab} &= e_a^\mu e_b^\nu g_{\mu\nu} & &\quad (\text{induced metric}) \\ \Pi_{ab}^i &= \bar{D}_a e_b^\mu & &\quad (\text{second fundamental form}) \end{aligned}$$



- For two-dimensional defects [Henningson, Skenderis, 1999]
[Schwimmer, Theisen, 2008]

$$T^\mu{}_\mu|_{\Sigma_2} = \frac{1}{24\pi} (b \bar{R} + d_1 \mathring{\Pi}^2 + d_2 W^{ab}{}_{ab})$$

$$b_{\text{UV}} \geq b_{\text{IR}}$$

[Jensen, O'Bannon, 2015]

$$\langle \hat{\mathcal{D}}_i(\mathbf{y}) \hat{\mathcal{D}}_j(\mathbf{0}) \rangle \propto d_1 \frac{\delta_{ij}}{|\mathbf{y}|^6} \quad \langle T_{\mu\nu} \rangle \propto \frac{d_2}{|x_\perp|^d}$$

- For three-dimensional boundaries [Herzog, Huang, Jensen, 2015 - 2017]

$$T^\mu{}_\mu|_{\Sigma_3} = \frac{1}{16\pi^2} \left(a_{\mathcal{M}} E_4|_{\partial\mathcal{M}} + b_1 \mathring{K}^3 + b_2 \mathring{K}^{ab} W^c{}_{acb} \right)$$

$$E_4|_{\partial\mathcal{M}} = \delta_{def}^{abc} \left(2 K^d{}_a R^{ef}{}_{bc} + \frac{8}{3} K^d{}_a K^e{}_b K^f{}_c \right)$$

$$\langle \hat{\mathcal{D}} \hat{\mathcal{D}} \hat{\mathcal{D}} \rangle \propto b_1 \quad \langle \hat{\mathcal{D}}(\mathbf{y}) \hat{\mathcal{D}}(\mathbf{0}) \rangle \propto b_2 \frac{1}{|\mathbf{y}|^8}$$

4d Defect Weyl Anomaly

[Chalabi, Herzog, O'Bannon, Robinson, JS; JHEP 2022]

- Full Weyl anomaly for p=4 dimensional defects and codimension q>1

$$\begin{aligned}
 T^\mu_{\mu}|_{\Sigma_4} = & \frac{1}{(4\pi)^2} \left(-a_\Sigma \bar{E}_4 + d_1 \mathcal{J}_1 + d_2 \mathcal{J}_2 + d_3 W_{abcd} W^{abcd} + d_4 (W_{ab}{}^{ab})^2 \right. \\
 & + d_5 W_{aibj} W^{aibj} + d_6 W_{iab}^b W_c{}^{iac} + d_7 W_{ijkl} W^{ijkl} + d_8 W_{aijk} W^{aijk} \\
 & + d_9 W_{abjk} W^{abjk} + d_{10} W_{iabc} W^{iabc} + d_{11} W_{acb}^c W_d{}^{adb} + d_{12} W_{iaj}^a W_b{}^{ibj} \\
 & + d_{13} W_{ab}{}^{ab} \overset{\circ}{\Pi}_{cd}^i \overset{\circ}{\Pi}_i^{cd} + d_{14} W_{bij}^a \overset{\circ}{\Pi}_{ac}^i \overset{\circ}{\Pi}^{jbc} + d_{15} W_{ibj}^a \overset{\circ}{\Pi}_{ac}^i \overset{\circ}{\Pi}^{jbc} \\
 & + d_{16} W_{abcd} \overset{\circ}{\Pi}_{ac}^i \overset{\circ}{\Pi}_{ibd} + d_{17} W_a{}^{bac} \overset{\circ}{\Pi}_{bd}^i \overset{\circ}{\Pi}_{ic}^d + d_{18} W_{icj}^c \overset{\circ}{\Pi}_{ab}^i \overset{\circ}{\Pi}^{jab} \\
 & \left. + d_{19} \text{Tr } \overset{\circ}{\Pi}_i^i \overset{\circ}{\Pi}_i^j \overset{\circ}{\Pi}_j^j + d_{20} \text{Tr } \overset{\circ}{\Pi}^i \overset{\circ}{\Pi}^j \overset{\circ}{\Pi}_i \overset{\circ}{\Pi}_j + d_{21} (\text{Tr } \overset{\circ}{\Pi}^i \overset{\circ}{\Pi}_i)^2 + d_{22} (\text{Tr } \overset{\circ}{\Pi}^i \overset{\circ}{\Pi}_i) (\text{Tr } \overset{\circ}{\Pi}_i \overset{\circ}{\Pi}_j) \right)
 \end{aligned}$$

- 2 “non-trivial” invariants

$$\begin{aligned}
 \mathcal{J}_1 = & \frac{1}{d-1} R \overset{\circ}{\Pi}_{ab}^i \overset{\circ}{\Pi}_i^{ab} - \frac{1}{d-2} N^{\mu\nu} R_{\mu\nu} \overset{\circ}{\Pi}_{ab}^i \overset{\circ}{\Pi}_i^{ab} - \frac{2}{d-2} R_a{}^b \overset{\circ}{\Pi}_{ac}^i \overset{\circ}{\Pi}_i^{bc} - \frac{1}{2} W_{acb}^c \Pi_i \overset{\circ}{\Pi}^{iab} \\
 & + \frac{4}{9} W_{ica}^c \bar{D}^b \overset{\circ}{\Pi}_{ab}^i + \overset{\circ}{\Pi}^{iab} D_i W_{acb}^c - \frac{1}{2} \Pi^i \text{Tr } \overset{\circ}{\Pi}_i \overset{\circ}{\Pi}^j \overset{\circ}{\Pi}_j + \frac{1}{16} \Pi^i \Pi_i \text{Tr } \overset{\circ}{\Pi}^j \overset{\circ}{\Pi}_j \\
 & + \frac{2}{9} \bar{D}^b \overset{\circ}{\Pi}_{ab}^i \bar{D}^c \overset{\circ}{\Pi}_{ic}^a
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{J}_2 = & \frac{d-4}{d-2} W_{ab}{}^{ab} N^{\mu\nu} R_{\mu\nu} - \frac{d-4}{d-1} R W_{ab}{}^{ab} + \frac{4(d-5)}{3(d-2)} R_{ab} W_c{}^{acb} \\
 & - \frac{5(d-4)}{48} W_{ab}{}^{ab} \Pi^i \Pi_i + \frac{2(d-5)}{3} W_{ica}^c \bar{D}^b \overset{\circ}{\Pi}_{ab}^i + \frac{4(d+1)}{9} \overset{\circ}{\Pi}^{iab} D_i W_{acb}^c \\
 & - \frac{1}{3} W_{ic}{}^{ac} \bar{D}_a \Pi^i - \frac{2(d-5)}{3} \Pi^i \text{Tr } \overset{\circ}{\Pi}_i \overset{\circ}{\Pi}^j \overset{\circ}{\Pi}_j + \frac{(d-10)}{12} \Pi^i D_i W_{ab}{}^{ab} + \frac{1}{3} D^i D_i W_{ab}{}^{ab},
 \end{aligned}$$

Related to two-point function of displacement operator

Related to one-point function of the stress tensor

Defect Weyl Anomalies and Correlators

- Insertions of the displacement operator generate perturbation of the defect's shape
- This produces a variation in the effective action (at the second order)

$$\delta_X W = -\frac{1}{2} \int_{\Sigma_p} d^p y_1 d^p y_2 \langle \mathcal{D}_i(\mathbf{y}_1) \mathcal{D}_j(\mathbf{y}_2) \rangle \delta X^i(\mathbf{y}_1) \delta X^j(\mathbf{y}_2) + \mathcal{O}(\delta X^3)$$

$$\langle \hat{\mathcal{D}}_i(\mathbf{y}) \hat{\mathcal{D}}_j(0) \rangle = \delta_{ij} \frac{C_{\hat{\mathcal{D}}\hat{\mathcal{D}}}}{|\mathbf{y}|^{2p+2}}$$

- We Taylor expand and take only the term that diverges logarithmically
- From $\delta\Pi_{ab}^i = \partial_a \partial_b \delta X^i$ we find for p even (p odd no logs):

$$\delta_X W|_{\log \varepsilon} = (-1)^{p/2+1} \frac{2^{-(p+2)} \pi^{\frac{p}{2}}}{p! \Gamma(\frac{p}{2} + 2)} C_{\mathcal{D}\mathcal{D}} \int_{\Sigma_p} d^p y \partial^{a_1} \dots \partial^{a_{p/2-1}} \delta\Pi^{icd} \partial_{a_1} \dots \partial_{a_{p/2-1}} \delta\Pi_{cd}^i$$

- We find for p=2 and p=4:

$p = 2$

$$\langle \hat{\mathcal{D}}_i(\mathbf{y}) \hat{\mathcal{D}}_j(0) \rangle = \frac{4}{3\pi^2} \frac{d_1}{|\mathbf{y}|^6} \delta_{ij}$$

$$T^\mu{}_\mu|_{\Sigma_2} \supset \frac{d_1}{24\pi} \mathring{\Pi}^2$$

$p = 4$

$$\langle \mathcal{D}^i(\mathbf{y}) \mathcal{D}^j(0) \rangle = -\frac{72}{\pi^4} \frac{d_1}{|\mathbf{y}|^{10}} \delta^{ij}$$

$$T^\mu{}_\mu|_{\Sigma_4} \supset \frac{2}{9} d_1 \overline{D}^b \mathring{\Pi}_{ab}^i \overline{D}^c \mathring{\Pi}_{ic}^a$$

Unitarity

$$d_1 \leq 0$$

Defect Anomalies and Correlators

- Insertions of the stress-tensor produce (bulk) metric perturbation
- This also implies variation in the anomaly contribution

$$\delta_g W = -\frac{1}{2} \int d^d x \langle T^{\mu\nu} \rangle \delta g_{\mu\nu} \longrightarrow \delta_g \int d^d x \sqrt{g} \langle T^{\mu}_{\mu} \rangle = -\frac{1}{2} \int d^d x \langle T^{\mu\nu} \rangle \delta g_{\mu\nu} \Big|_{\log \varepsilon}$$

- At first order only one term contributes

$$\delta_g \int_{\Sigma_4} d^4 y \sqrt{\gamma} \left\langle T^{\mu}_{\mu} \Big|_{\Sigma_4} \right\rangle = \frac{d_2}{16\pi^2} \int_{\Sigma_4} d^4 y \frac{1}{3} \partial^k \partial_k \delta W^{ab}{}_{ab} \longrightarrow \langle T_{\mu\nu} \rangle \propto \frac{h}{|x_{\perp}|^d}$$

$$h = -\frac{\Gamma\left(\frac{q}{2} + 1\right)}{\pi^{\frac{q}{2}+2} (q+3)} d_2$$


- Averaged null-energy condition $\int_{-\infty}^{\infty} du \langle T_{\mu\nu} \rangle v^{\mu} v^{\nu} \geq 0$ implies a constraint

[Jensen, O'Bannon, Robinson, Rodgers, 2018]

[Chalabi, O'Bannon, Herzog, Robinson, JS, JHEP 2022]

- Null geodesics: $t = \ell u, \quad x_1 = \ell u \cos \psi, \quad x_5 = \ell u \sin \psi, \quad x_6 = \ell$

$$\int_{-\infty}^{\infty} du \langle T_{\mu\nu} \rangle v^{\mu} v^{\nu} = -d_2 \frac{q \Gamma\left(\frac{d+1}{2}\right) \Gamma\left(\frac{q}{2}\right)}{(q+3) \Gamma\left(\frac{d}{2}\right) \pi^{\frac{q+3}{2}} \ell^{d-2}} |\sin \psi| \geq 0 \longrightarrow d_2 \leq 0$$




Defect Anomalies and Entanglement Entropy

- Contribution to EE of a spherical region with a centred defect
 - In CFT for spherical regions we can use the CHM map
 - Two contributions:
 - [Kobayashi, Nishioka, Sato, Watanabe, 2018]
 - [Jensen, O'Bannon, Robinson, Rodgers, 2018]
- 1) Free-energy
- 2) Killing Energy

$$S_{A,\Sigma_2} = \frac{1}{3} \left(b + \frac{d-3}{d-1} d_2 \right) \log \left(\frac{L}{\varepsilon} \right)$$

[Chalabi, O'Bannon, Herzog, Robinson, JS; JHEP 2022]

$$S_{A,\Sigma_4} = -4 \left[a_\Sigma + \frac{1}{4} \frac{(d-5)(d-4)}{d-1} d_2 \right] \log \left(\frac{L}{\varepsilon} \right)$$



5-dimensional probe branes in AdS

$$I = -\frac{1}{16\pi G_N} \int dr d^d x \sqrt{g} \left(\mathcal{R} + \frac{d(d-1)}{L_{\text{AdS}}^2} \right) + T_{\text{br}} \int dr d^p y \sqrt{\bar{g}}$$

- Holographic result from the generalised Willmore Energy (probe brane) [Graham, Reichert, 2017]

- For $q > 1$

a_Σ	d_1	d_2	d_3	d_4	d_5	d_6	d_7	d_8	d_9	d_{10}	d_{11}
$\frac{1}{4}$	-1	$-\frac{1}{q}$	$\frac{1}{4}$	$\frac{1}{4}$	$\frac{2}{3q}$	$-\frac{(3q+14)}{9q}$	0	0	$-\frac{2}{3q}$	$-\frac{(q+2)}{6q}$	$-\frac{(3q+2)}{3q}$
d_{12}	d_{13}	d_{14}	d_{15}	d_{16}	d_{17}	d_{18}	d_{19}	d_{20}	d_{21}	d_{22}	-
$-\frac{2}{3q}$	$-\frac{1}{2}$	$\frac{2-q}{3q}$	$\frac{2(q+4)}{3q}$	$\frac{2(2q+1)}{3q}$	$\frac{(7q+6)}{3q}$	$-\frac{(q+6)}{3q}$	-1	$-\frac{1}{2}$	$\frac{1}{4}$	$\frac{1}{2}$	-

Monodromy defects

[Bianchi, Chalabi, Prochazka, Robinson, JS; JHEP 2021]

see also [Giombi et all, 2021]

- Co-dimension 2 operator which implements a flavour symmetry rotation

- Simplest case: U(1) rotation

$$\Psi(x) \rightarrow e^{-2\pi i \alpha} \Psi(x)$$

- In a Lagrangian theory it can be achieved by “gauging” the global symmetry by coupling to an external potential

$$S \rightarrow S + \int d^d x J^\mu A_\mu \quad A = \alpha d\theta$$

- This is a pure gauge everywhere but at the origin

$$F_{xy} = 2\pi \alpha \delta^{(2)}(x, y)$$

Anomaly coefficients for the free scalar:

d=4:

$$d_1 = d_2 = \frac{3}{2} [(1 - \alpha)^2 \alpha^2 + 4\xi \alpha^3]$$

$$b = \frac{(1 - \alpha)^2 \alpha^2 + 4\xi \alpha^3}{2}$$

$$\alpha \in (0, 1)$$

monodromy parameter

d=6:

$$d_1 = 2d_2 = -\frac{\alpha(1 - \alpha^2)(2 - \alpha)}{36} \left[\alpha(1 - \alpha) + \frac{6\alpha^2 \xi}{2 - \alpha} \right]$$


$$a_\Sigma = \frac{\alpha^2}{720} (1 - \alpha)^2 (3 + \alpha - \alpha^2) + \frac{\alpha^3}{360} (5 - 3\alpha^2) \xi$$

$$\xi = 0$$

IR fixed point

$$\xi = 1$$

UV fixed point



Monodromy defects

$\xi = 1$ UV fixed point



$$[\hat{O}_{-\alpha}(\sigma)] = \frac{d-2}{2} - \alpha$$

$\xi = 0$ IR fixed point




$$[\hat{O}_{\alpha}(\sigma)] = \frac{d-2}{2} + \alpha$$

- The two choices are connected by an RG flow:

$$S_{def} = \lambda \int d^{d-2} \sigma \hat{O}_{-\alpha}(\sigma) \hat{O}_{-\alpha}^{\dagger}(\sigma)$$

$$\lambda = \Lambda^{2\alpha} \bar{\lambda} \quad \text{relevant deformation}$$

$$\begin{aligned} \langle \hat{O}_{-\alpha}(\sigma) \hat{O}_{-\alpha}(0) \rangle_{\lambda} &\rightarrow \frac{1}{|\sigma|^{d-2-2\alpha}} \quad \sigma \rightarrow 0 \\ \langle \hat{O}_{-\alpha}(\sigma) \hat{O}_{-\alpha}(0) \rangle_{\lambda} &\rightarrow \frac{\Lambda^{-4\alpha}}{|\sigma|^{d-2+2\alpha}} \quad |\sigma| \rightarrow \infty \end{aligned}$$



- The Euler anomaly decreases

$$a_{\Sigma} = \frac{\alpha^2}{720} (1 - \alpha)^2 (3 + \alpha - \alpha^2) + \frac{\alpha^3}{360} (5 - 3\alpha^2) \xi$$



$$\boxed{a_{\Sigma}^{(\text{UV})} \geq a_{\Sigma}^{(\text{IR})}}$$

$$\xi = 1 \quad \xi = 0$$

defect a-theorem
[Wang, 2021]

- Analogous discussion for fermions

“Research in Physics is hard, so
you need to feel you are doing something epic”

Thank you for your attention!

and Happy Birthday Andrea!

Higher-Derivative Theories with Boundaries

[Chalabi, Herzog, Ray, Robinson, JS, Stergiou; JHEP 2023]

- Theories with an higher number of derivatives such as:

$$S = \int d^d x \phi \square^k \phi$$

$$\Delta_\phi = \frac{d - 2k}{2}$$

- Dimension of the fields is lower  Many relevant boundary deformations
- Violation of the unitarity bound $\Delta \geq \frac{d - 2}{2}$ for scalars

Boundary Primaries:

$$D |O_I\rangle = i\Delta |O_I\rangle ,$$

$$M_{ab} |O_I\rangle = (M_{ab})_I{}^J |O_J\rangle , \quad a, b = \text{parallel directions}$$

$$K_a |O_I\rangle = 0$$

- We need to compute the “single-trace” boundary primaries

$$|\Phi^{(k,q)}\rangle = \sum_{j=0}^{\lfloor \frac{q}{2} \rfloor} \alpha_j^{(k,q)} P_{\parallel}^{2j} P_n^{q-2j} |\phi\rangle$$




$$K_a |\Phi^{(k,q)}\rangle = 0 \quad a = \text{parallel directions}$$

Higher-Derivative Theories with Boundaries

- The first few boundary primaries are

$$\Phi^{(k,0)} = \phi, \quad \Phi^{(k,1)} = \partial_n \phi, \quad \Phi^{(k,2)} = \left(\partial_n^2 - \frac{1}{2k-3} \square_{\parallel} \right) \phi, \quad \Phi^{(k,3)} = \left(\partial_n^3 - \frac{3}{2k-5} \partial_n \square_{\parallel} \right) \phi$$

- For $k=1$ only 2 primaries:

$$\delta S_{\phi, \square}^{(k=1)} = \frac{1}{2} \int d^{d-1} x_{\parallel} (\Phi^{(1,0)} \delta \Phi^{(1,1)} - \Phi^{(1,1)} \delta \Phi^{(1,0)})$$


Neumann (N) $\Phi^{(k,1)} = \partial_n \phi = 0$
Dirichlet (D) $\Phi^{(k,0)} = \phi = 0$

- For $k=2$ we find 4 primaries:

$$\delta S_{\phi, \square}^{(k=2)} = \frac{1}{2} \int d^{d-1} x_{\parallel} (\Phi^{(2,0)} \delta \Phi^{(2,3)} - \Phi^{(2,1)} \delta \Phi^{(2,2)} + \Phi^{(2,2)} \delta \Phi^{(2,1)} - \Phi^{(2,3)} \delta \Phi^{(2,0)})$$

• NN

$$\Phi^{(2,2)} = (\partial_n^2 - \square_{\parallel}) \phi = 0$$

$$\Phi^{(2,3)} = (\partial_n^3 + 3\partial_n \square_{\parallel}) \phi = 0$$

• ND

$$\Phi^{(2,1)} = \partial_n \phi = 0$$

$$\Phi^{(2,3)} = \partial_n^3 \phi = 0$$

• DN

$$\Phi^{(2,0)} = \phi = 0$$


$$\Phi^{(2,2)} = \partial_n^2 \phi = 0$$

• DD


$$\Phi^{(2,0)} = \phi = 0$$

$$\Phi^{(2,1)} = \partial_n \phi = 0$$

Boundary RG flows



- In $d=5$
$$\Delta a_{\Sigma}(N \rightarrow D) = \frac{17}{5760}$$




- In $d=5$
$$\Delta a_{\Sigma}(NN \rightarrow DN) = \frac{21}{640}$$

$$\Delta a_{\Sigma}(NN \rightarrow ND) = -\frac{17}{5760}$$

$$\Delta a_{\Sigma}(DN \rightarrow DD) = -\frac{17}{5760}$$

$$\Delta a_{\Sigma}(ND \rightarrow DD) = \frac{21}{640}$$



- In $d=5$
$$\Delta a_{\Sigma}(\phi^2) = \frac{1375}{1152}$$

$$\Delta a_{\Sigma}((\partial_n \phi)^2) = \frac{21}{640}$$

$$\Delta a_{\Sigma}((\partial_n^2 \phi + \dots)^2) = -\frac{17}{5760}$$

Conclusion and Outlook

Summary

- We discussed the defect Weyl anomalies up to 4-dimensional defects
- Some of the anomaly coefficients are related to simple correlators in flat space-time
- Examples and computation of anomaly coefficients in free and holographic theories
- Characterisation of boundary condition for free-higher derivative theories

Outlook

- To relate additional anomaly coefficient to more complicated correlators
- Bounds on other anomaly coefficients
- Bulk theories coupled to lower dimensional one like minimal models for $p=2$
- Interacting higher-derivative theories with boundaries and defects?

Algorithm for finding the Weyl Anomalies

The algorithm (p=2, q=1 example):

- Find a basis of terms for the anomaly $\delta_\omega W$

$$\mathcal{B}_1 \delta\omega = \bar{R} \delta\omega ,$$

$$\mathcal{B}_2 \delta\omega = R \delta\omega ,$$

$$\mathcal{B}_3 \delta\omega = K^2 \delta\omega$$

$$\tilde{\mathcal{B}}_1 \delta\omega = \mathring{K}_{ab} \mathring{K}^{ab} \delta\omega$$

$$\mathcal{D}_1 \delta\omega = K D_\perp \delta\omega , \quad \mathcal{D}_2 \delta\omega = D_\perp^2 \delta\omega$$



$$\delta_\omega W = \int \sqrt{\bar{g}} \left[b_1 \mathcal{B}_1 + b_2 \mathcal{B}_2 + b_3 \mathcal{B}_3 + \tilde{b}_1 \tilde{\mathcal{B}}_1 + d_1 \mathcal{D}_1 + d_2 \mathcal{D}_2 \right] \delta\omega$$

Algorithm for finding the Weyl Anomalies

The algorithm (p=2, q=1 example):

- Find a basis of terms for the anomaly $\delta_\omega W$


$$\mathcal{B}_1 \delta\omega = \bar{R} \delta\omega,$$

$$\mathcal{B}_2 \delta\omega = R \delta\omega,$$

$$\mathcal{B}_3 \delta\omega = K^2 \delta\omega$$

$$\tilde{\mathcal{B}}_1 \delta\omega = \mathring{K}_{ab} \mathring{K}^{ab} \delta\omega$$

$$\mathcal{D}_1 \delta\omega = K D_\perp \delta\omega, \quad \mathcal{D}_2 \delta\omega = D_\perp^2 \delta\omega$$



$$\delta_\omega W = \int \sqrt{\bar{g}} \left[b_1 \mathcal{B}_1 + b_2 \mathcal{B}_2 + b_3 \mathcal{B}_3 + \tilde{b}_1 \tilde{\mathcal{B}}_1 + d_1 \mathcal{D}_1 + d_2 \mathcal{D}_2 \right] \delta\omega$$

- Impose Wess-Zumino consistency $[\delta_{\omega_1}, \delta_{\omega_2}] W = 0$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{B}_1 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{B}_2 \delta\omega_2) - (1 \leftrightarrow 2) \propto [\delta\omega_1 D_\perp^2 \delta\omega_2 - (1 \leftrightarrow 2)]$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{B}_3 \delta\omega_2) - (1 \leftrightarrow 2) \propto K [\delta\omega_1 D_\perp \delta\omega_2 - (1 \leftrightarrow 2)]$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \tilde{\mathcal{B}}_1 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

$$(\sqrt{\gamma})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{D}_1 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{D}_2 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

Algorithm for finding the Weyl Anomalies

The algorithm (p=2, q=1 example):

- Find a basis of terms for the anomaly $\delta_\omega W$

$$\mathcal{B}_1 \delta\omega = \bar{R} \delta\omega,$$

~~$$\mathcal{B}_2 \delta\omega = R \delta\omega,$$~~

~~$$\mathcal{B}_3 \delta\omega = K^2 \delta\omega$$~~

$$\tilde{\mathcal{B}}_1 \delta\omega = \mathring{K}_{ab} \mathring{K}^{ab} \delta\omega$$

$$\mathcal{D}_1 \delta\omega = K D_\perp \delta\omega, \quad \mathcal{D}_2 \delta\omega = D_\perp^2 \delta\omega$$

$$\delta_\omega W = \int \sqrt{\bar{g}} \left[b_1 \mathcal{B}_1 + \cancel{b_2 \mathcal{B}_2} + \cancel{b_3 \mathcal{B}_3} + \tilde{b}_1 \tilde{\mathcal{B}}_1 + d_1 \mathcal{D}_1 + d_2 \mathcal{D}_2 \right] \delta\omega$$

- Impose Wess-Zumino consistency $[\delta_{\omega_1}, \delta_{\omega_2}] W = 0$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{B}_1 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$



$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \tilde{\mathcal{B}}_1 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{B}_2 \delta\omega_2) - (1 \leftrightarrow 2) \propto [\delta\omega_1 D_\perp^2 \delta\omega_2 - (1 \leftrightarrow 2)]$$



$$(\sqrt{\gamma})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{D}_1 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{B}_3 \delta\omega_2) - (1 \leftrightarrow 2) \propto K [\delta\omega_1 D_\perp \delta\omega_2 - (1 \leftrightarrow 2)]$$

$$(\sqrt{\bar{g}})^{-1} \delta_1 (\sqrt{\bar{g}} \mathcal{D}_2 \delta\omega_2) - (1 \leftrightarrow 2) = 0$$

Algorithm for finding the Weyl Anomalies

The algorithm (p=2, q=1 example):

- Find a basis of terms for the anomaly $\delta_\omega W$

$$\mathcal{B}_1 \delta\omega = \bar{R} \delta\omega ,$$

~~$$\mathcal{B}_2 \delta\omega = R \delta\omega ,$$~~

~~$$\mathcal{B}_3 \delta\omega = K^2 \delta\omega$$~~

$$\tilde{\mathcal{B}}_1 \delta\omega = \mathring{K}_{ab} \mathring{K}^{ab} \delta\omega$$

$$\mathcal{D}_1 \delta\omega = K D_\perp \delta\omega , \quad \mathcal{D}_2 \delta\omega = D_\perp^2 \delta\omega$$

$$\delta_\omega W = \int \sqrt{\bar{g}} \left[b_1 \mathcal{B}_1 + \cancel{b_2 \mathcal{B}_2} + \cancel{b_3 \mathcal{B}_3} + \tilde{b}_1 \tilde{\mathcal{B}}_1 + d_1 \mathcal{D}_1 + d_2 \mathcal{D}_2 \right] \delta\omega$$

- Add all the possible counter-terms

$$W_{CT} = \sum_{i=1}^3 \int \sqrt{\gamma} c_i \mathcal{B}_i$$



$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_1) = 0$$

$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_2) \propto D_\perp^2 \delta\omega$$

$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_3) \propto K D_\perp \delta\omega$$

Algorithm for finding the Weyl Anomalies

The algorithm (p=2, q=1 example):

- Find a basis of terms for the anomaly $\delta_\omega W$

$$\mathcal{B}_1 \delta\omega = \bar{R} \delta\omega ,$$

~~$$\mathcal{B}_2 \delta\omega = R \delta\omega ,$$~~

~~$$\mathcal{B}_3 \delta\omega = K^2 \delta\omega$$~~

$$\tilde{\mathcal{B}}_1 \delta\omega = \mathring{K}_{ab} \mathring{K}^{ab} \delta\omega$$

~~$$\mathcal{D}_1 \delta\omega = K D_\perp \delta\omega ,$$~~

~~$$\mathcal{D}_2 \delta\omega = D_\perp^2 \delta\omega$$~~

$$\delta_\omega W = \int \sqrt{\bar{g}} \left[b_1 \mathcal{B}_1 + \cancel{b_2 \mathcal{B}_2} + \cancel{b_3 \mathcal{B}_3} + \tilde{b}_1 \tilde{\mathcal{B}}_1 + \cancel{d_1 \mathcal{D}_1} + \cancel{d_2 \mathcal{D}_2} \right] \delta\omega$$

- Add all the possible counter-terms

$$W_{CT} = \sum_{i=1}^3 \int \sqrt{\gamma} c_i \mathcal{B}_i$$



$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_1) = 0$$

$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_2) \propto D_\perp^2 \delta\omega$$



$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_3) \propto K D_\perp \delta\omega$$



Algorithm for finding the Weyl Anomalies

The algorithm (p=2, q=1 example):

- Find a basis of terms for the anomaly $\delta_\omega W$

$$\mathcal{B}_1 \delta\omega = \bar{R} \delta\omega, \quad \cancel{\mathcal{B}_2 \delta\omega = R \delta\omega}, \quad \cancel{\mathcal{B}_3 \delta\omega = K^2 \delta\omega}$$

$$\tilde{\mathcal{B}}_1 \delta\omega = \mathring{K}_{ab} \mathring{K}^{ab} \delta\omega$$

$$\cancel{\mathcal{D}_1 \delta\omega = K D_\perp \delta\omega}, \quad \cancel{\mathcal{D}_2 \delta\omega = D_\perp^2 \delta\omega}$$

$$\delta_\omega W = \int \sqrt{\bar{g}} \left[b_1 \mathcal{B}_1 + \cancel{b_2 \mathcal{B}_2} + \cancel{b_3 \mathcal{B}_3} + \tilde{b}_1 \tilde{\mathcal{B}}_1 + \cancel{d_1 \mathcal{D}_1} + \cancel{d_2 \mathcal{D}_2} \right] \delta\omega$$

- Add all the possible counter-terms

$$W_{CT} = \sum_{i=1}^3 \int \sqrt{\gamma} c_i \mathcal{B}_i$$



$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_1) = 0$$

$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_2) \propto D_\perp^2 \delta\omega$$



$$\delta_\omega(\sqrt{\bar{g}} \mathcal{B}_3) \propto K D_\perp \delta\omega$$



$$\mathcal{A} = \int \sqrt{\gamma} \left[-b \bar{R} + \mathring{K}_{ab} \mathring{K}^{ab} \right]$$

Boundary Weyl Anomaly (q=1)

- We can specify our general result to q=1. We recover [Astaneh, Solodukhin, 2021]

$$T^\mu_{\mu}|_{\Sigma_4} = \frac{1}{(4\pi)^2} \left(-a_\Sigma \bar{E}_4 + b_1 \mathcal{I} + b_2 (\text{Tr } \mathring{K}^2)^2 + b_3 \text{Tr } \mathring{K}^4 + b_4 W_{abcd} W^{abcd} + b_5 W_{anbn} W^a_n {}^b_n \right. \\ \left. + b_6 W_{abcd} \mathring{K}^{ac} \mathring{K}^{bd} + b_7 W_{anbn} \mathring{K}^a_c \mathring{K}^{cb} + b_8 W_{nabc} W_n{}^{abc} \right)$$

- Also in this case, the term \mathcal{I} is related to the displacement two-point function

$$\mathcal{I} = -\frac{2}{3} R_{ab} \mathring{K}^a_c \mathring{K}^{cb} + \frac{1}{4} R \text{Tr } \mathring{K}^2 - \frac{1}{3} R_{nn} \text{Tr } \mathring{K}^2 + \frac{1}{2} W_{anbn} K \mathring{K}^{ab} + \frac{1}{16} K^2 \text{Tr } \mathring{K}^2 \\ + \mathring{K}^{ab} D_n W_{anbn} - \frac{1}{2} K \text{Tr } \mathring{K}^3 + \frac{2}{9} \bar{D}^a \mathring{K}_{ab} \bar{D}_c \mathring{K}^{bc}$$



$$b_1 \leq 0$$

- Boundary anomaly coefficients for 5d probe branes

a_Σ	b_1	b_2	b_3	b_4	b_5	b_6	b_7	b_8
$\frac{1}{4}$	-1	$\frac{3}{4}$	$-\frac{3}{2}$	$\frac{1}{4}$	-1	2	-1	$-\frac{1}{2}$


- Holographic description: boundary in the space-time

$$S_{\text{grav}} = -\frac{1}{16\pi G_N} \int_{\mathcal{N}} \sqrt{\mathfrak{g}} (\mathcal{R} - 2\Lambda) - \frac{1}{8\pi G_N} \int_{\mathcal{Q}} \sqrt{\bar{\mathfrak{g}}} (\mathcal{K} - T) - \frac{1}{8\pi G_N} \int_{\mathcal{M}_d} \sqrt{\bar{\mathfrak{g}}} \mathcal{K}$$

Einstein equations + (Neumann) boundary condition

$$R_{MN} - \frac{1}{2} \mathcal{R} \mathfrak{g}_{MN} + \Lambda \mathfrak{g}_{MN} = 0 \quad \mathcal{K}_{MN} = (\mathcal{K} - T) \bar{\mathfrak{g}}_{MN}$$


- Conformal solutions



$$ds^2 = \frac{L^2}{z^2} (dz^2 + dy^2 + dx_\perp^2)$$

$$x_\perp = (\cot \theta) z$$

$$T = -\frac{d-1}{L} \cos \theta$$




$$ds^2 = \frac{L^2}{z^2} (dz^2 + dy^2 + dx_\perp^2)$$

$$\rho_{\mathcal{Q}}(z) \equiv \sqrt{\frac{L^2}{\sin^2 \theta} - (z + L \cot \theta)^2}$$

- (Half-)sphere free energy

$$S_{\text{grav}} = \frac{\pi L^4}{6G_N} \left[-\frac{1 + 2 \sin^2 \theta}{\sin^2 \theta} \cot \theta \log \left(\frac{L}{\epsilon} \right) + \dots \right]$$

$$a_\Sigma = -\frac{\pi L^4}{24G_N} \frac{1 + 2 \sin^2 \theta}{\sin^2 \theta} \cot \theta$$



Boundary RG flow

$$\theta_{\text{UV}} \geq \theta_{\text{IR}}$$



$$a_\Sigma^{(\text{UV})} \geq a_\Sigma^{(\text{IR})}$$


- Metric perturbation about AdS (back-reaction)

$$\begin{aligned} ds^2 = & \frac{L_{\text{AdS}}^2}{z^2} \left[dz^2 + \left(1 + \epsilon^2 x_\perp^2 \mathcal{G}^{(2)}(u) + \epsilon^3 x_\perp^3 \mathcal{G}^{(3)}(u) \right) dx_\perp^2 + \right. \\ & \left. + \left(\delta_{ab} + \epsilon x_\perp \mathcal{F}_{ab}^{(1)}(u) + \epsilon^2 x_\perp^2 \mathcal{F}_{ab}^{(2)}(u) + \epsilon^3 x_\perp^3 \mathcal{F}_{ab}^{(3)}(u) \right) dy^a dy^b \right] + \mathcal{O}(\epsilon^4) \end{aligned}$$

- Embedding for Q

$$u_Q(x) = \tan \theta + \epsilon B^{(1)} x_\perp + \epsilon^2 B^{(2)} x_\perp^2 + \epsilon^3 B^{(3)} x_\perp^3 + \mathcal{O}(\epsilon^4)$$

- We can solve the system of equation order by order in ϵ



AdS/BCFT

[Chalabi, Herzog, O'Bannon, Robinson, JS, 2021]

- Solution up to third order in metric perturbation
- Values of the anomaly coefficients EXACT in θ


$$b_1 = -\frac{1}{3} \frac{1}{\theta - \sin \theta \cos \theta} \frac{\pi L^4}{G_N} \quad b_1 \leq 0$$

$$2b_4 + b_5 = \frac{1}{4} \frac{1}{\tan \theta - \theta} \frac{\pi L^4}{G_N}$$

$$b_6 + b_7 = \frac{1}{24} \frac{13 \sin \theta - 3 \sin(3\theta) - 4\theta \cos \theta}{(\theta - \sin \theta \cos \theta)(\sin \theta - \theta \cos \theta)} \frac{\pi L^4}{G_N}$$

$$\frac{b_3}{2} - 2b_4 + b_6 = \frac{1}{48} \frac{13 - 3 \cos(2\theta) - 10\theta \cot \theta}{(1 - \theta \cot \theta)(\theta - \sin \theta \cos \theta)} \frac{\pi L_{\text{AdS}}^4}{G_N}$$

$$\begin{aligned} b_2 + \frac{7}{12}b_3 &= \frac{1}{93312} \frac{1}{(\sin(2\theta) - 2\theta)^4 (\theta \cos \theta - \sin \theta)} \times \\ &\times \left[12 (7963 + 16996\theta^2 - 6144\theta^4) \cos \theta - 24(5023 + 36\theta^2) \cos(3\theta) \right. \\ &+ 8(3719 + 4644\theta^2) \cos(5\theta) - 4699 \cos(7\theta) - 57 \cos(9\theta) \\ &+ 24\theta(-13933 + 1488\theta^2) \sin \theta + 144\theta(600\theta^2 - 421) \sin(3\theta) \\ &\left. + 21968\theta \sin(5\theta) - 11131\theta \sin(7\theta) + 429\theta \sin(9\theta) \right] \frac{\pi L_{\text{AdS}}^4}{G_N} \end{aligned}$$



Scale vs Conformal Invariance

- Example of scale (and Lorentz invariant) but not conformal boundary conditions

$$\frac{dP^a}{dt} = \int_{t,z=0} d^{d-2}x [T^{za} + \partial_t \tau^{at}]$$

$$\frac{dD}{dt} = \int_{t,z=0} d^{d-2}x [x_a T^{az} + \partial_t (x_a \tau^{at})]$$

$$\frac{dK^a}{dt} = \int_{t,z=0} d^{d-2}x [(2x^a x_b - x^2 \delta_b^a) T^{bz} + \partial_t (2x^t x_b - x^2 \delta_b^t) \tau^{bt}]$$

- If we take the free and not supported bc in plate theory we have to set to zero the following

$$\tilde{\Phi}^{(2,2)} \equiv (\partial_n^2 + \sigma \square_{\parallel})\phi, \quad \tilde{\Phi}^{(2,3)} \equiv (\partial_n^3 + (2 - \sigma)\partial_n \square_{\parallel})\phi$$

- This gives $T^{az} = -\partial_b \tau^{ab} \neq 0$

- Boundary scale invariance requires $\tau_a^a = \partial^a j_a$



$\sigma = -1$ conformal
 $j_a = 0$

$\sigma \neq -1$ non conformal
 $j_a \neq 0$

- It is achieved by deforming the boundary action with

$$S_{\text{marg}} = \sigma \int d^{d-1}x \phi \partial_{\parallel}^2 \partial_{\perp} \phi$$