




Prime Suspects

Quantum Tales in Number Theory

Giuseppe Mussardo

SISSA-INFN

Trieste



Magic of Many (Real and Imaginary) Paths





MODULAR INVARIANT PARTITION FUNCTIONS IN TWO DIMENSIONS

A. CAPPELLI*, C. ITZYKSON and J.-B. ZUBER

Service de Physique Théorique, CEN-Saclay, 91191 Gif-sur-Yvette Cedex, France

Received 3 September 1986

We present a systematic study of modular invariance of partition functions, relevant both for two-dimensional minimal conformal invariant theories and for string propagation on a $SU(2)$ group manifold. We conjecture that all solutions are labelled by simply laced Lie algebras.

1. Introduction

The minimal two-dimensional conformal invariant field theories [1] carry a set of representations of two Virasoro algebras of common central charge

$$c = 1 - \frac{6(p - p')^2}{pp'}, \quad (1.1)$$


with (p, p') a pair of coprime positive integers. Belavin, Polyakov and Zamolodchikov have shown that it is consistent to retain only a finite number of primary fields $\phi_{h, \bar{h}}$, of conformal dimensions h and $\bar{h}$ chosen among the Kac values [2]

$$h_{rs} = \frac{(rp - sp')^2 - (p - p')^2}{4pp'} = h_{p' - r, p - s}, \quad (1.2a)$$

with

$$1 \leq r \leq p' - 1, \quad 1 \leq s \leq p - 1. \quad (1.2b)$$

Paths of life









I L
DECAMERON

DI MESSER
GIOVANNI BOCCACCI
Cittadino Fiorentino.

Ricorretto in Roma, et Emendato secondo
l'ordine del Sacro Conc. di Trento,

*Et riscontrato in Firenze con Testi Antichi & alla sua
vera lezione ridotto da' Deputati di loro Alt. Ser.*

NVOVAMENTE STAMPATO.

*Con Privilegi del Sommo Pontefice, delle Maestadi del Re Christianissimo &
Re Cattolico, delli Serenissimi Gran Duca & Principe di Toscana,
dell' Ill. et Ecc. S. Duca di Ferrara, et d' altri Sign. et Rep.*



IN FIORENZA
Nella Stamperia de i Giunti
MDLXXIII.





SPACCIO
DI VINI
E
LIQUORI

Mosca



Le Guide
routard



Brasile




Le Routard



Le Routard








Statistical Field Theories

Edited by

Andrea Cappelli and Giuseppe Mussardo

NATO Science Series

II. Mathematics, Physics and Chemistry – Vol. 73



Statistical Field Theory





*An Introduction to Exactly Solved Models
in Statistical Physics*


Second Edition

Giuseppe Mussardo

OXFORD GRADUATE TEXTS

There is an increasing interest for the profound and engaging links recently discovered between Number Theory and Physics





$$\nu_{n,m} = E_n - E_m$$

Natural questions

- Given an arithmetic sequence $\{S_n\}$, does exist a quantum mechanics system which has this sequence as a spectrum?
- Is the Hamiltonian of such a system unique?


Our attention is on one-dimensional and one-body
Hamiltonians of the form

$$H = \frac{p^2}{2m} + V(x)$$

Equivalently, on tridiagonal Hamiltonians of the form



$$\begin{pmatrix} v_1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -1 & v_2 & -1 & 0 & 0 & 0 & 0 & 0 \\ 0 & -1 & v_3 & -1 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & v_4 & -1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 & v_5 & -1 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & v_k & -1 & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 & v_m & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & -1 & v_n \end{pmatrix}$$

Equivalently, on tridiagonal Hamiltonians of the form




Why one-body Hamiltonians?

- These Hamiltonians possess **discrete** spectrum
- Many-body Hamiltonians on the other hands, have **dense** spectrum



But, we can use nevertheless many-body Hamiltonians
to encode interesting number sequences!

- Instead of using energy levels, use wave functions!



$$n_i = \{S_n\}$$


Prime Suspects in a Quantum Ladder

Giuseppe Mussardo^{1,*}, Andrea Trombettoni^{2,1} and Zhao Zhang^{3,4}

¹SISSA and INFN, Sezione di Trieste, via Beirut 2/4, I-34151 Trieste, Italy

²Department of Physics, University of Trieste, Strada Costiera 11, I-34151 Trieste, Italy

³Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanghai 200240, China

⁴Nordita, KTH Royal Institute of Technology and Stockholm University, Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden



(Received 10 May 2020; accepted 6 November 2020; published 9 December 2020)

In this Letter we set up a suggestive number theory interpretation of a quantum ladder system made of $\mathcal{N}$ coupled chains of spin 1/2. Using the hard-core boson representation and a leg-Hamiltonian made of a magnetic field and a hopping term, we can associate to the spins σ_a the prime numbers p_a so that the chains become quantum registers for square-free integers. The rung Hamiltonian involves permutation terms between next-neighbor chains and a coprime repulsive interaction. The system has various phases; in particular, there is one whose ground state is a coherent superposition of the first $\mathcal{N}$ prime numbers. We also discuss the realization of such a model in terms of an open quantum system with a dissipative Lindblad dynamics.

DOI: 10.1103/PhysRevLett.125.240603

Introduction.—The aim of this Letter is to point out some interesting connections between quantum many-body systems and number theory, in particular, prime numbers. Prime numbers are the building blocks of arithmetics and, arguably, one of the pillars of the entire mathematics [1,2]. Their nature has two fascinating but opposite features [3]: If their appearance in the sequence of natural numbers is rather unpredictable, their coarse-graining properties [e.g., their total number $\pi(x)$ less than x] can be captured instead rather efficiently by simple statistical considerations [4–8]. In particular, the scaling of the k th prime is particularly plain:

$$p_k \simeq k \log k. \quad (1)$$


Equally fascinating is the connection between prime numbers and quantum mechanics: Prime numbers, for instance, were the main concern of Shor's algorithm, one of the first quantum computing algorithms [9]. Moreover, the scaling behavior (1) permits one to show the existence of a single-particle one-dimensional quantum mechanical potential $V(x)$ with eigenvalues given just by the prime numbers and, therefore, permits one to address the primality test of a natural number in terms of a quantum scattering [10]. Such a potential $V(x)$ can be determined either semiclassically [10] or exactly, using in this case methods of supersymmetric quantum mechanics [11,12]. In experimental setups of cold atom systems, $V(x)$ could be realized using a holographic trap [13].

Turning now our attention to quantum many-body systems, for the dense nature of their spectra it is obviously impossible to have energy levels given by prime numbers, but we can have instead many-body ground state wave

functions expressed in terms of prime numbers. This is what we are going to present below, where we consider a quantum ladder system with a suggestive number theoretic interpretation. We will see that such a system has a rich spectrum of ground states and, in particular, there is one whose wave function is given in terms of a highly coherent superposition of prime number occupations. To the best of our knowledge, this is the first time where a ground state of this type has been constructed.

Quantum ladder systems, made of coupled one-dimensional chains, have attracted considerable interest in recent years as truly interpolating between one- and two-dimensional systems [14–20]. In our case, we have $\mathcal{N}$ coupled half-infinite chains of spins 1/2 subjected to a magnetic field and a hopping term. As discussed below, properly tuning these two interactions, we can put in correspondence the spins with the prime numbers and reformulate the spin-spin rung interaction in terms of coprimality conditions (two integers are coprime if they do not share common factors other than 1).

Degrees of freedom.—As it is well known, spin 1/2 can be described by hard-core bosons: The mapping between the Pauli matrices σ_a and the hard-core annihilation and creation operators f and $f^\dagger$ [$f^2 = (f^\dagger)^2 = 0$] is provided by $\sigma_z = f^\dagger f - 1/2$; $\sigma_+ = f^\dagger$; $\sigma_- = f$ [21]. Hence, instead of the spins, we can equivalently take as degrees of freedom the hard-core boson operators $f_i(a)$ and $f_i^\dagger(a)$, where the index i refers to the i th chain ($i = 1, 2, \dots, \mathcal{N}$), while $a = 1, 2, \dots$ to the vertical position along the half-infinite chain (see Fig. 1). Since $[f_i(a)]^2 = [f_i^\dagger(a)]^2 = 0$, the occupation number of each vertical site in the ladder can take only values $\{0, 1\}$. Let $|\text{vac}\rangle$ be the vacuum state, i.e.,




Natural questions

- Given an arithmetic sequence $\{S_n\}$, does exist a quantum mechanics system which has this sequence as a spectrum?
- Is the Hamiltonian of such a system unique?

Fibonacci numbers

$$F_{n+2} = F_{n+1} + F_n$$

$$\{F_n\} = 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$$



$$|\Psi_f\rangle = M^{-1} |\Psi_i\rangle$$

Fibonacci numbers

$$F_{n+2} = F_{n+1} + F_n$$

$$\{F_n\} = 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$$

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^n = \begin{pmatrix} F_{n+1} & F_n \\ F_n & F_{n-1} \end{pmatrix}$$

$$F_{n+2} = F_{n+1} + F_n$$

$$F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

Fibonacci numbers

$$F_{n+2} = F_{n+1} + F_n$$

$$\{F_n\} = 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, \dots$$

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^n = \begin{pmatrix} F_{n+1} & F_n \\ F_n & F_{n-1} \end{pmatrix}$$

$$F_n = \frac{1}{\sqrt{5}} \left[\left(\frac{1 + \sqrt{5}}{2} \right)^n - \left(\frac{1 - \sqrt{5}}{2} \right)^n \right]$$

- Unfortunately, it does not exist a quantum system which has the Fibonacci numbers as spectrum...
- The reason is that their sequence grows too fast
- Similarly, it does not exist a quantum Schroedinger Hamiltonian with a spectrum given, for instance, by the Mersenne numbers or the perfect numbers

$$M_n = 2^n - 1$$

$$P_n = 2^{n-1} M_n$$


Bound on the growth of eigenvalues

- For a one-dimensional Hamiltonian of the form


$$H = \frac{p^2}{2m} + V(x)$$

the sequence of energy eigenvalues must satisfy

$$E_n \leq n^2$$



Eratosthenes sieve



It does not exist a close formula for the n-th prime number


However their scaling law is captured by this simple formula

$$p_n \simeq n \log n$$

Hence, there must exist a quantum Hamiltonian that has the primes as eigenvalues!


Counting the Primes

$\pi(x)$: gives the number of primes less or equal to x




Counting the Primes

$\pi(x)$: gives the number of primes less or equal to x



Counting the Primes

$\pi(x)$: gives the number of primes less or equal to x



Prime Number Theorem: Riemann

$$\pi(x) \simeq R(x)$$

$$R(x) = \sum_{n=1}^{\infty} \frac{\mu(n)}{n} Li(x^{1/n})$$


$$Li(t) \equiv \int_1^t \frac{ds}{\log s}$$

Prime Number Theorem: Riemann


$$\pi(x) \simeq R(x)$$

$$R(x) = \sum_{n=1}^{\infty} \frac{\mu(n)}{n} Li(x^{1/n})$$

$$\mu(n) = \begin{cases} 1 & \text{if } n \text{ is squarefree with an even number of prime factors} \\ -1 & \text{if } n \text{ is squarefree with an odd number of prime factors} \\ 0 & \text{if } n \text{ has a squared prime factor} \end{cases}$$



“The music of the primes”




Prime Number Theorem: Riemann

$$R(x) = \sum_{n=1}^{\infty} \frac{\mu(n)}{n} Li(x^{1/n})$$

$$\pi(x) = R(x) - \sum_{\rho} R(x^{\rho})$$

ρ : non-trivial zeros of the $\zeta(z)$ Riemann function in the critical strip

Climbing the staircase




DR. JEKYLL and MR. HYDE

- On a large scale, primes have extremely smooth distribution

$$n < p_k < 2n$$

$$p_{k+1} < 2p_k$$



DR. JEKYLL and MR. HYDE

- On a small scale, however, primes have highly unpredictable and irregular behavior

Example: Gap between the primes

1. Many (infinite?) twins of primes

(11,13) (17,19) (41,43) ... (347,349) ...

2. Arbitrarily large interval without a single prime!!

$$(10^{12} + 1)! + n \quad , \quad n = 2, 3, \dots, 10^{12} + 1$$

Inverse problems

Given an admissible sequence of numbers S_n ,
how to find the potential $V(x)$?





- Semi-classical method
- Dressing method (solitonic equations)

Quantum experiment



Primality test


$$E = N \hbar \omega$$




$$\int_{\varphi=0}^{\varphi=2\pi} P_{\varphi} d\varphi = j\hbar \int_{\varphi=0}^{\varphi=2\pi} P_r dr \quad \text{K}\hbar \quad 3p$$

$$E_{j,K} = - \frac{2\pi^2 m_e e^2 Z^2}{h^2} \frac{1}{(j+K)^2}$$

$$j+K=n$$

$$E_n = - \frac{2\pi^2 m_e e^2 Z^2}{h^2} \frac{1}{n^2} \quad h\nu_{m,n} = E_n - E_m$$


A diagram of an atom with a central nucleus and several elliptical orbits. The orbits are labeled with quantum numbers: 3s, 3p, and 3d. The 3s orbit is the innermost, followed by 3p, and then 3d. The orbits are drawn in a way that shows their orientation and relative size.



Semi-classical potential

$$\oint p(x) dx = \oint \sqrt{E - V(x)} dx = n \hbar$$

This formula can be inverted, i.e. once the energy levels E_n are assigned, we can find the potential $V(x)$!

Semi-classical potential

GM, (1995)

$$x(V) = \frac{\hbar}{\sqrt{2m}} \int_{E_0}^V \frac{dE}{\omega(E) \sqrt{V - E}} ,$$

$$E_n \rightarrow p_n \qquad p_n = \pi^{-1}(n)$$

$$\omega(p) = \frac{dp_n}{dn} = \frac{1}{\frac{dn}{dp_n}}$$


Semi-classical potential

GM, (1995)


$$x(V) = \frac{\hbar}{\sqrt{2m}} \int_{E_0}^V \frac{dE}{\omega(E) \sqrt{V - E}} ,$$


For prime numbers...

$$\omega^{-1}(E) = \sum_{n=1}^{\infty} \frac{\mu(n)}{n} \frac{\mathcal{E}^{\frac{1-n}{n}}}{\ln \mathcal{E}}$$




$$V(x) \simeq x^2 \log x$$






Holographic realization of the prime number quantum potential

Donatella Cassettari^{a,*}, Giuseppe Mussardo^b and Andrea Trombettoni^{b,c}






SUSY Quantum Mechanics



$$V_1(x) \rightarrow V_2(x) \rightarrow V_3(x) \rightarrow \dots$$


$$E = \{E_1, E_2, E_3, \dots, E_n\}$$

Quantum potential relative to the first 45 primes




Quantum potential relative to the first 150 primes
approximated by the first 35 Chebychev polys


Potential



$n=35$



Parametric resonance




$$H \rightarrow H + W(x) \cos \omega t$$

$$W_I = e^{iHt/\hbar} W e^{-iHt/\hbar}$$


$$|\alpha; t\rangle_i = \sum_n c_n(t) |n\rangle$$

Parametric resonance



$$\dot{c}_k = -i (M(t))_{km} c_m$$

$$M(t) = \begin{pmatrix} W_{11} & W_{12} e^{i\omega_{12}t} & \cdots \\ W_{21} e^{i\omega_{21}t} & W_{22} & \cdots \\ \vdots & \vdots & \ddots \end{pmatrix}$$



New perspectives on a series of problems...

- Lucky numbers
- Goldbach conjecture
- Factorization
-

Integer Factorization

(GM, Trombettoni)

Integer Factorization by Quantum Measurements

G. Mussardo^{1,2} and A. Trombettoni^{3,2}

¹SISSA, Via Bonomea 265, I-34136 Trieste, Italy.

²INFN, Sezione di Trieste, Via Valerio 2, I-34127 Trieste, Italy

³Dipartimento di Fisica, Università di Trieste, Strada Costiera 11, I-34151 Trieste, Italy

Quantum algorithms are at the heart of the ongoing efforts to use quantum mechanics to solve computational problems unsolvable on ordinary classical computers^{1–3}. Their common feature is the use of genuine quantum properties such as entanglement and superposition of states⁴. Among the known quantum algorithms, a special role is played by the Shor algorithm^{5,6}, i.e. a polynomial-time quantum algorithm for integer factorization, with far reaching potential applications in several fields, such as cryptography⁷. For an integer N of the order of 2^n , i.e. with n digits, the Shor algorithm permits its factorization in (order of) n steps. This results in an exponential gain in computational efficiency with respect to the best known classical algorithms. Here we present a different algorithm for integer factorization based on another genuine quantum property: quantum measurement^{8–10}. In this new scheme, the factorization of the integer N is achieved in a number of steps equal to the number of its prime factors, referred to as k —e.g., if N is the product of two primes, two quantum measurements are enough, regardless of the number of digits n of the number N . Since k is the lower bound to the number of operations one can do to factorize a general integer, then one sees that a quantum mechanical setup can saturate such a bound. Once established this, we discuss how the algorithm can physically be ran. We argue that one needs a single-purpose device where quantum measurements of an observable with assigned spectrum can be performed. The preparation from scratch of this device requires the solution, once for all and not for each factorization operation, of $\sim 2^n$ differential equations, a task that with a quantum computer can be accomplished in n steps.

Introduction. Recent progress in the implementation of quantum devices has led to the experimental demonstration of some instances of quantum advantage. This happens when a specific computational problem may be solved faster and more efficiently on quantum processors rather than a classical computer¹³. To achieve this goal the quantum processor must have an architecture made at least of several tens of qubits and long enough decoherence times.

A notable example, from a historical and conceptual point of view, of a clear quantum advantage is provided by the Shor algorithm^{5,6}. This algorithm indicates how to solve efficiently on a quantum computer the long-standing problem of finding the prime factors of an integer number N . Assuming that such a number N is of order 2^n , the Shor algorithm exploits in an ingenious way the implementation of the discrete Fourier transform on n qubits. To date, its validity has been shown with the factorization of a small numbers (the present computational bottleneck being the quantum modular exponentiation). The factorization of the number, $15 = 3 \times 5$, was done using 7 qubits with an NMR implementation of a quantum computer¹⁴. Similar demonstrations were performed using photonic^{15,16} and solid-state qubits¹⁷, while in 2012, with the n qubits control register replaced by a single qubit recycled n times, it was achieved the factorization of the integer $21 = 3 \times 7$ ¹⁸. Despite their simplicity, these examples nevertheless provide a *proof of principle* realization of the algorithm.

In this paper we present a different route for integer factorization, based on an algorithm which exploits another genuine quantum property: projective quantum measurement^{8–10}. As it is well known from quantum mechanics axioms, if a physical system is in a normalised state $|\psi\rangle$, a measurement of an observable $\hat{O}$ will yield one eigenvalue α of its spectrum with probability $|\langle\psi|\alpha\rangle|^2$, where $|\alpha\rangle$ is the normalised eigenfunction corresponding to the eigenvalue α

$$\hat{O}|\alpha\rangle = \alpha|\alpha\rangle. \quad (1)$$

As a result of the measurement, the system state will change from $|\psi\rangle$ to $|\alpha\rangle$. For problems related to number theory, interesting spectra to consider are: (a) the natural numbers, corresponding to the Hamiltonian of an harmonic oscillator^{9,10}; (b) the primes^{11,12}; and (c) the logarithm of the primes^{40–45}. Employing such spectra, one may translate number theory problems in quantum physical settings. As an example of this general philosophy, in this paper we show that with a suitable choice of the operator $\hat{O}$ is possible to determine the prime factors of an integer number N by making a finite set of quantum measurements.

Integer Factorization

(GM, Trombettoni)

Given an integer N , find its unique prime factorization

$$N = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_k^{\alpha_k}$$

Integer Factorization

(GM, Trombettoni)



(Quantum) Degeneracy

$$\begin{aligned}\log N &= \log p_1 + \log \hat{N}_1 \\ &= \log p_2 + \log \hat{N}_2 \\ &\quad \vdots \\ &= \log p_k + \log \hat{N}_k\end{aligned}\quad \left. \vphantom{\begin{aligned}\log N \\ = \log p_2 + \log \hat{N}_2 \\ \vdots \\ = \log p_k + \log \hat{N}_k\end{aligned}} \right\} \begin{array}{l} \text{k} \\ \text{times} \end{array}$$

$$\log \hat{N}_j = \log \left(p_1^{\alpha_1} \cdots p_j^{\alpha_j - 1} \cdots p_k^{\alpha_k} \right)$$

Quantum Hamiltonians

$$H = H_1 + H_2$$

$$H_1 = \frac{1}{2}p_x^2 + V(x) \qquad E_n^{(1)} = \{\log p_k\}$$

$$H_2 = \frac{1}{2}p_y^2 + W(x) \qquad E_n^{(2)} = \{\log N\}$$

$$E_n = E_n^{(1)} + E_n^{(2)}$$

Quantum Hamiltonians

$$H = H_1 + H_2$$

$$E_n = E_n^{(1)} + E_n^{(2)}$$



$d(n) = \#$ of distinct prime factors
of the integer number n


Initial state

$$|\Psi\rangle = |\log N\rangle = \sum_{n=1}^k c_n |\log p_n\rangle |\log \hat{N}_n\rangle$$


where the coefficients may also be regarded as random

It is however crucial that all of them are different from 0


Branching tree paths



Branching tree paths



Branching tree paths



Conclusions

- Interesting interplays between Number Theory & Physics
- Proof of principle: it is possible to realize in lab quantum mechanics potentials which encode interesting arithmetic sequences
- These realizations transform genuine mathematical questions into quantum experiments
- These quantum abacuses provide an alternative way to implement quantum algorithms

