

Shakhov collision model for relativistic fluids

Victor E. Ambruş

Department of Physics, West University of Timișoara, Romania

Work in collaboration with E. Molnár and D. Wagner

GGI: Hydro,
Florence, 16th May 2025



Outline

Introduction

Anderson-Witting (RTA) model

First-order relativistic Shakhov model

Application: Bjorken flow

Application: Sound waves

Second-order relativistic Shakhov model

Application: Shear-bulk coupling

Application: Shear-diffusion coupling

Application: Ultrarelativistic hard spheres (Riemann problem)

Code availability


Conclusions

Section 1


Introduction

Relativistic hydro playground: Heavy-ion collisions

[See talk by C. Werthmann (Tue, 11:15)]




- ▶ Shortly after the collision, the system is far-from-equilibrium.
- ▶ Pre-eq. dynamics require a non-eq. description.
- ▶ Strongly-interacting QGP leaves imprints of thermalization and collectivity in final-state observables.



[Venaruzzo, PhD Thesis, 2011]

Hydro vs Kinetic theory



[Ambruş, Bazzanini, Gabbana, Simeoni, Succi, Nature Comput. Sci. 2 (2022) 641]

- ▶ Hydro employed in HIC modelling, but it breaks down far from eq.
- ▶ Kinetic theory overcomes this limitation, but realistic simulations are expensive due to $C[f]$.

AMPT: [He, Edmonds, Lin, Liu, Molnar, Wang, PLB 753 (2016) 506]
 BAMPS: Greif, Greiner, Schenke, Schlichting, Xu [PRD 96 (2017) 091504]

- ▶ RTA: $C_{AW}[f] = -\frac{E_{\mathbf{k}}}{\tau_R} (f_{\mathbf{k}} - f_{0\mathbf{k}}) \Rightarrow 1 - 2 \text{ o.m. faster than BAMPS.}$

VEA, Busuioc, Fotakis, Gallmeister, Greiner [PRD 104 (2021) 094022]

- ▶ τ_R fixes the IR limit of RTA by matching e.g. η to that of $C[f] \Rightarrow$ good agreement with BAMPS.

Section 2

Anderson-Witting (RTA) model

Anderson-Witting model

[Anderson, Witting, Physica **74** (1974) 466]

- The Anderson & Witting RTA reads $[E_{\mathbf{k}} = k^\mu u_\mu]$

$$k^\mu \partial_\mu f_{\mathbf{k}} = C_{\text{AW}}[f], \quad C_{\text{AW}}[f] = -\frac{E_{\mathbf{k}}}{\tau_R} (f_{\mathbf{k}} - f_{0\mathbf{k}}). \quad (1)$$

- N^μ and $T^{\mu\nu}$ are obtained from $f_{\mathbf{k}}$: $[dK = g d^3k / [k_0 (2\pi)^3]]$

$$N^\mu = \int dK k^\mu f_{\mathbf{k}}, \quad T^{\mu\nu} = \int dK k^\mu k^\nu f_{\mathbf{k}}. \quad (2)$$

- $f_{0\mathbf{k}}$ describes LTE, for which $[\Delta^{\mu\nu} = g^{\mu\nu} - u^\mu u^\nu]$

$$N_0^\mu = n_0 u^\mu, \quad T_0^{\mu\nu} = \epsilon_0 u^\mu u^\nu - P_0 \Delta^{\mu\nu}, \quad (3)$$

- Imposing $\partial_\mu N^\mu = \partial_\nu T^{\mu\nu} = 0$ requires Landau matching:

$$n = n_0, \quad \epsilon = \epsilon_0, \quad T^\mu{}_\nu u^\nu = \epsilon u^\mu. \quad (4)$$

- $C_{\text{AW}}[f]$ drives $f_{\mathbf{k}}$ towards $f_{0\mathbf{k}}$ on the timescale τ_R .

Chapman-Enskog expansion

- ▶ Out of eq., N^μ and $T^{\mu\nu}$ receive dissipative corrections:

$$N^\mu - N_0^\mu = V^\mu, \quad T^{\mu\nu} - T_0^{\mu\nu} = -\Pi\Delta^{\mu\nu} + \pi^{\mu\nu}. \quad (5)$$

- ▶ The dissipative quantits. can be obtained as moments of $\delta f_{\mathbf{k}}$:

$$\Pi = -\frac{m^2}{3} \int dK \delta f_{\mathbf{k}}, \quad V^\mu = \int dK k^{\langle\mu\rangle} \delta f_{\mathbf{k}}, \quad \pi^{\mu\nu} = \int dK k^{\langle\mu} k^{\nu\rangle} \delta f_{\mathbf{k}}, \quad (6)$$

with $k^{\langle\mu\rangle} = \Delta_{\alpha}^{\mu} k^{\alpha}$ and $k^{\langle\mu} k^{\nu\rangle} = \Delta_{\alpha\beta}^{\mu\nu} k^{\alpha} k^{\beta}$ *irreducible* tensors.

- ▶ Employing the Chapman-Enskog procedure gives

$$\delta f_{\mathbf{k}} \equiv f_{\mathbf{k}} - f_{0\mathbf{k}} \simeq -\frac{\tau_R}{E_{\mathbf{k}}} k^{\mu} \partial_{\mu} f_{0\mathbf{k}}. \quad (7)$$

- ▶ Taking moments as in Eq. (6) gives

$$\Pi = -\zeta_R \theta, \quad V^\mu = \kappa_R \nabla^{\mu} \alpha, \quad \pi^{\mu\nu} = 2\eta_R \sigma^{\mu\nu}, \quad (8)$$

where ζ_R , κ_R and η_R are given by

$$\zeta_R = \frac{m^2}{3} \tau_R \alpha_0^{(0)}, \quad \kappa_R = \tau_R \alpha_0^{(1)}, \quad \eta_R = \tau_R \alpha_0^{(2)}. \quad (9)$$

where $\alpha_0^{(\ell)}$ are τ_R -independent thermodynamic functions.

QGP Transport coefficients

- Bayesian estimation shows that η/s and ζ/s can be parametrized as

J. E. Bernhard, J. S. Moreland, S. A. Bass, *Nature Phys.* **15** (2019) 1113

$$\frac{\eta}{s} = (\eta/s)_{\min} + (\eta/s)_{\text{slope}}(T - T_c) \left(\frac{T}{T_c} \right)^{(\eta/s)_{\text{crv}}}, \quad (10)$$


$$\frac{\zeta}{s} = (\zeta/s)_{\max} \times \left[1 + \left(\frac{T - T_{\text{peak}}}{(\zeta/s)_{\text{width}}} \right)^2 \right]^{-1}. \quad (11)$$

- RTA allows, e.g. η to be specified by setting

$$\tau_R = \frac{\eta}{\alpha_0^{(2)}},$$

however, ζ/η is fixed uniquely by

$$\frac{\zeta}{\eta} = \frac{m^2 \alpha_0^{(0)}}{3\alpha_0^{(2)}},$$



which does not resemble the (ζ/η) in the QGP.

Section 3

First-order relativistic Shakhov model

- ▶ We consider a Shakhov-like extension:

[Shakhov, Fluid Dyn. 3 (1968) 112]

$$C_S[f] = -\frac{E_{\mathbf{k}}}{\tau_R}(f_{\mathbf{k}} - f_{S\mathbf{k}}), \quad (12)$$

where $f_{S\mathbf{k}} \rightarrow f_{0\mathbf{k}}$ as $\delta f_{\mathbf{k}} = f_{\mathbf{k}} - f_{0\mathbf{k}} \rightarrow 0$.

- ▶ Shakhov vs. AW: $f_{\mathbf{k}}$ relaxes towards $f_{0\mathbf{k}}$ on a modified path.
- ▶ $\partial_\mu N^\mu = \partial_\nu T^{\mu\nu} = 0$ imply:

$$u_\mu N^\mu = u_\mu N_S^\mu, \quad u_\nu T^{\mu\nu} = u_\nu T_S^{\mu\nu}, \quad (13)$$

which allows for plenty of degrees of freedom (δn , $\delta\epsilon$, W^μ , etc).

- ▶ For simplicity, we stick to the Landau matching conditions:

$$\delta n = \delta\epsilon = 0, \quad T^{\mu\nu} u_\nu = \epsilon u^\mu. \quad (14)$$

Shakohv-like extension

- ▶ Employing the Chapman-Enskog procedure gives

$$\delta f_{\mathbf{k}} - \delta f_{S\mathbf{k}} = -\frac{\tau_R}{E_{\mathbf{k}}} k^\mu \partial_\mu f_{0\mathbf{k}}, \quad (15)$$

leading to

$$\Pi - \Pi_S = -\zeta_R \theta, \quad V^\mu - V_S^\mu = \kappa_R \nabla^\mu \alpha, \quad \pi^{\mu\nu} - \pi_S^{\mu\nu} = 2\eta_R \sigma^{\mu\nu}. \quad (16)$$

- ▶ We seek to replace ζ_R etc by independent transport coefficients:

$$\begin{aligned} \Pi &\simeq -\zeta_S \theta, & V^\mu &\simeq \kappa_S \nabla^\mu \alpha, & \pi^{\mu\nu} &\simeq 2\eta_S \sigma^{\mu\nu}, \\ \zeta_S &= \frac{\tau_\Pi}{\tau_R} \zeta_R, & \kappa_S &= \frac{\tau_V}{\tau_R} \kappa_R, & \eta_S &= \frac{\tau_\pi}{\tau_R} \eta_R. \end{aligned} \quad (17)$$

- ▶ Eq. (17) can be obtained from Eq. (16) when

$$\begin{aligned} \Pi_S &= \Pi \left(1 - \frac{\tau_\Pi}{\tau_R} \right), & V_S^\mu &= V^\mu \left(1 - \frac{\tau_V}{\tau_R} \right), \\ \pi_S^{\mu\nu} &= \pi^{\mu\nu} \left(1 - \frac{\tau_\pi}{\tau_R} \right). \end{aligned} \quad (18)$$

Minimal $\delta f_{S\mathbf{k}}$

- The solution can be written as $\delta f_{S\mathbf{k}} = f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} \mathbb{S}_{\mathbf{k}}$, where

$$\begin{aligned} \mathbb{S}_{\mathbf{k}} = & -\frac{3\Pi}{m^2} \left(1 - \frac{\tau_R}{\tau_\Pi}\right) \mathcal{H}_{\mathbf{k}0}^{(0)} + k_{\langle\mu} V^{\mu} \left(1 - \frac{\tau_R}{\tau_V}\right) \mathcal{H}_{\mathbf{k}0}^{(1)} \\ & + k_{\langle\mu} k_{\nu\rangle} \pi^{\mu\nu} \left(1 - \frac{\tau_R}{\tau_\pi}\right) \mathcal{H}_{\mathbf{k}0}^{(2)}. \end{aligned} \quad (19)$$

- $\mathcal{H}_{\mathbf{k}0}^{(\ell)}$ are polynomials in $E_{\mathbf{k}}$ satisfying the constraints: [\[DNMR, PRD 85 \(2012\) 114047\]](#)

$$\begin{aligned} \text{Bulk visc. p.} & \Rightarrow \begin{pmatrix} \rho_{S;0} \\ \rho_{S;1} \\ \rho_{S;2} \end{pmatrix} = \int dK \begin{pmatrix} 1 \\ E_{\mathbf{k}} \\ E_{\mathbf{k}}^2 \end{pmatrix} \delta f_{S\mathbf{k}} = \begin{pmatrix} -3\Pi_S/m^2 \\ 0 \\ 0 \end{pmatrix}, \\ \text{Particle cons.} & \\ \text{Energy cons.} & \\ \text{Diff. current} & \Rightarrow \begin{pmatrix} \rho_{S;0}^\mu \\ \rho_{S;1}^\mu \end{pmatrix} = \int dK \begin{pmatrix} 1 \\ E_{\mathbf{k}} \end{pmatrix} k^{\langle\mu} \delta f_{S\mathbf{k}} = \begin{pmatrix} V_S^\mu \\ 0 \end{pmatrix}, \\ \text{Mom. cons.} & \\ \text{SS tens.} & \Rightarrow \rho_{S;0}^{\mu\nu} = \int dK k^{\langle\mu} k^{\nu\rangle} \delta f_{S\mathbf{k}} = \pi_S^{\mu\nu}. \end{aligned} \quad (20)$$

- The solution can be written as $\delta f_{S\mathbf{k}} = f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} S_{\mathbf{k}}$, where

$$S_{\mathbf{k}} = -\frac{3\Pi}{m^2} \left(1 - \frac{\tau_R}{\tau_\Pi}\right) \mathcal{H}_{\mathbf{k}0}^{(0)} + k_{\langle\mu} V^\mu \left(1 - \frac{\tau_R}{\tau_V}\right) \mathcal{H}_{\mathbf{k}0}^{(1)} + k_{\langle\mu} k_{\nu\rangle} \pi^{\mu\nu} \left(1 - \frac{\tau_R}{\tau_\pi}\right) \mathcal{H}_{\mathbf{k}0}^{(2)}. \quad (21)$$

- $\mathcal{H}_{\mathbf{k}0}^{(0)}$ satisfies 3 constraints $(\rho_{S;0,1,2}) \Rightarrow$ polynomial of order 2.
- $\mathcal{H}_{\mathbf{k}0}^{(1)}$ satisfies 2 constraints $(\rho_{S;0,1}) \Rightarrow$ polynomial of order 1.
- $\mathcal{H}_{\mathbf{k}0}^{(2)}$ satisfies 1 constraint $(\rho_{S;0,1}) \Rightarrow$ polynomial of order 0.
- The simplest solution is:

$$\mathcal{H}_{\mathbf{k}0}^{(0)} = \frac{G_{33} - G_{23}E_{\mathbf{k}} + G_{22}E_{\mathbf{k}}^2}{J_{00}G_{33} - J_{10}G_{23} + J_{20}G_{22}},$$

$$\mathcal{H}_{\mathbf{k}0}^{(1)} = \frac{J_{31}E_{\mathbf{k}} - J_{41}}{J_{21}J_{41} - J_{31}^2}, \quad \mathcal{H}_{\mathbf{k}0}^{(2)} = \frac{1}{2J_{42}}, \quad (22)$$

where $G_{nm} = J_{n0}J_{m0} - J_{n-1,0}J_{m+1,0}$ and

$$J_{nq} = \frac{1}{(2q+1)!!} \int dK E_{\mathbf{k}}^{n-2q} (-\Delta^{\alpha\beta} k_\alpha k_\beta)^q f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}}. \quad (23)$$

Entropy production

[VEA, Molnár, PLB 855 (2024) 138795]

- ▶ The entropy current is given by

[classical stat. used for simplicity]

$$S^\mu = - \int dK k^\mu (f_{\mathbf{k}} \ln f_{\mathbf{k}} - f_{\mathbf{k}}). \quad (24)$$

- ▶ In the Shakhov model, $k^\mu \partial_\mu f = C_S[f]$ and

$$\partial_\mu S^\mu = - \int dK C_S[f] \ln f_{\mathbf{k}} = \frac{1}{\tau_R} \int dK E_{\mathbf{k}} (\delta f_{\mathbf{k}} - \delta f_{S\mathbf{k}}) \ln f_{\mathbf{k}}. \quad (25)$$

- ▶ When $\phi_{\mathbf{k}} = \delta f_{\mathbf{k}}/f_{0\mathbf{k}}$ is small, detailed manipulations lead to

$$\partial_\mu S^\mu \simeq \frac{\beta}{\zeta_S} \Pi^2 - \frac{1}{\kappa_S} V_\mu V^\mu + \frac{\beta}{2\eta_S} \pi_{\mu\nu} \pi^{\mu\nu} \geq 0. \quad (26)$$

- ▶ Close to eq., the S-model satisfies the 2nd law of thermodynamics.
- ▶ Proof far from eq. unavailable even for non-rel. Shakhov!

Section 4

Application: Bjorken flow

Application: Bjorken flow

- ▶ Bjorken model: flow invariant under longitudinal boosts:

$$u^\mu \partial_\mu = \frac{t}{\tau} \partial_t + \frac{z}{\tau} \partial_z, \quad \tau = \sqrt{t^2 - z^2}, \quad \eta_s = \tanh^{-1}(z/t). \quad (27)$$

- ▶ In Bjorken coordinates $(\tau, \mathbf{x}_\perp, \eta_s)$,

$$T^{\mu\nu} = \text{diag}(e, P_T, P_T, \tau^{-2} P_L),$$
$$P_T = P + \Pi - \frac{\pi_d}{2}, \quad P_L = P + \Pi + \pi_d. \quad (28)$$



- ▶ In 2nd-order hydro, we have: [\[Denicol, Florkowski, Ryblewski, Strickland, PRC 90 \(2014\) 044905\]](#)

$$\tau \dot{\epsilon} + \epsilon + P_L = 0, \quad (29a)$$

$$\tau \dot{\Pi} + \left(\frac{\delta_{\Pi\Pi}}{\tau_\Pi} + \frac{\tau}{\tau_\Pi} \right) \Pi + \frac{\lambda_{\Pi\pi}}{\tau_\Pi} \pi_d = -\frac{\zeta}{\tau_\Pi},$$
$$\tau \dot{\pi}_d + \left(\frac{\delta_{\pi\pi}}{\tau_\pi} + \frac{\tau_{\pi\pi}}{3\tau_\pi} + \frac{\tau}{\tau_\pi} \right) \pi_d + \frac{2\lambda_{\pi\Pi}}{3\tau_\pi} \Pi = -\frac{4\eta}{3\tau_\pi}. \quad (29b)$$

- ▶ We employ the Shakhov model to control ζ independently from η .

Shakhov model: ζ vs. η



► Choosing $\tau_R = \tau_{\text{II}}$, the Shakhov distribution becomes

$$f_{\text{Sk}} = f_{0\mathbf{k}} \left[1 + \frac{\beta^2 k_{\mu} k_{\nu} \pi^{\mu\nu}}{2(e + P)} \left(1 - \frac{\tau_{\text{II}}}{\tau_{\pi}} \right) \right]. \quad (30)$$

► Left panel: τ_{π} is fixed and τ_{II} is varied using the Shakhov model.

► Right panel: τ_{II} is fixed and τ_{π} is varied using the Shakhov model.

► $m = 1$ GeV; $\tau_0 = 0.5$ fm; $\beta_0^{-1} = 0.6$ GeV; For $\tau_{\pi} = 0.5$ fm, $4\pi\eta/s \simeq 3.3$ at $\tau = \tau_0$.

Section 5

Application: Sound waves

Application: Sound waves

- ▶ We now consider an infinitesimal perturbation propagating in an ultrarelativistic fluid at rest.
- ▶ Writing $u^\mu \simeq (1, 0, 0, \delta v)$, $\epsilon = \epsilon_0 + \delta\epsilon$ and $n = n_0 + \delta n$, we have

$$\begin{aligned}\partial_t \delta n + n_0 \partial_z \delta v + \partial_z \delta V &= 0, \\ \partial_t \delta \epsilon + (\epsilon_0 + P_0) \partial_z \delta v &= 0, \\ (\epsilon_0 + P_0) \partial_t \delta v + \partial_z \delta P + \partial_z \delta \pi &= 0, \\ \tau_V \partial_t \delta V + \delta V + \kappa \partial_z \delta \alpha - \ell_{V\pi} \partial_z \delta \pi &= 0, \\ \tau_\pi \partial_t \delta \pi + \delta \pi + \frac{4\eta}{3} \partial_z \delta v + \frac{2}{3} \ell_{\pi V} \partial_z \delta V &= 0,\end{aligned}\tag{31}$$

where $\delta V = V^z$ and $\delta \pi = \pi^{zz}/\gamma^2$.

- ▶ **In RTA**, $\ell_{V\pi} = \ell_{\pi V} = 0$.



[VEA, Molnár, Rischke, PRD **106** (2022) 076005]

- ▶ We track the time evolution of the amplitudes

$$\widetilde{\delta V} = \frac{2}{L} \int_0^L dz \delta V \cos(kz), \quad \widetilde{\delta \pi} = \frac{2}{L} \int_0^L dz \delta \pi \sin(kz).\tag{32}$$

- ▶ We employ the Shakhov model to control κ independently from η .

Shakhov model: κ vs. η



► Setting $\tau_R = \tau_\pi$, the Shakhov distribution becomes

$$f_{S\mathbf{k}} = f_{0\mathbf{k}} \left[1 + \frac{k_\mu V^\mu}{P} (\beta E_{\mathbf{k}} - 5) \left(1 - \frac{\tau_\pi}{\tau_V} \right) \right]. \quad (33)$$

Section 6

Second-order relativistic Shakhov model

Beyond first order: second-order transport coefficients?


- ▶ Navier-Stokes hydrodynamics is acausal \Rightarrow a-relativistic!
- ▶ One example of causal hydro is MIS 2nd order hydro, by which e.g. $\pi^{\mu\nu}$ evolves according to $\tau_\pi \dot{\pi}^{\langle\mu\nu\rangle} + \pi^{\mu\nu} = 2\eta\sigma^{\mu\nu} + \mathcal{J}^{\mu\nu} + \mathcal{R}^{\mu\nu}$, with

$$\begin{aligned}\mathcal{J}^{\mu\nu} &= 2\tau_\pi \pi_\lambda^{\langle\mu} \omega^{\nu\rangle\lambda} - \delta_{\pi\pi} \pi^{\mu\nu} \theta - \tau_{\pi\pi} \pi^{\lambda\langle\mu} \sigma_\lambda^{\nu\rangle} + \lambda_{\pi\Pi} \Pi \sigma^{\mu\nu} \\ &\quad - \tau_{\pi V} V^{\langle\mu} \dot{u}^{\nu\rangle} + \ell_{\pi V} \nabla^{\langle\mu} V^{\nu\rangle} + \lambda_{\pi V} V^{\langle\mu} \nabla^{\nu\rangle} \alpha, \\ \mathcal{R}^{\mu\nu} &= \varphi_6 \Pi \pi^{\mu\nu} + \varphi_7 \pi^{\lambda\langle\mu} \pi_\lambda^{\nu\rangle} + \varphi_8 V^{\langle\mu} V^{\nu\rangle}.\end{aligned}\tag{34}$$

- ▶ In RTA, $\mathcal{R}^{\mu\nu} = 0$.
- ▶ 2nd-order t.c. are important e.g. in preeq!
- ▶ In conformal RTA, $\delta_{\pi\pi} + \tau_{\pi\pi}/3 = 38/21$.
- ▶ Solving hydro with $\delta_{\pi\pi} + \tau_{\pi\pi}/3 = 31/15$ gives much better agreement with RTA!

[J.-P. Blaizot, L. Yan, PRC **104** (2021) 055201]

- ▶ Etc...



Second-order hydro from KT

► In the method of moments, second-order hydro can be derived using:

- Irreducible moments of $\delta f_{\mathbf{k}}$: $\rho_r^{\mu_1 \dots \mu_\ell} = \int dK E_{\mathbf{k}}^r k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} \delta f_{\mathbf{k}}$.
- Irreducible moments of $C[f]$: $C_r^{\mu_1 \dots \mu_\ell} = \int dK E_{\mathbf{k}}^r k^{\langle \mu_1} \dots k^{\mu_\ell \rangle} C[f]$.
- Define collision matrix via $C_{r-1}^{\mu_1 \dots \mu_\ell} = - \sum_n \mathcal{A}_{rn}^{(\ell)} \rho_n^{\mu_1 \dots \mu_\ell}$.
- Define inverse matrix $\tau_{rn}^{(\ell)}$ via $\sum_n \tau_{rn}^{(\ell)} \mathcal{A}_{nm}^{(\ell)} = \delta_{rm}$.

► The 1st-order transport coeffs. are

$$\zeta_r = \frac{m^2}{3} \sum_n \tau_{r\mathbf{n}}^{(0)} \alpha_n^{(0)}, \quad \kappa_r = \sum_n \tau_{r\mathbf{n}}^{(1)} \alpha_n^{(1)}, \quad \eta_r = \sum_n \tau_{r\mathbf{n}}^{(2)} \alpha_n^{(2)}.$$

► The relaxation times can be obtained via [\[Wagner, Palermo, VEA, PRD 106 \(2022\) 016013\]](#)

$$\tau_{\Pi} = \sum_r \tau_{0\mathbf{r}}^{(0)} C_r^{(0)}, \quad \tau_V = \sum_r \tau_{0\mathbf{r}}^{(1)} C_r^{(1)}, \quad \tau_{\pi} = \sum_r \tau_{0\mathbf{r}}^{(2)} C_r^{(2)}, \quad (35)$$

with $C_r^{(0)} = \zeta_r / \zeta_0$, $C_r^{(1)} = \kappa_r / \kappa_0$ and $C_r^{(2)} = \eta_r / \eta_0$.

- ...all other 2nd-order t.c. are computed using $\tau_{0\mathbf{n}}^{(\ell)}$ and $C_n^{(\ell)}$.
- **NOTE!** $C_0 = C_1 = C_1^\mu = 0$ to conserve mass & energy-momentum!
- Idea: Use Shakhov model to “manipulate” $\mathcal{A}_{rn}^{(\ell)}$.

From RTA to Shakhov

- In RTA, $C[f] = -\frac{E_{\mathbf{k}}}{\tau_R} \delta f_{\mathbf{k}}$ and

[VEA, Molnár, Rischke, PRD **106** (2022) 076005]

$$C_{r-1}^{\mu_1 \cdots \mu_\ell} = -\frac{1}{\tau_R} \rho_r^{\mu_1 \cdots \mu_\ell} \Rightarrow \mathcal{A}_{rn}^{(\ell)} = \frac{\delta_{rn}}{\tau_R} \Rightarrow \tau_{rn}^{(\ell)} = \tau_R \delta_{rn}. \quad (36)$$

- In the Shakhov model, $C_S = -\frac{E_{\mathbf{k}}}{\tau_R} [\delta f_{\mathbf{k}} - \delta f_{S\mathbf{k}}]$ and

$$C_{r-1}^{\mu_1 \cdots \mu_\ell} = -\frac{1}{\tau_R} [\rho_r^{\mu_1 \cdots \mu_\ell} - \rho_{S;r}^{\mu_1 \cdots \mu_\ell}], \quad (37)$$

where $\rho_{S;r}^{\mu_1 \cdots \mu_\ell}$ are essentially arbitrary.

- Imposing $C_{r-1}^{\mu_1 \cdots \mu_\ell} = -\sum_n \mathcal{A}_{rn}^{(\ell)} \rho_n^{\mu_1 \cdots \mu_\ell}$ suggests taking

$$\rho_{S;r}^{\mu_1 \cdots \mu_\ell} = \sum_n [\delta_{rn} - \tau_R \mathcal{A}_{rn}^{(\ell)}] \rho_n^{\mu_1 \cdots \mu_\ell}, \quad (38)$$

where $\mathcal{A}_{rn}^{(\ell)}$ is the desired collision matrix.

- Our approach is to fix a subset of $\rho_{\mathbb{S};r}^{\mu_1 \cdots \mu_\ell}$ with:

$$0 \leq \ell \leq L = 2, \quad -s_\ell \leq r \leq N_\ell, \quad (39)$$

where $s_\ell \equiv$ “shift” and $N_\ell \geq \{2, 1, 0\}$. [VEA, Molnár, Rischke, PRD **106** (2022) 076005]

- We construct $\delta f_{\mathbb{S}\mathbf{k}} \equiv f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} \mathbb{S}_{\mathbf{k}}$ as

$$\mathbb{S}_{\mathbf{k}} = \sum_{\ell=0}^L \sum_{n=-s_\ell}^{N_\ell} \rho_{\mathbb{S};n}^{\mu_1 \cdots \mu_\ell} E_{\mathbf{k}}^{-s_\ell} k_{\langle \mu_1} \cdots k_{\mu_\ell \rangle} \tilde{\mathcal{H}}_{\mathbf{k},n+s_\ell}^{(\ell)}, \quad (40)$$

with $\tilde{\mathcal{H}}_{\mathbf{k}n}^{(\ell)}$ ensuring $\rho_{\mathbb{S};n}^{\mu_1 \cdots \mu_\ell} = \int dK E_{\mathbf{k}}^n k^{\langle \mu_1} \cdots k^{\mu_\ell \rangle} \delta f_{\mathbb{S}\mathbf{k}}$.

Shakhov collision matrix

- ▶ Eq. (40) sets $\rho_{S;n}^{\mu_1 \dots \mu_\ell}$ for $-s_\ell \leq n \leq N_\ell$.
- ▶ In general $\rho_{S;r}^{\mu_1 \dots \mu_\ell} \neq 0$ when $r < -s_\ell$ and $r > N_\ell$.
- ▶ $\Rightarrow \mathcal{A}_{S;rn}^{(\ell)}$ contains non-trivial entries when $r < -s_\ell$ and $r > N_\ell$:

$$\mathcal{A}_{rn}^{(\ell)} = \begin{pmatrix} \frac{1}{\tau_R} \delta_{rn} & \mathcal{A}_{<;rn}^{(\ell)} & 0 \\ 0 & \mathcal{A}_{S;rn}^{(\ell)} & 0 \\ 0 & \mathcal{A}_{>;rn}^{(\ell)} & \frac{1}{\tau_R} \delta_{rn} \end{pmatrix}, \quad (41)$$

where $\mathcal{A}_{</>;rn}^{(\ell)}$ correspond to $r < -s_\ell$ and $r > N_\ell$, respectively.

- ▶ These entries supplement the $\tau_R^{-1} \delta_{rn}$ structure of AW with

$$\mathcal{A}_{</>;rn}^{(\ell)} = -\frac{1}{\tau_R} \tilde{\mathcal{F}}_{-(r+s_\ell), n+s_\ell}^{(\ell)} + \sum_{j=-s_\ell}^{N_\ell} \tilde{\mathcal{F}}_{-(r+s_\ell), j+s_\ell}^{(\ell)} \mathcal{A}_{S;jn}^{(\ell)}, \quad (42)$$

with $\tilde{\mathcal{F}}_{rn}^{(\ell)} \equiv \frac{\ell!}{(2\ell+1)!!} \int dK f_{0\mathbf{k}} \tilde{f}_{0\mathbf{k}} E_{\mathbf{k}}^{-2s_\ell-r} (\Delta^{\alpha\beta} k_\alpha k_\beta)^\ell \tilde{\mathcal{H}}_{\mathbf{k}n}^{(\ell)}$.

Inverse collision matrix

► The inverse matrix $\tau_{rn}^{(\ell)}$ reads

$$\tau_{rn}^{(\ell)} = \begin{pmatrix} \tau_R \delta_{rn} & \tau_{<;rn}^{(\ell)} & 0 \\ 0 & \tau_{S;rn}^{(\ell)} & 0 \\ 0 & \tau_{>;rn}^{(\ell)} & \tau_R \delta_{rn} \end{pmatrix}, \quad (43)$$

with $\tau_{S;rn}^{(\ell)} = [\mathcal{A}_{S;rn}^{(\ell)}]^{-1}$ a finite $(N_\ell + s_\ell + 1)^2$ matrix and

$$\tau_{<, >;rn}^{(\ell)} = -\tau_R \tilde{\mathcal{F}}_{-(r+s_\ell), n+s_\ell}^{(\ell)} + \sum_{j=-s_\ell}^{N_\ell} \tilde{\mathcal{F}}_{-(r+s_\ell), j+s_\ell}^{(\ell)} \tau_{S;jn}^{(\ell)}. \quad (44)$$

► For example, the shear viscosities $\eta_r = \sum_n \tau_{rn}^{(2)} \alpha_n^{(2)}$ are

$$\eta_{-s_\ell \leq r \leq N_\ell} = \sum_{n=-s_2}^{N_2} \tau_{S;rn}^{(2)} \alpha_n^{(2)},$$

$$\eta_{r, </>} = \tau_R \alpha_r^{(2)} + \sum_{n=-s_2}^{N_2} \tilde{\mathcal{F}}_{-r-s_2, n+s_2}^{(2)} (\eta_n - \tau_R \alpha_n^{(2)}). \quad (45)$$

Tunable coefficients in the Shakhov model

- The transport coefficients depend on

$$\begin{aligned}\tau_{0,n \neq 1,2}^{(0)} : & N_0 + s_0 - 1 \text{ entries; } & \mathcal{C}_{n \neq 1,2}^{(0)} \equiv \frac{\zeta_n}{\zeta_0} : & N_0 + s_0 - 2 \text{ extra lines,} \\ \tau_{0,n \neq 1}^{(1)} : & N_1 + s_1 \text{ entries; } & \mathcal{C}_{n \neq 1}^{(1)} \equiv \frac{\kappa_n}{\kappa_0} : & N_1 + s_1 - 1 \text{ extra lines,} \\ \tau_{0n}^{(2)} : & N_2 + s_2 + 1 \text{ entries; } & \mathcal{C}_n^{(2)} \equiv \frac{\eta_n}{\eta_0} : & N_2 + s_2 \text{ extra lines,}\end{aligned}$$

so in total:

$$[2(N_0 + s_0 + N_1 + s_1 + N_2 + s_2) - 3] \text{ transport coefficients, } (46)$$

plus a hidden degree of freedom given by τ_R .

- For an UR gas, the scalar sector is not important, leaving in total

$$[2(N_1 + s_1 + N_2 + s_2)] \text{ transport coefficients, } (47)$$

plus τ_R .

Section 7

Application: Shear-bulk coupling

Example: shear-bulk coupling

[VEA, Wagner, PRD 110 (2024) 056002]

- ▶ To illustrate the capabilities of the Shakhov model in the case of finite m , we consider again the Bjorken flow problem.
- ▶ In MIS hydro, the diffusive quantities evolve according to

$$\begin{aligned}\tau_{\Pi} \frac{d\Pi}{d\tau} + \Pi &= -\frac{1}{\tau} (\zeta + \delta_{\Pi\Pi}\Pi + \lambda_{\Pi\pi}\pi_d) , \\ \tau_{\pi} \frac{d\pi_d}{d\tau} + \pi_d &= -\frac{1}{\tau} \left[\frac{4\eta}{3} + \left(\delta_{\pi\pi} + \frac{\tau_{\pi\pi}}{3} \right) \pi_d + \frac{2\lambda_{\pi\Pi}}{3} \Pi \right] .\end{aligned}\quad (48)$$

- ▶ Our aim is to separately tune ζ , η and $\lambda_{\Pi\pi}$, i.e.

$$\frac{\lambda_{\Pi\pi}}{\tau_{\Pi}} = A \frac{\lambda_{\Pi\pi}^R}{\tau_R}, \quad \eta = H\eta_R, \quad \zeta = \zeta_R, \quad (49)$$


where $\lambda_{\Pi\pi}^R = \frac{m^2}{3} \tau_R \left(\mathcal{R}_{-2}^{(2)} + \frac{J_{10}}{J_{30}} \right)$ is the RTA expression, while A and H are arbitrary functions.

- ▶ This can be achieved using the following collision matrix:

$$\mathcal{A}_S^{(2)} = \frac{1}{\tau_R H} \begin{pmatrix} 1 & (1-A) \left(\mathcal{R}_{-2}^{(2)} + \frac{J_{10}}{J_{30}} \right) \\ 0 & 1 \end{pmatrix}, \quad \mathcal{R}_{-2}^{(2)} = \frac{\alpha_{-2}^{(2)}}{\alpha_0^{(2)}}. \quad (50)$$

Example: shear-bulk coupling

[VEA, Wagner, PRD 110 (2024) 056002]



- For definiteness, we consider $AH = 1 \Rightarrow$ bulk response $\lambda_{\Pi\pi}\pi_d$ remains unchanged (see central panel).
- The Shakhov $f_{S\mathbf{k}} = f_{0\mathbf{k}}\tilde{f}_{0\mathbf{k}}\mathbb{S}_{\mathbf{k}}$ has $\tilde{f}_{0\mathbf{k}} = 1$ (classical gas) and

$$\mathbb{S}_{\mathbf{k}} = \left(\pi_{S;-2} \mathfrak{h}_{\mathbf{k}0}^{(2)} + \pi_{S;0} \mathfrak{h}_{\mathbf{k}2}^{(2)} \right) \left(\frac{k_\eta^2}{\tau^2 k_\tau^2} - \frac{k_\perp^2}{2k_\tau^2} \right),$$

$$\pi_{S;r} = \pi_r - \tau_R \mathcal{A}_{S;rn}^{(2)} \pi_n, \quad k^\tau = \frac{tk^t - zk^z}{\tau}, \quad k^\eta = \frac{tk^z - zk^t}{\tau^2},$$

$$\mathfrak{h}_{\mathbf{k}0}^{(2)} = \frac{J_{42} - J_{22}E_{\mathbf{k}}^2}{2(J_{02}J_{42} - J_{22}^2)}, \quad \mathfrak{h}_{\mathbf{k}2}^{(2)} = \frac{-J_{22} + J_{02}E_{\mathbf{k}}^2}{2(J_{02}J_{42} - J_{22}^2)}. \quad (51)$$

Section 8

Application: Shear-diffusion coupling

Example: shear-diffusion coupling

[VEA, Wagner, PRD **110** (2024) 056002]

- ▶ Consider a longitudinal wave propagating along z .
- ▶ The linearized hydro equations for $\delta\pi \equiv \pi^{zz}$ and $\delta V \equiv V^z$ read

$$\begin{aligned}\tau_V \partial_t \delta V + \delta V &= -\kappa \partial_z \delta\alpha + \ell_{V\pi} \partial_z \delta\pi, \\ \tau_\pi \partial_t \delta\pi + \delta\pi &= -\frac{4\eta}{3} \partial_z \delta v - \frac{2}{3} \ell_{\pi V} \partial_z \delta V,\end{aligned}\quad (52)$$

where the cross couplings read (for an UR classical gas):

$$\ell_{V\pi} = \sum_{r \neq 1} \tau_{0r}^{(1)} \left(\frac{\beta J_{r+2,1}}{\epsilon + P} - \mathcal{C}_{r-1}^{(2)} \right), \quad \ell_{\pi V} = \frac{2}{5} \sum_r \tau_{0r}^{(2)} \mathcal{C}_{r+1}^{(1)}. \quad (53)$$

- ▶ In RTA, $\ell_{V\pi} = \tau_R \left(\frac{\beta J_{21}}{\epsilon + P} - \mathcal{C}_{-1}^{(2)} \right)$ and $\ell_{\pi V} = \tau_R \mathcal{C}_1^{(1)}$ both vanish:

$$\begin{aligned}J_{21} = nT = \frac{1}{3}\epsilon, \quad \mathcal{C}_{-1}^{(2)} = \frac{\alpha_{-1}^{(2)}}{\alpha_0^{(2)}} = \frac{\beta}{4} &\Rightarrow \ell_{V\pi} = 0, \\ \kappa_1 = \alpha_1^{(1)} = 0, \quad \mathcal{C}_1^{(1)} = \frac{\alpha_1^{(1)}}{\alpha_0^{(1)}} = 0 &\Rightarrow \ell_{\pi V} = 0.\end{aligned}\quad (54)$$

- ▶ We aim to control independently 4 t.c.: κ , η , $\ell_{V\pi}$ and $\ell_{\pi V}$.

Example: shear-diffusion coupling

[VEA, Wagner, PRD 110 (2024) 056002]

- ▶ To fix κ , η , $\ell_{V\pi}$, $\ell_{\pi V}$, we use $(N_1, N_2, s_1, s_2) = (1001)$ having

$$2(N_1 + s_1 + N_2 + s_2) = 4 \text{ degrees of freedom.} \quad (55)$$

- ▶ We take $\mathcal{A}_S^{(1)} = 1/\tau_V$ with $\tau_V = 12\kappa/\beta P$.
- ▶ Introducing the notation

$$H = \frac{5\eta}{4\tau_\pi P}, \quad L_{V\pi} = \frac{4\ell_{V\pi}}{\beta\tau_V}, \quad L_{\pi V} = \frac{5\beta\ell_{\pi V}}{8\tau_\pi}, \quad (56)$$

we have the constraint $H = 1 + L_{V\pi}L_{\pi V}$, i.e.

$$\tau_\pi = \frac{\tau_R}{1 + L_{V\pi}L_{\pi V}}, \quad (57)$$

where we take $\tau_R = 5\eta/4P$.


- ▶ Then, the matrix reads:

$$\mathcal{A}_S^{(2)} = \frac{1 - \alpha}{\alpha H \tau_\pi (1 - \alpha H)} \begin{pmatrix} H - L_{\pi V} & -\frac{\beta}{4}x \\ -\frac{4}{\beta}L_{\pi V} & H(1 - L_{V\pi}) - x \end{pmatrix}, \quad (58)$$

with $x = H(1 - \alpha - L_{V\pi}) - L_{\pi V} - \frac{1-H}{1-\alpha}$ and $\alpha = 1/2$.

Example: shear-diffusion coupling

[VEA, Wagner, PRD **110** (2024) 056002]





► We first consider $\ell_{\pi V} = 0$ (left panel) and $\ell_{V\pi} = 0$ (right panel):

$$\ell_{\pi V} = 0 : \quad \mathcal{A}_S^{(2)} = \frac{1}{\tau_\pi} \begin{pmatrix} 2 & -\frac{\beta}{4}(1 - 2L_{V\pi}) \\ 0 & 1 \end{pmatrix}, \quad \ell_{V\pi} = 0 : \quad \mathcal{A}_S^{(2)} = \frac{2}{\tau_\pi} \begin{pmatrix} 1 - L_{\pi V} & -\beta(\frac{1}{2} - L_{\pi V}) \\ -4\beta L_{\pi V} & \frac{1}{2} + L_{\pi V} \end{pmatrix}. \quad (59)$$

► Very good agreement with hydro observed!

Example: shear-diffusion coupling

[VEA, Wagner, PRD 110 (2024) 056002]



- The requirement $\partial_\mu S^\mu \geq 0$ imposes

$$\frac{\ell_{V\pi}}{\kappa} + \frac{\ell_{\pi V}}{2\eta T} = 0 \quad \Rightarrow \quad L_{\pi V} = -3H L_{V\pi}. \quad (60)$$

- In this case, the Shakhov matrix reads:

$$\mathcal{A}_S^{(2)} = \frac{2}{\tau_\pi(2-H)} \begin{pmatrix} 1 + 3L_{V\pi} & \frac{\beta}{8}(12L_{V\pi}^2 - 4L_{V\pi} - 1) \\ \frac{12}{\beta}L_{V\pi} & 6L_{V\pi}^2 - 3L_{V\pi} + \frac{1}{2} \end{pmatrix}, \quad (61)$$

- Again, very good agreement with hydro observed!

Section 9

Application: Ultrarelativistic hard spheres (Riemann problem)

Ultrarelativistic hard spheres (URHS)

- The t.c. of the URHS model are:

[Wagner, VEA, Molnár, PRD **109** (2024) 056018]
[Wagner, Palermo, VEA, PRD **106** (2022) 016013]

$\kappa\sigma$	$\tau_V[\lambda_{\text{mfp}}]$	$\delta_{VV}[\tau_V]$	$\ell_{V\pi}[\tau_V] = \tau_{V\pi}[\tau_V]$	$\lambda_{VV}[\tau_V]$	$\lambda_{V\pi}[\tau_V]$
0.15892	2.0838	1	0.028371β	0.89862	0.069273β

$\eta\sigma\beta$	$\tau_\pi[\lambda_{\text{mfp}}]$	$\delta_{\pi\pi}[\tau_\pi]$	$\ell_{\pi V}[\tau_\pi]$	$\tau_{\pi V}[\tau_\pi]$	$\tau_{\pi\pi}[\tau_\pi]$	$\lambda_{\pi V}[\tau_\pi]$
1.2676	1.6557	$4/3$	$-0.56960/\beta$	$-2.2784/\beta$	1.6945	$0.20503/\beta$

- The t.c. of RTA with $\eta_R = \eta_{\text{HS}}$ are

$\kappa\sigma$	$\tau_V[\lambda_{\text{mfp}}]$	$\delta_{VV}[\tau_V]$	$\ell_{V\pi}[\tau_V] = \tau_{V\pi}[\tau_V]$	$\lambda_{VV}[\tau_V]$	$\lambda_{V\pi}[\tau_V]$
0.13204	1.5845	1	0	$3/5$	$\beta/16$

$\eta\sigma\beta$	$\tau_\pi[\lambda_{\text{mfp}}]$	$\delta_{\pi\pi}[\tau_\pi]$	$\ell_{\pi V}[\tau_\pi]$	$\tau_{\pi V}[\tau_\pi]$	$\tau_{\pi\pi}[\tau_\pi]$	$\lambda_{\pi V}[\tau_\pi]$
1.2676	1.5845	$4/3$	0	0	$10/7$	0

- RTA-HS mismatch for almost all coefficients, except $\delta_{VV} = \tau_V$ and $\delta_{\pi\pi} = 4\tau_\pi/3$, which are fixed for an UR gas.
- To align all transport coefficients, we need 11 parameters!

Various (N_1, N_2, s_1, s_2) models



- ▶ A Shakhov model with (N_1, N_2, s_1, s_2) provides $2(N_1 + N_2 + s_1 + s_2)$ params.
- ▶ To test the effect of various t.c., we employed several models:
- ▶ AW: τ_R is used to fix $\eta_R = \eta_{\text{HS}}$.
- ▶ (1000): Fixes η and κ .
- ▶ (1001): discussed previously, fixes $(\kappa, \eta, \ell_{V\pi}, \ell_{\pi V})$.
- ▶ (1012): has $2 \times 4 = 8$ free entries and fixes everything except λ_{VV} and $\lambda_{V\pi}$.
- ▶ (2102): has $2 \times 5 = 10$ free entries and fixes everything.

Models used


Model	$\eta\sigma\beta$	$\tau_\pi/\lambda_{\text{mfp}}$	$\ell_{\pi V}/\tau_\pi$	$\tau_{\pi\pi}/\tau_\pi$	$\beta\lambda_{\pi V}/\tau_\pi$
HS	1.2676	1.6557	−0.56960	1.6945	0.20503
AW	1.2676	1.5845	0	1.4286	0
1000	1.2676	1.5845	0	1.4286	0
1001	1.2676	1.6457	−0.56960	1.7607	0
1012	1.2676	1.6557	−0.56960	1.6945	0.20503
2012	1.2676	1.6557	−0.56960	1.6945	0.20503

Model	$\kappa\sigma$	$\tau_V/\lambda_{\text{mfp}}$	$\ell_{V\pi}/\beta\tau_V$	λ_{VV}/τ_V	$\lambda_{V\pi}/\beta\tau_V$
HS	0.15892	2.0838	0.028371	0.89862	0.069273
AW	0.13204	1.5845	0	0.6	0.0625
1000	0.15892	1.5845	0	0.6	0.0625
1001	0.15892	1.9070	0.028371	0.6	0.055407
1012	0.13204	2.0838	0.028371	0.762023	0.062933
2012	0.15892	2.0838	0.028371	0.89862	0.069273

Sod shock tube: convergence properties





- ▶ To validate the numerical scheme, we compared AW results to BAMPS for various fixed η/s .
- ▶ As $\eta/s \rightarrow 0$, our results approach the inviscid (analytical) solution.
- ▶ AW and all Shakhov implementations are in excellent agreement w. BAMPS for the eq. quantits. (n, P, u) .



[Bouras et al, PRC **82** (2010) 024910]

Sod shock tube: Comparison to BAMPS



[Bouras et al, PRC 82 (2010) 024910]



- ▶ In the frame of the Sod shock tube, we considered a comparison to BAMPS for hard-sphere interactions.
- ▶ Using τ_R to tune η , shear comes out well with AW and Shakhov.
- ▶ For diffusion: 1000 \equiv first-order Shakhov underestimates peak.
- ▶ All high-order Shakhov models perform well!

Heat flow problem: Comparison to BAMPS



[DNBMXRG, PRD **89** (2014) 074005]



- ▶ Case 1: const. initial λ , pressure jump.
- ▶ All models recover π/P .
- ▶ For q^z , both AW (fixing only η) and 1000 (fixing η and κ) fail.
- ▶ All high-order Shahkov models perform well!



Heat flow problem: Comparison to BAMPS

[DNBMXRG, PRD **89** (2014) 074005]



- ▶ Case 2: cons. initial P , jump in λ .
- ▶ AW and 1000 give $\pi/P = 0$; all high-order models recover π/P .
- ▶ For q^z , AW is off by $\simeq 10\%$, while 1000 and high-order Shahkov models perform well!

IReD Supremacy: Problem with DNMR



- So far, we used the IReD method for the t.coeffs computation.
- Now we tune the S-model to capture the $O(\text{Re}^{-1}\text{Kn})$ t.coeffs to the DNMR values, ignoring the $O(\text{Kn}^2)$ t.coeffs.
- While π is recovered well, in all S-models the DNMR coefficients lead to wrong results for q^z .

Section 10

Code availability

Code availability

- ▶ The kinetic equation is solved using a discrete velocity method algorithm based on the relativistic lattice Boltzmann method.
- ▶ The source code, run scripts, as well as plotting scripts are available to download from CodeOcean, as follows:
 - 0 + 1-D massless Bjorken flow: DOI: 10.24433/CO.5625382.v2
[VEA *et al*, Nature Comput. Sci. 2 (2022) 641]
 - 0 + 1-D massive Bjorken flow (hydro, aHydro, Boltzmann-RTA): DOI: 10.24433/CO.1942625.v1
[VEA, Molnár, Rischke, PRD 109 (2024) 076001]
 - First-order Shakhov model (Bjorken flow, longitudinal waves): DOI: 10.24433/CO.6267589.v1
[VEA, Molnár, PLB 855 (2024) 138795]
 - High-order Shakhov model (Bjorken flow, longitudinal waves, shock waves): DOI: 10.24433/CO.8322373.v1
[VEA, Wagner, PRD 110 (2024) 056002]

Kinetic solver: $1 + 1$ -D flows

- For $1 + 1$ -D flows, the kinetic equation reduces to

$$k^t \partial_t f_{\mathbf{k}} + k^z \partial_z f_{\mathbf{k}} = -\frac{E_{\mathbf{k}}}{\tau_R} (f_{\mathbf{k}} - f_{S\mathbf{k}}). \quad (62)$$

- We parametrize $f_{\mathbf{k}} \equiv f(x^\mu; m_\perp, v^z, \varphi_{\mathbf{k}})$, with

$$\begin{pmatrix} k^t \\ k^z \end{pmatrix} = m_\perp \begin{pmatrix} \cosh y \\ \sinh y \end{pmatrix} = \frac{m_\perp}{\sqrt{1 - v_z^2}} \begin{pmatrix} 1 \\ v^z \end{pmatrix}, \quad \begin{pmatrix} k^x \\ k^y \end{pmatrix} = k_\perp \begin{pmatrix} \cos \varphi_{\mathbf{k}} \\ \sin \varphi_{\mathbf{k}} \end{pmatrix}, \quad (63)$$

where $m_\perp = \sqrt{\mathbf{k}_\perp^2 + m^2}$ is the transverse mass, $y = \tanh^{-1} v^z$ is the rapidity, and $v^z = k^z / k^t$.

- Assuming $u^\mu \partial_\mu = \gamma(\partial_t + \beta^z \partial_z)$, Eq. (62) leads to

$$\partial_t f_{\mathbf{k}} + v^z \partial_z f_{\mathbf{k}} = -\frac{\gamma(1 - \beta^z v^z)}{\tau_R} (f_{\mathbf{k}} - f_{S\mathbf{k}}). \quad (64)$$

Kinetic solver: Rapidity-based moments

- ▶ Going from $\mathbf{k} = (k^x, k^y, k^z)$ to $(m_\perp, v^z, \varphi_k)$ implies:

$$\int \frac{d^3 k}{k^0} \rightarrow \int_{-1}^1 \frac{dv^z}{1 - v_z^2} \int_0^{2\pi} d\varphi_k \int_m^\infty dm_\perp m_\perp . \quad (65)$$

- ▶ The m_\perp and $\varphi_{\mathbf{k}}$ dofs can be integrated out by introducing *rapidity-based moments*:

$$F_n(v^z) = \frac{g}{(2\pi)^3} \int_0^{2\pi} d\varphi_k \int_m^\infty \frac{dm_\perp m_\perp^{n+1}}{(1 - v_z^2)^{(n+2)/2}} f_{\mathbf{k}} . \quad (66)$$

- ▶ For the longitudinal waves and shock waves problems, Eq. (64) can be integrated w.r.t. m_\perp and $\varphi_{\mathbf{k}}$, leading to

$$\frac{\partial F_n}{\partial t} + v^z \frac{\partial F_n}{\partial z} = - \frac{\gamma(1 - \beta^z v^z)}{\tau} (F_n - F_n^S) . \quad (67)$$

- ▶ The equation is closed since all required macroscopic quantits. entering $f_{S\mathbf{k}} \rightarrow F_n^S$ can be recovered from F_n :

$$\begin{pmatrix} N_r^t \\ N_r^z \end{pmatrix} = \int_{-1}^1 dv^z \begin{pmatrix} 1 \\ v^z \end{pmatrix} (u \cdot v)^r F_{r+1} , \quad \begin{pmatrix} T_r^{tt} \\ T_r^{tz} \\ T_r^{zz} \end{pmatrix} = \int_{-1}^1 dv^z \begin{pmatrix} 1 \\ v^z \\ v_z^2 \end{pmatrix} (u \cdot v)^r F_{r+2} . \quad (68)$$

Kinetic solver: Non-conformal Bjorken flow

- ▶ Due to the symmetries of Bjorken flow, it is convenient to employ (τ, η) , defined by

$$t = \tau \cosh \eta, \quad z = \tau \sinh \eta. \quad (69)$$

- ▶ Due to boost invariance, $f_{\mathbf{k}}$ depends on y and η only through $y - \eta$.
- ▶ Then, $f_{\mathbf{k}} \rightarrow f(\tau; m_{\perp}, \varphi_{\mathbf{k}}, v^z)$, where $v^z = \tanh(y - \eta)$ instead of $\tanh y$.
- ▶ The kinetic eq. for Bjorken flow becomes:

$$\frac{\partial f_{\mathbf{k}}}{\partial \tau} - \frac{v^z(1 - v_z^2)}{\tau} \frac{\partial f_{\mathbf{k}}}{\partial v^z} = -\frac{1}{\tau_R} (f_{\mathbf{k}} - f_{S\mathbf{k}}). \quad (70)$$

- ▶ Defining again the *rapidity-based moments*,

$$F_n(v^z) = \frac{g}{(2\pi)^3} \int_0^{2\pi} d\varphi_k \int_m^{\infty} \frac{dm_{\perp} m_{\perp}^{n+1}}{(1 - v_z^2)^{(n+2)/2}} f_{\mathbf{k}}, \quad (71)$$

one obtains

$$\frac{\partial F_n}{\partial \tau} + \frac{1}{\tau} [1 + (n - 1)v_z^2] F_n - \frac{1}{\tau} \frac{\partial [v^z(1 - v_z^2) F_n]}{\partial v^z} = -\frac{1}{\tau_R} (F_n - F_n^S). \quad (72)$$

- ▶ The equation is again closed w.r.t. n .

Momentum-space discretization: v^z

- ▶ v^z is discretized via the Gauss-Legendre quadrature.
- ▶ The continuous functions $F_n(v^z)$ are replaced by

$$F_{n;j} = w_j F_n(v_j^z), \quad w_j = \frac{2(1 - v_{z;j}^2)}{[(K+1)P_{K+1}(v_j^z)]^2}, \quad (73)$$

where v_j^z ($1 \leq j \leq K$) satisfy $P_K(v_j^z) = 0$

- ▶ The derivative w.r.t. v^z is replaced by the finite sum

$$\left[\frac{\partial [v^z (1 - v_z^2) F_n]}{\partial v^z} \right]_j = \sum_{j'=1}^K \mathcal{K}_{j,j'} F_{n;j'}, \quad (74)$$

where $\mathcal{K}_{j,j'}$ is obtained by projection onto Legendre polynomials:

[VEA, Blaga, PRC **98** (2018) 035201]

$$\begin{aligned} \mathcal{K}_{j,j'} = w_j \sum_{m=1}^{K-3} \frac{m(m+1)(m+2)}{2(2m+3)} P_m(v_j^z) P_{m+2}(v_{j'}^z) \\ - w_j \sum_{m=1}^{K-1} \frac{m(m+1)}{2} P_m(v_j^z) \left[\frac{(2m+1)P_m(v_{j'}^z)}{(2m-1)(2m+3)} + \frac{m-1}{2m-1} P_{m-2}(v_{j'}^z) \right]. \end{aligned} \quad (75)$$

Section 11

Conclusions

Conclusions

- ▶ Shakhov model generalized for the relativistic Anderson-Witting RTA, allowing ζ , κ and η to be controlled independently.
- ▶ Numerical simulations of the Bjorken flow and of sound waves damping confirmed that the model is robust.
- ▶ Extending the Shakhov model allows 2nd-order t. coeffs. to be controlled \Rightarrow agreement with BAMPS in Sod shock tube.
- ▶ This work was supported through a grant of the Ministry of Research, Innovation and Digitization, CNCS - UEFISCDI, project number PN-III-P1-1.1-TE-2021-1707, within PNCDI III.