

DESI shakes up the Dark Universe

Satya Gontcho A Gontcho (Berkeley Lab, University of Virginia)

On behalf of the DESI Collaboration

September 11, 2025

New Physics from Galaxy Clustering at GGI



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

The expanding universe

Hubble & Lemaître, 1929



Velocity increases with Distance



Distance

Velocity

**Expanding
Universe**

Velocity = $H \times \text{Distance}$

$H \sim 70 \text{ km/s/Mpc}^*$
(7% per Giga-year)
*current estimation



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

The expanding universe

Hubble law

$$v = H \cdot D$$

Spectroscopy

Photometry

Redshift $z = \frac{\lambda - \lambda_0}{\lambda_0}$
(Doppler effect: $z = v/c$)


Source of known luminosity

$$\mathcal{L}_{\text{obs}} \propto \frac{\mathcal{L}_0}{D^2}$$

$z=0.05$



$z=0$



Cepheids (Period – luminosity)
Type Ia Supernovae (Peak luminosity)

Variation of H? $\Rightarrow$ Study at different epochs (= redshifts)





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

The expanding universe

Hubble diagram
distance – redshift relation



SN Ia
(known luminosity)

SN Ia are weaker (~20%)
for a given redshift
Hence more distant (~10%)



Accelerated expansion

Incompatible with matter domination



DARK ENERGY!



2011 Nobel Prize

Perlmutter et al., 1998



Riess et al., 1998




DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Standard model of cosmology

U.S. Department of Energy Office of Science



Accelerated expansion



Cosmic Microwave Background

Physics we know
+ cosmological model


successfully predicts most
observations of last 20 yrs



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Standard model of cosmology – Λ CDM



Two main components of unknown nature

- **Dark Matter** (galaxy formation, gravitational lensing, rotation curves, ...)
- **Dark Energy** (late-time acceleration)

Other missing information

- **Neutrino masses**



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Dark Energy

U.S. Department of Energy Office of Science

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$$

Geometry ($G_{\mu\nu}$)?
Cosmological constant Λ

**Accelerated
expansion**

Energy content ($T_{\mu\nu}$)?
Additional component

$$w = \frac{p}{\rho}$$

Modified gravity?
Beyond general relativity

Baryon Acoustic Oscillations



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT



U.S. Department of Energy Office of Science

Baryon Acoustic Oscillations (BAO)

Propagation of baryon-photon over-density sound waves in primordial plasma

At recombination ($z \sim 1100$): $p + e^- \rightarrow H$

- Plasma evolves from optically thick to optically thin
- Baryons decouple from photons
- Waves stall



Residual spherical shell $\longrightarrow$ Peak in clustering of matter



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT



U.S. Department of Energy Office of Science

Baryon Acoustic Oscillations (BAO)

Propagation of baryon-photon over-density sound waves in primordial plasma

At recombination ($z \sim 1100$): $p + e^- \rightarrow H$

- Plasma evolves from optically thick to optically thin
- Baryons decouple from photons
- Waves stall



Residual spherical shell $\longrightarrow$ Peak in clustering of matter

Size of feature = distance sound wave traveled

Preferred 3D scale $r_s \sim 150 \text{ kpc}$ (at recombination)
 $r_s \sim 150 \text{ Mpc}$ (today)




DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

Baryon Acoustic Oscillations (BAO)


Imprint of fluctuations in primordial plasma
→ Standard Ruler to measure distances



DESI year 1



@ Claire Lamman / DESI collaboration







DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

The BAO standard ruler

Artist's view of BAO



$D_M(z)$ and $H(z)$ encode **expansion history** of the Universe



DARK ENERGY SPECTROSCOPIC INSTRUMENT

U.S. Department of Energy Office of Science

DESI



Science & Technology
Facilities Council


Thanks to our sponsors and
72 Participating Institutions!



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Photometry before DESI




Photometry
of 1/3 of the Sky



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

DESI Legacy Surveys




18k sq. deg. in *grz* using three telescopes (Bok, Mayall & Blanco)



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Viewer: legacysurvey.org/viewer






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Viewer: legacysurvey.org/viewer

U.S. Department of Energy Office of Science






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Viewer: legacysurvey.org/viewer

U.S. Department of Energy Office of Science




Satya Gontcho A Gontcho (LBNL/UVA)



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

DESI: spectroscopic galaxy survey




Photometry
of 1/3 of the Sky



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science



DESI targets: 40 million galaxies & quasars!





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

The Lyman- α Forest at $z > 2.1$



$$F = e^{-\tau}$$

$$\tau \propto n_{HI}$$

- **Quasars** visible to high redshift ($z \sim 5$)
- Absorption of Quasar spectrum by neutral H in IGM
- Transmitted flux fraction F : **proxy for neutral H density**



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

DESI instrument

Mayall telescope
at Kitt Peak Observatory (AZ)



Satya Gontcho A Gontcho (LBNL/UVA)




**DARK ENERGY
SPECTROSCOPIC
INSTRUMENT**

U.S. Department of Energy Office of Science

DESI instrument

**Focal plane:
5000 fiber positioners
(high multiplexing)**



**7 deg²
field of view**



**4m mirror
(large collecting area)**






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

DESI instrument

40m-long
optical fibers



10 3-band spectrographs



7 deg²
field of view



4m mirror
(large collecting area)






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

DESI instrument



10 3-band spectrographs [360nm – 980nm]



$\text{Ly}\alpha$ $\lambda 121.6$ nm
down to $z = 2.0$



$[\text{OII}] \lambda 373$ nm
up to $z = 1.6$





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

DESI DR2: data & analysis



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

DESI Data Release 1 footprint

DR1 analysis sample

420,000 Lyman- α forests
5.7 million galaxies and quasars

2 to 3x larger than SDSS (20 years)



DESI 2024 II: Samples ([arXiv:2404.03002](https://arxiv.org/abs/2404.03002))



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

DESI Data Release 2 footprint

U.S. Department of Energy Office of Science

DR1 analysis sample


420,000 Lyman- α forests
5.7 million galaxies and quasars

2 to 3x larger than SDSS (20 years)

DR2 analysis sample

820,000 Lyman- α forests
14.3 million galaxies and quasars

2 (QSO) to 3 (ELG) x DR1






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Blinding strategy

U.S. Department of Energy Office of Science

Blinded analysis to prevent confirmation bias

- Catalog-level for Galaxies & quasars: redshifts & weights
- Cosmology-level for Lyman-alpha forest: shift of BAO peak



Procedure

Determine analysis parameters
& validate choices based on

- Simulated data (*mocks*)
- Data splits (*blinded data*)

Robustness tests


- Variations in data vector
- Methods to compute correlations & covariances
- BAO modeling (priors, broadband, ...)
- Imaging systematics
- Data splits



Galaxy clustering

Dominant systematics

- Theoretical modeling
- Galaxy-halo connection
- Fiducial cosmology



Total systematic (tracer-dependent)

$$\Delta\alpha_{\text{iso}} = 0.14\% \text{ to } 0.22\%$$
$$\Delta\alpha_{\text{AP}} = 0.22\% \text{ to } 0.33\%$$

Induced increase of σ_{tot} over σ_{stat}

$$\Delta\sigma(\alpha_{\text{iso}}) = 1 - 9\% \text{ (BGS - LRG3+ELG1)}$$
$$\Delta\sigma(\alpha_{\text{AP}}) = 0.1 - 2\% \text{ (QSO - LRG3+ELG1)}$$

Ly α forest clustering


Dominant systematics

- non-linear evolution of BAO peak

Total systematic

$$\Delta\alpha_{\parallel} = 0.3\%$$

$$\Delta\alpha_{\perp} = 0.3\%$$



Induced increase of σ_{tot} over σ_{stat}

$$\Delta\sigma(\alpha_{\text{iso}}) = 9\% \text{ (Ly}\alpha\text{)}$$

Statistics-limited!





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

DR2 clustering measurements

LRG+ELG ($0.8 < z < 1.1$)

15 σ detection of BAO
at $z_{\text{eff}} = 0.93$

U.S. Department of Energy Office of Science







DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

DESI DR2 BAO

BAO data: $\Delta\theta$ and $\Delta z \longrightarrow D_M / r_d$ and D_H / r_d

$$D_V = (z D_M(z)^2 D_H(z))^{1/3}$$

Ω_M and $H_0 r_d$



Agreement & complementarity
between tracers

Aggregated precision on BAO distance scale: 0.3%

(vs. 0.6% for final SDSS)

Satya Gontcho A Gontcho (LBNL/UVA)



DESI DR2 BAO

BAO data: $\Delta\theta$ and $\Delta z \longrightarrow D_M / r_d$ and D_H / r_d


$$D_V = (z D_M(z)^2 D_H(z))^{1/3}$$

Ω_M and $H_0 r_d$

$$\Omega_m = 0.298 \pm 0.0086 \quad (\mathbf{2.9\%})$$

$$h r_d = 101.54 \pm 0.73 \text{ Mpc} \quad (\mathbf{0.7\%})$$

DESI



Agreement & complementarity
between tracers



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Λ CDM: DESI DR2 vs. CMB

DESI DR2 results II: BAO (arXiv:2503.14738)


U.S. Department of Energy Office of Science

DESI DR2 BAO is:

- Consistent with **DESI DR1**
- 2.3σ from the **CMB** (was 1.9σ with DESI DR1)

CMB:

- primary CMB from Planck PR4 (CamSpec)
- CMB lensing from Planck PR4 + ACT DR6






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Λ CDM: DESI DR2 vs. Supernovae

- **DESI DR2** consistent with DESI DR1
- **DESI DR2** is lower than the **CMB**
- **DESI DR2** is lower than Supernovae:
 - 1.7σ lower than **Pantheon+**
 - 2.1σ lower than **Union3**
 - 2.9σ lower than **DESY5**





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science


Data consistency

Differences in H_0 & Ω_m
between DESI BAO, CMB and SN

expected

when dynamic dark energy universe
fitted assuming Λ CDM

Tang+ (arXiv:2412.04430)



Expansion rate of the Universe

H_0



**DARK ENERGY
SPECTROSCOPIC
INSTRUMENT**

H_0

U.S. Department of Energy Office of Science


Extrapolation to $z=0$
(normalization) of
early-universe data



Distance-ladder calibration
approach in
late-time universe

Early-time

Late-time Universe



Riess & Breuval 2023



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

H_0

U.S. Department of Energy Office of Science

BAO data: $\Delta\theta$ and Δz $\longrightarrow$ D_M / r_d and D_H / r_d $\longrightarrow$ Ω_M and $H_0 r_d$


Using r_d from BBN ($\Omega_b \rightarrow r_d$)

$$H_0 = 68.51 \pm 0.58 \text{ km s}^{-1} \text{ Mpc}^{-1} \quad (0.8\%)$$

DESI + BBN

$\rightarrow$ 0.8% precision on H_0 , independent of CMB

$\rightarrow$ 4.5 σ tension with SH0ES





U.S. Department of Energy Office of Science

Dark Energy



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Dark Energy

U.S. Department of Energy Office of Science

Equation of state $P = w\rho$


Dynamic dark energy

(Chevalier & Polarski 2001, Linder 2003)

$$w(z) = w_0 + w_a \frac{z}{1+z}$$

Cosmological constant Λ

$$\left\{ \begin{array}{l} w_0 = -1 \text{ and } w_a = 0 \end{array} \right.$$





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

Evolving Dark Energy

DESI DR2 results II: BAO (arXiv:2503.14738)

$$w(a) = w_0 + (1 - a)w_a$$

- Degeneracy in $w_0 - w_a$ plane with BAO alone
- DESI DR2 within 2σ of Λ CDM





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science


Evolving Dark Energy

DESI DR2 results II: BAO (arXiv:2503.14738)

$$w(a) = w_0 + (1 - a)w_a$$

- **3.1 σ** preference for evolving dark energy with DESI DR2 + CMB

$$\left. \begin{aligned} w_0 &= -0.42 \pm 0.21 \\ w_a &= -1.75 \pm 0.58 \end{aligned} \right\} \text{DESI + CMB}$$





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Evolving Dark Energy


DESI DR2 results II: BAO ([arXiv:2503.14738](#))

$$w(a) = w_0 + (1 - a)w_a$$

DESI + CMB + **Pantheon+**: **2.8 σ**

DESI + CMB + **Union3**: **3.8 σ**

DESI + CMB + **DES-SN5Yr**: **4.2 σ**





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Result robustness

DESI DR2 results II: BAO (arXiv:2503.14738)

U.S. Department of Energy Office of Science

CMB alternatives limited to early-time information


- Early-Universe priors on $(\theta_*, \omega_b, \omega_{bc})$ derived from CMB:

$$\text{DESI} + (\theta_*, \omega_b, \omega_{bc})_{\text{CMB}} \Rightarrow 2.4\sigma$$

- CMB without lensing:

$$\text{DESI} + \text{CMB (no lensing)} \Rightarrow 2.7\sigma$$

Weaker preference (3.1σ for DESI + CMB)
but similar posteriors





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Result robustness

DESI DR2 results II: BAO ([arXiv:2503.14738](#))

U.S. Department of Energy Office of Science


Constraint limited to low-redshift probes

- Replacing CMB with DESY3 3×2 pt (weak lensing + galaxy clustering)

DESI + DESY3 (3×2 pt) $\Rightarrow 2.2\sigma$

DESI + DESY3 (3×2 pt) + DESY5 $\Rightarrow 3.3\sigma$

Preference for same region






DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Isotropic BAO
distance measurement

Supernovae
distance modulus

The nature of the evidence



- **Λ CDM model** can fit DESI BAO
- DESI at $z < 1$ prefer distances 1-2% lower than CMB prediction




DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Isotropic BAO
distance measurement

Supernovae
distance modulus

The nature of the evidence



- **Λ CDM model** can fit DESI BAO
- DESI at $z < 1$ prefer distances 1-2% lower than CMB prediction

- **Λ CDM model** can fit SNe
- Tension with DESI and CMB

⇒ No good **Λ CDM** fit
to DESI BAO, CMB & SN
simultaneously



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

Isotropic BAO
distance measurement

Supernovae
distance modulus

The nature of the evidence

DESI DR2 results II: BAO (arXiv:2503.14738)



Not have enough
freedom in **wCDM** to fit
BAO, CMB and SN
simultaneously either



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT


U.S. Department of Energy Office of Science

Isotropic BAO
distance measurement

Supernovae
distance modulus

The nature of the evidence

DESI DR2 results II: BAO (arXiv:2503.14738)



w_0w_a CDM fit to BAO+SN
also a good fit to CMB

Dark Energy model

Observations = distance(z), not $w(z)$!




DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

Dark energy model

DESI DR2 results II: BAO ([arXiv:2503.14738](#))

DESI supporting paper Lodha+ ([arXiv:2503.14743](#))

U.S. Department of Energy Office of Science



$w < -1$ at high z : increasing dark energy density with time !

Could indicate more complex dark sector

Maximum dark energy density reached at $z \sim 0.45$ (phantom crossing)




DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Weakening dark energy

DESI DR2 results II: BAO ([arXiv:2503.14738](#))

DESI supporting paper Lodha+ ([arXiv:2503.14743](#))





DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science


Dark energy model

DESI DR2 results II: BAO ([arXiv:2503.14738](#))

DESI supporting paper Lodha+ ([arXiv:2503.14743](#))

Three classes of dark energy

- Thawing (away from $w = -1$)
- Emergent (from $\rho = 0$, never crosses $w = -1$)
- Mirage ($\langle w \rangle = -1$)



Improvement over LCDM
DESI BAO + CMB + SN (DESY5)

Dark Energy Model	$\Delta\chi^2$
Thawing	-12.0
Emergent	-3.9
Mirage	-21.3
$w_0 w_a$	-21.4




DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Dark energy model

DESI DR2 results II: BAO ([arXiv:2503.14738](#))

DESI supporting paper Lodha+ ([arXiv:2503.14743](#))



Binned reconstruction of $w(z)$

- Consistent with w_0w_a CDM
- Weaker than Λ CDM at small z ($>3\sigma$)
- Preference for "phantom" at large z

Neutrino masses



DARK ENERGY
SPECTROSCOPIC
INSTRUMENT

U.S. Department of Energy Office of Science

Upper bounds on neutrino masses

DESI DR2 results II: BAO ([arXiv:2503.14738](#))



In Λ CDM, Σm_ν changes angular diameter distance to last scattering,
degenerate with Ω_m , H_0 ...
BAO breaks degeneracy

$$\Sigma m_\nu < 0.064 \text{ eV} \quad (95\%, \text{ DESI (BAO)+CMB})$$

➡ Tightest constraint to-date (in Λ CDM)

In $w_0 w_a$ CDM, relaxed to

$$\Sigma m_\nu < 0.163 \text{ eV} \quad (95\%, \text{ DESI (BAO)+CMB})$$



Thank you!



© Claire Lamman

Thank you!



© Claire Lamman


DESI will observe until 2029!

- extended southern footprint by ~20% (down to DEC -20)
- Increase overlap with LSST, Simons Observatory by ~50%



DARK ENERGY SPECTROSCOPIC INSTRUMENT

U.S. Department of Energy Office of Science






FIG. 6. DESI DR2 BAO measurements compared to DR1 measurements. In each panel, DR1 and DR2 results are shown with empty and filled circles, respectively. Results are shown for the following tracers in order of increasing redshift: BGS (yellowgreen points), LRG1 (orange), LRG2 (orangered), LRG3+ELG1 (slateblue), ELG2 (steelblue), QSO (seagreen) and Ly α (purple) [63, Appendix D]. The distances are normalized by the DESI DR2 best-fit Λ CDM model predictions (black solid lines). All systematic errors are included. The *Planck* Λ CDM predictions are shown with brown dashed lines with 68% confidence intervals in the brown shaded region. A small artificial offset in redshift has been applied to the DESI DR1 data points for a clearer comparison.