


Heat flow in nonequilibrium steady states from CFT

Benjamin Doyon
King's college London

based on arXiv:1202.0239 with Denis Bernard

Florence, 23 April 2012

Physical situation



$R \gg v_F(t - t_0) \gg$ observation length, microscopic scales



Physical situation

$$\langle \dots \rangle_{\text{ness}} = \lim_{t_0 \rightarrow -\infty} \lim_{R \rightarrow \infty} \frac{\text{Tr} (e^{iHt_0} \rho_0 e^{-iHt_0} \dots)}{\text{Tr} (\rho_0)}$$

$$\rho_0 = e^{-\beta_l H^l - \beta_r H^r}$$

$$H = H^l + H^r + H_{\text{contact}}$$

Observables supported on a finite region

Scaling limit: (relativistic) QFT

Linear dispersion relation: CFT

Energy



wave number

Relativistic dispersion relation: QFT

Energy



wave number

Description of the steady state

$$\langle \dots \rangle_{\text{ness}} = \frac{\text{Tr} (e^{-Y} \dots)}{\text{Tr} (e^{-Y})}$$

Operator Y :

- Commutes with the Hamiltonian H
- «Asymptotically looks like» $\beta_l H^l + \beta_r H^r$

★ Formal definition first proposed by Hershfield (PRL 1993) (case where both temperatures are the same and something else is flowing, like a charge)

★ Studied widely for charge transfer in impurity systems

Description of the steady state


[DB & BD]

$$Y = \beta_l \int_0^\infty d\theta E_\theta A(\theta)^\dagger A(\theta) + \beta_r \int_{-\infty}^0 d\theta E_\theta A(\theta)^\dagger A(\theta)$$

Total energy of
right-moving
asymptotic particles

Total energy of
left-moving
asymptotic particles

State



The energy current

$$J = \langle p^1(x) \rangle_{\text{ness}}$$

$$= \lim_{R \rightarrow \infty} R^{-d} \frac{\text{Tr}_R (e^{-Y} P^1)}{\text{Tr}_R (e^{-Y})}$$

$$= \lim_{R \rightarrow \infty} R^{-d} \frac{\text{Tr}_R (e^{-Y_+} P_+^1)}{\text{Tr}_R (e^{-Y_+})} + \lim_{R \rightarrow \infty} R^{-d} \frac{\text{Tr}_R (e^{-Y_-} P_-^1)}{\text{Tr}_R (e^{-Y_-})}$$

$$\mathcal{H} = \mathcal{H}_+ \otimes \mathcal{H}_-$$

$$Y = Y_+ + Y_-$$

$$P^1 = P_+^1 + P_-^1$$

The energy current

Using the fact that the energy is unchanged under change of sign of a momentum component:

$$J = f(\beta_l) - f(\beta_r)$$

Towards the conformal (gapless) point:

$$J = \alpha m^{d-1} (\beta_l^{-2} - \beta_r^{-2})$$

1D: the CFT central charge

[DB & BD]

$$J = \frac{\pi c}{12} (\beta_l^{-2} - \beta_r^{-2}) = \frac{\pi c k_B^2}{12 \hbar} (T_l^2 - T_r^2)$$

central charge

$$T(x) = -\frac{c}{24} + \sum_{n \in \mathbb{Z}} L_n e^{-\frac{2\pi i n x}{R}}$$

Virasoro

$$H = \int dx (h_+(x) + h_-(x)) \quad h_+(x) = \frac{2\pi}{R^2} T(x)$$

$$H^{l,r} = \int dx (h_+^{l,r}(x) + h_-^{l,r}(x)) \quad h_-(x) = \frac{2\pi}{R^2} \bar{T}(x)$$

$$J = \langle h_+(x) - h_-(x) \rangle_{\text{ness}}$$

1D: the CFT central charge

Using the fact that

$$h_{\pm}(x) = \begin{cases} h_{\pm}^l(x) & (x < 0) \\ h_{\pm}^r(x) & (x > 0) \end{cases}$$

and

$$\rho_0 = e^{-\beta_l H^l - \beta_r H^r}$$

we find

$$Y = \frac{2\pi\beta_l}{R} L_0 - \frac{2\pi\beta_r}{R} \bar{L}_0$$

1D: the CFT central charge

Hence:

$$J = f(\beta_l) - f(\beta_r), \quad f(\beta) = - \lim_{R \rightarrow \infty} \frac{1}{R} \frac{d}{d\beta} \log Z(\beta)$$

where

$$Z(\beta) = \text{Tr} \left(e^{-\frac{2\pi\beta}{R} L_0} \right)$$

and we can use

$$Z(\beta) \sim N e^{\frac{\pi c R}{12\beta}}$$

Fluctuations of the energy transfer

We want to measure the fluctuations of the transfer of energy, whose «charge» can be taken as:

$$Q = \frac{1}{2} (H^l - H^r)$$



Fluctuations of the energy transfer

$$P(q, t) = \sum_{q_0} \text{Tr} \left(P_{q_0+q} e^{-iHt} P_{q_0} \rho_{\text{ness}} P_{q_0} e^{iHt} P_{q_0+q} \right)$$

$\frac{e^{-Y}}{\text{Tr}(e^{-Y})}$

$$P(\lambda, t) = \sum_q e^{i\lambda q} P(q, t)$$

$$\log P(\lambda, t) \sim t F(\lambda) + O(1)$$

Cumulant generating function
 $= -i\lambda J + \dots$

An expected fluctuation relation

$$F(\lambda) = F(i(\beta_l - \beta_r) - \lambda)$$

Equivalent to:

$$P(q, t \rightarrow \infty) = e^{(\beta_l - \beta_r)q} P(-q, t \rightarrow \infty)$$

Such a relation was argued for first measurement at $t = t_0$

Jarzynski, Wojcik (PRL 2004)

See the nice review by: Esposito, Harbola, Mukamel (RMP 2009)

More rigorous proof given in: Andrieux, Gaspard, Monnai, Tasaki (2008)

Basic ideas: Gallavoti, ...

The full counting statistics in CFT

Recall:

$$P(\lambda, t) = \sum_{q, q_0} e^{i\lambda q} \text{Tr} \left(P_{q_0+q} e^{-iHt} P_{q_0} \rho_{\text{ness}} P_{q_0} e^{iHt} P_{q_0+q} \right)$$

Use $\sum_q f(q) P_q = f(Q)$ and $P_q \propto \int d\mu e^{i\mu(Q-q)}$

$$F(\lambda) = \lim_{t \rightarrow \infty} t^{-1} \log \left[\lim_{t_0 \rightarrow -\infty} \lim_{R \rightarrow \infty} \int d\mu \left(\star \right) \right]$$

$$\frac{\text{Tr} \left(\rho_0(t_0) e^{-i\left(\frac{\lambda}{2} + \mu\right)Q} e^{i\lambda Q(t)} e^{-i\left(\frac{\lambda}{2} - \mu\right)Q} \right)}{\text{Tr} \rho_0(t_0)}$$

The full counting statistics in CFT

Parenthesis: charge transfer in free-fermion systems

- Cumulant generating function known in terms of transmission matrix: Lesovik–Levitov formula (1993,1994) (also: Klich, Schonhammer, DB & BD, . . .)
- It is observed that the same result is obtained with any fixed μ

Hence we expect to get the same result with:

$$\frac{\text{Tr} (\rho_0(t_0) e^{i\lambda Q(t)} e^{-i\lambda Q})}{\text{Tr} \rho_0(t_0)}$$

The full counting statistics in CFT

$$e^{i\lambda Q(t)} e^{-i\lambda Q} = e^{i\lambda Q + i\lambda \int_0^t dx (h_-(x) - h_+(-x))} e^{-i\lambda Q}$$

Supported on a finite region

Finitely-supported observable, can use Y-operator, get factorization:

$$F(\lambda) = f(\lambda, \beta_l) + f(-\lambda, \beta_r)$$

$$f(\lambda, \beta) = \left\langle e^{i\lambda \left(-\frac{\pi}{R} + \frac{2}{R} \sum_{n \in \mathbb{Z}} L_n \frac{\sin \frac{\pi n t}{R}}{n} \right)} \right\rangle_{\beta - \frac{i\lambda}{2}} \left\langle e^{i\lambda \frac{\pi L_0}{R}} \right\rangle_{\beta}$$

The full counting statistics in CFT

[DB & BD]

$$F(\lambda) = \frac{i\lambda\pi c}{12} \left(\frac{1}{\beta_r(\beta_r - i\lambda)} - \frac{1}{\beta_l(\beta_l + i\lambda)} \right)$$

Using dimensional analysis, unique solution to:

- Factorization $F(\lambda) = f(\lambda, \beta_l) + f(-\lambda, \beta_r)$
- Leading behaviour $F(\lambda) = O(\lambda)$
- Fluctuation relation

A stochastic interpretation

Independent Poisson processes for jumps of every energy E , positive or negative, with intensity

$$dE e^{-\beta_l E} \quad (E > 0)$$

$$dE e^{\beta_r E} \quad (E < 0)$$

Conclusion and perspectives

- We have proof of FR, hence of results, up to addressing cancellation of UV divergences...
- Generalization to presence of conformal impurity, to integrable models (thermodynamic Bethe ansatz), to charge currents, etc.
- Higher dimensions: same formula?
- Stochastic re-interpretation of CFT? of QFT?