

# REDUCTION OF ONE-LOOP AMPLITUDES AT THE INTEGRAND LEVEL

Costas G. Papadopoulos

NCSR “Demokritos”, Athens



RADCOR 2007, FIRENZE, 1-5 October 2007

## 1 INTRODUCTION: WISHLISTS AND TROUBLES

## 2 OPP REDUCTION

## 3 NUMERICAL TESTS

- 4-photon amplitudes
- 6-photon amplitudes
- ZZZ production

# INTRODUCTION: LHC NEEDS NLO

- The experimental programs of LHC require high precision predictions for multi-particle processes (also ILC of course)

# INTRODUCTION: LHC NEEDS NLO

- The experimental programs of LHC require high precision predictions for multi-particle processes (also ILC of course)
- In the last years we have seen a remarkable progress in the theoretical description of multi-particle processes at tree-order, thanks to very efficient recursive algorithms

# INTRODUCTION: LHC NEEDS NLO

- The experimental programs of LHC require high precision predictions for multi-particle processes (also ILC of course)
- In the last years we have seen a remarkable progress in the theoretical description of multi-particle processes at tree-order, thanks to very efficient recursive algorithms
- The current need of precision goes beyond tree order. At LHC, most analyses require at least next-to-leading order calculations (NLO)

# INTRODUCTION: LHC NEEDS NLO

- The experimental programs of LHC require high precision predictions for multi-particle processes (also ILC of course)
- In the last years we have seen a remarkable progress in the theoretical description of multi-particle processes at tree-order, thanks to very efficient recursive algorithms
- The current need of precision goes beyond tree order. At LHC, most analyses require at least next-to-leading order calculations (NLO)
- As a result, a big effort has been devoted by several groups to the problem of an efficient computation of one-loop corrections for multi-particle processes!

[from G. Heinrich's Summary talk]

## Wishlist Les Houches 2007

1.  $pp \rightarrow V V + \text{jet}$
2.  $pp \rightarrow t\bar{t} b\bar{b}$
3.  $pp \rightarrow t\bar{t} + 2 \text{ jets}$
4.  $pp \rightarrow W W W$
5.  $pp \rightarrow V V b\bar{b}$
6.  $pp \rightarrow V V + 2 \text{ jets}$
7.  $pp \rightarrow V + 3 \text{ jets}$
8.  $pp \rightarrow t\bar{t} b\bar{b}$
9.  $pp \rightarrow 4 \text{ jets}$

Processes for which a NLO calculation is both desired and feasible

Will we “finish” in time for LHC?

# WHAT HAS BEEN DONE? (2005-2007)

Some recent results → Cross Sections available

- $pp \rightarrow ZZ Z$   $pp \rightarrow t\bar{t}Z$  [Lazopoulos, Melnikov, Petriello]
- $pp \rightarrow H + 2$  jets [Campbell, et al., J. R. Andersen, et al.]
- $pp \rightarrow VV + 2$  jets via VBF [Bozzi, Jäger, Oleari, Zeppenfeld]

Mostly  $2 \rightarrow 3$ , very few  $2 \rightarrow 4$  complete calculations.

- $e^+ e^- \rightarrow 4$  fermions [Denner, Dittmaier, Roth]
- $e^+ e^- \rightarrow HH\nu\bar{\nu}$  [GRACE group (Boudjema et al.)]

This is NOT a complete list

(A lot of work has been done at NLO → calculations & new methods)



## Problems arising in NLO calculations

- Large **Number of Feynman diagrams**
- **Reduction to Scalar Integrals** (or sets of known integrals)
- **Numerical Instabilities** (inverse Gram determinants, spurious phase-space singularities)
- Extraction of **soft and collinear singularities** (we need virtual and real corrections)

- **Traditional** Method: Feynman Diagrams & Passarino-Veltman Reduction:
  - general applicability major achievements
  - but major problem: not designed @ amplitude level

- **Traditional** Method: Feynman Diagrams & Passarino-Veltman Reduction:
- **Semi-Numerical** Approach (Algebraic/Partly Numerical – Improved traditional) → Reduction to set of well-known integrals
- **Numerical** Approach (Numerical/Partly Algebraic) → Compute tensor integrals numerically
  - Ellis, Giele, Glover, Zanderighi;
  - Binoth, Guillet, Heinrich, Schubert;
  - Denner, Dittmaier; Del Aguila, Pittau;
  - Ferroglia, Passera, Passarino, Uccirati;
  - Nagy, Soper; van Hameren, Vollinga, Weinzierl;

- **Traditional** Method: Feynman Diagrams & Passarino-Veltman Reduction:
- **Semi-Numerical** Approach (Algebraic/Partly Numerical – Improved traditional) → Reduction to set of well-known integrals
- **Numerical** Approach (Numerical/Partly Algebraic) → Compute tensor integrals numerically
- **Analytic** Approach (Twistor-inspired)
  - extract information from lower-loop, lower-point amplitudes
  - determine scattering amplitudes by their poles and cuts
    - **major advantage: designed to work @ amplitude level**
    - **quadruple and triple cuts major simplifications**
    - Bern, Dixon, Dunbar, Kosower, Berger, Forde;
    - Anastasiou, Britto, Cachazo, Feng, Kunszt, Mastrolia;

- **Traditional** Method: Feynman Diagrams & Passarino-Veltman Reduction:
- **Semi-Numerical** Approach (Algebraic/Partly Numerical – Improved traditional) → Reduction to set of well-known integrals
- **Numerical** Approach (Numerical/Partly Algebraic) → Compute tensor integrals numerically
- **Analytic** Approach (Twistor-inspired)
  - extract information from lower-loop, lower-point amplitudes
  - determine scattering amplitudes by their poles and cuts
- ★ **OPP Integrand-level reduction combine: PV@integrand + n-particle cuts**

# OPP REDUCTION - INTRO

G. Ossola., C. G. Papadopoulos and R. Pittau,

Nucl. Phys. B **763**, 147 (2007) – arXiv:hep-ph/0609007

and JHEP **0707** (2007) 085 – arXiv:0704.1271 [hep-ph]

Any  $m$ -point one-loop amplitude can be written, **before integration**, as

$$A(\bar{q}) = \frac{N(q)}{\bar{D}_0 \bar{D}_1 \cdots \bar{D}_{m-1}}$$

A bar denotes objects living in  $n = 4 + \epsilon$  dimensions

$$\bar{D}_i = (\bar{q} + p_i)^2 - m_i^2$$

$$\bar{q}^2 = q^2 + \tilde{q}^2$$

$$\bar{D}_i = D_i + \tilde{q}^2$$

External momenta  $p_i$  are 4-dimensional objects

# THE OLD “MASTER” FORMULA

$$\begin{aligned}\int A &= \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} d(i_0 i_1 i_2 i_3) D_0(i_0 i_1 i_2 i_3) \\ &+ \sum_{i_0 < i_1 < i_2}^{m-1} c(i_0 i_1 i_2) C_0(i_0 i_1 i_2) \\ &+ \sum_{i_0 < i_1}^{m-1} b(i_0 i_1) B_0(i_0 i_1) \\ &+ \sum_{i_0}^{m-1} a(i_0) A_0(i_0) \\ &+ \text{rational terms}\end{aligned}$$

General expression for the 4-dim  $N(q)$  at the integrand level in terms of  $D_i$

$$\begin{aligned}
 N(q) &= \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i \\
 &+ \sum_{i_0 < i_1 < i_2}^{m-1} \left[ c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2) \right] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\
 &+ \sum_{i_0 < i_1}^{m-1} \left[ b(i_0 i_1) + \tilde{b}(q; i_0 i_1) \right] \prod_{i \neq i_0, i_1}^{m-1} D_i \\
 &+ \sum_{i_0}^{m-1} \left[ a(i_0) + \tilde{a}(q; i_0) \right] \prod_{i \neq i_0}^{m-1} D_i
 \end{aligned}$$



# OPP “MASTER” FORMULA - II

$$\begin{aligned}
 N(q) = & \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[ d(i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i + \sum_{i_0 < i_1 < i_2}^{m-1} \left[ c(i_0 i_1 i_2) \right] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\
 & + \sum_{i_0 < i_1}^{m-1} \left[ b(i_0 i_1) \right] \prod_{i \neq i_0, i_1}^{m-1} D_i + \sum_{i_0}^{m-1} \left[ a(i_0) \right] \prod_{i \neq i_0}^{m-1} D_i
 \end{aligned}$$

- The quantities  $d(i_0 i_1 i_2 i_3)$  are the coefficients of 4-point functions with denominators labeled by  $i_0$ ,  $i_1$ ,  $i_2$ , and  $i_3$ .
- $c(i_0 i_1 i_2)$ ,  $b(i_0 i_1)$ ,  $a(i_0)$  are the coefficients of all possible 3-point, 2-point and 1-point functions, respectively.

# OPP “MASTER” FORMULA - II

$$\begin{aligned} N(q) &= \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i + \sum_{i_0 < i_1 < i_2}^{m-1} \left[ c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2) \right] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\ &+ \sum_{i_0 < i_1}^{m-1} \left[ b(i_0 i_1) + \tilde{b}(q; i_0 i_1) \right] \prod_{i \neq i_0, i_1}^{m-1} D_i + \sum_{i_0}^{m-1} \left[ a(i_0) + \tilde{a}(q; i_0) \right] \prod_{i \neq i_0}^{m-1} D_i \end{aligned}$$

The quantities  $\tilde{d}$ ,  $\tilde{c}$ ,  $\tilde{b}$ ,  $\tilde{a}$  are the “spurious” terms

- They still depend on  $q$  (integration momentum)
- They should vanish upon integration

What is the explicit expression of the spurious term?

Following F. del Aguila and R. Pittau, arXiv:hep-ph/0404120

- Express any  $q$  in  $N(q)$  as

$$q^\mu = -p_0^\mu + \sum_{i=1}^4 G_i l_i^\mu, \quad l_i^2 = 0$$

$$k_1 = l_1 + \alpha_1 l_2, \quad k_2 = l_2 + \alpha_2 l_1, \quad k_i = p_i - p_0 \\ l_3^\mu = \langle l_1 | \gamma^\mu | l_2 \rangle, \quad l_4^\mu = \langle l_2 | \gamma^\mu | l_1 \rangle$$

- The coefficients  $G_i$  either reconstruct denominators  $D_i$

→ They give rise to  $d$ ,  $c$ ,  $b$ ,  $a$  coefficients

Following F. del Aguila and R. Pittau, arXiv:hep-ph/0404120

- Express any  $q$  in  $N(q)$  as

$$q^\mu = -p_0^\mu + \sum_{i=1}^4 G_i l_i^\mu, \quad l_i^2 = 0$$

$$k_1 = l_1 + \alpha_1 l_2, \quad k_2 = l_2 + \alpha_2 l_1, \quad k_i = p_i - p_0 \\ l_3^\mu = \langle l_1 | \gamma^\mu | l_2 \rangle, \quad l_4^\mu = \langle l_2 | \gamma^\mu | l_1 \rangle$$

- The coefficients  $G_i$  either reconstruct denominators  $D_i$  or vanish upon integration
  - They give rise to  $d, c, b, a$  coefficients
  - They form the spurious  $\tilde{d}, \tilde{c}, \tilde{b}, \tilde{a}$  coefficients

# SPURIOUS TERMS - II

- $\tilde{d}(q)$  term (only 1)

$$\tilde{d}(q) = \tilde{d} T(q),$$

where  $\tilde{d}$  is a constant (does not depend on  $q$ )

$$T(q) \equiv \text{Tr}[(\not{q} + \not{p}_0)\not{1}\not{2}\not{3}\not{5}]$$

- $\tilde{d}(q)$  term (only 1)

$$\tilde{d}(q) = \tilde{d} T(q),$$

where  $\tilde{d}$  is a constant (does not depend on  $q$ )

$$T(q) \equiv \text{Tr}[(\not{q} + \not{p}_0)\not{\ell}_1\not{\ell}_2\not{k}_3\gamma_5]$$

- $\tilde{c}(q)$  terms (they are 6)

$$\tilde{c}(q) = \sum_{j=1}^{j_{\max}} \{ \tilde{c}_{1j} [(q + p_0) \cdot \ell_3]^j + \tilde{c}_{2j} [(q + p_0) \cdot \ell_4]^j \}$$

In the renormalizable gauge,  $j_{\max} = 3$

- $\tilde{d}(q)$  term (only 1)

$$\tilde{d}(q) = \tilde{d} T(q),$$

where  $\tilde{d}$  is a constant (does not depend on  $q$ )

$$T(q) \equiv Tr[(\not{q} + \not{p}_0)\not{\ell}_1\not{\ell}_2\not{k}_3\gamma_5]$$

- $\tilde{c}(q)$  terms (they are 6)

$$\tilde{c}(q) = \sum_{j=1}^{j_{max}} \{ \tilde{c}_{1j} [(q + p_0) \cdot \ell_3]^j + \tilde{c}_{2j} [(q + p_0) \cdot \ell_4]^j \}$$

In the renormalizable gauge,  $j_{max} = 3$

- $\tilde{b}(q)$  and  $\tilde{a}(q)$  give rise to 8 and 4 terms, respectively

# A SIMPLE EXAMPLE

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4}$$



# A SIMPLE EXAMPLE

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4}$$
$$1 = \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$

# A SIMPLE EXAMPLE

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4}$$
$$1 = \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$
$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$

# A SIMPLE EXAMPLE

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4}$$

$$1 = \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} = \sum d(i_0 i_1 i_2 i_3) D_0(i_0 i_1 i_2 i_3)$$

# A SIMPLE EXAMPLE

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4}$$

$$1 = \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} = \sum d(i_0 i_1 i_2 i_3) D_0(i_0 i_1 i_2 i_3)$$

$$d(i_0 i_1 i_2 i_3) = \frac{1}{2} \left( \frac{1}{D_{i_4}(q^+)} + \frac{1}{D_{i_4}(q^-)} \right)$$

# A SIMPLE EXAMPLE

$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4}$$
$$1 = \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$
$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} \sum \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] D_{i_4}$$
$$\int \frac{1}{D_0 D_1 D_2 D_3 D_4} = \sum d(i_0 i_1 i_2 i_3) D_0(i_0 i_1 i_2 i_3)$$
$$d(i_0 i_1 i_2 i_3) = \frac{1}{2} \left( \frac{1}{D_{i_4}(q^+)} + \frac{1}{D_{i_4}(q^-)} \right)$$

- Melrose, Nuovo Cim. 40 (1965) 181
- G. Källén, J. Toll, J. Math. Phys. 6, 299 (1965)

Now we know the form of the spurious terms:

$$\begin{aligned}
 N(q) &= \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i + \sum_{i_0 < i_1 < i_2}^{m-1} \left[ c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2) \right] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\
 &+ \sum_{i_0 < i_1}^{m-1} \left[ b(i_0 i_1) + \tilde{b}(q; i_0 i_1) \right] \prod_{i \neq i_0, i_1}^{m-1} D_i + \sum_{i_0}^{m-1} \left[ a(i_0) + \tilde{a}(q; i_0) \right] \prod_{i \neq i_0}^{m-1} D_i
 \end{aligned}$$

Our calculation is now reduced to an algebraic problem

Now we know the form of the spurious terms:

$$\begin{aligned}
 N(q) = & \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i + \sum_{i_0 < i_1 < i_2}^{m-1} \left[ c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2) \right] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\
 & + \sum_{i_0 < i_1}^{m-1} \left[ b(i_0 i_1) + \tilde{b}(q; i_0 i_1) \right] \prod_{i \neq i_0, i_1}^{m-1} D_i + \sum_{i_0}^{m-1} \left[ a(i_0) + \tilde{a}(q; i_0) \right] \prod_{i \neq i_0}^{m-1} D_i
 \end{aligned}$$

Our calculation is now reduced to an **algebraic problem**

Extract all the coefficients by evaluating  $N(q)$  for a set of values of the integration momentum  $q$

Now we know the form of the spurious terms:

$$\begin{aligned}
 N(q) = & \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \left[ d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3) \right] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i + \sum_{i_0 < i_1 < i_2}^{m-1} \left[ c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2) \right] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i \\
 & + \sum_{i_0 < i_1}^{m-1} \left[ b(i_0 i_1) + \tilde{b}(q; i_0 i_1) \right] \prod_{i \neq i_0, i_1}^{m-1} D_i + \sum_{i_0}^{m-1} \left[ a(i_0) + \tilde{a}(q; i_0) \right] \prod_{i \neq i_0}^{m-1} D_i
 \end{aligned}$$

Our calculation is now reduced to an **algebraic problem**

Extract all the coefficients by evaluating  $N(q)$  for a set of values of the integration momentum  $q$

There is a very good set of such points: **Use values of  $q$  for which a set of denominators  $D_i$  vanish**  $\rightarrow$  The system becomes “triangular”: solve first for 4-point functions, then 3-point functions and so on



## EXAMPLE: 4-PARTICLES PROCESS

$$\begin{aligned} N(q) &= d + \tilde{d}(q) + \sum_{i=0}^3 [c(i) + \tilde{c}(q; i)] D_i + \sum_{i_0 < i_1}^3 [b(i_0 i_1) + \tilde{b}(q; i_0 i_1)] D_{i_0} D_{i_1} \\ &+ \sum_{i_0=0}^3 [a(i_0) + \tilde{a}(q; i_0)] D_{i \neq i_0} D_{j \neq i_0} D_{k \neq i_0} \end{aligned}$$

We look for a  $q$  of the form  $q^\mu = -p_0^\mu + x_i \ell_i^\mu$  such that

$$D_0 = D_1 = D_2 = D_3 = 0$$

→ we get a system of equations in  $x_i$  that has **two solutions**  $q_0^\pm$

## EXAMPLE: 4-PARTICLES PROCESS

$$N(q) = d + \tilde{d}(q)$$

Our “master formula” for  $q = q_0^\pm$  is:

$$N(q_0^\pm) = [d + \tilde{d} T(q_0^\pm)]$$

→ solve to extract the coefficients  $d$  and  $\tilde{d}$

## EXAMPLE: 4-PARTICLES PROCESS

$$\begin{aligned} N(q) - d - \tilde{d}(q) &= \sum_{i=0}^3 [c(i) + \tilde{c}(q; i)] D_i + \sum_{i_0 < i_1}^3 [b(i_0 i_1) + \tilde{b}(q; i_0 i_1)] D_{i_0} D_{i_1} \\ &+ \sum_{i_0=0}^3 [a(i_0) + \tilde{a}(q; i_0)] D_{i \neq i_0} D_{j \neq i_0} D_{k \neq i_0} \end{aligned}$$

Then we can move to the extraction of **c coefficients** using

$$N'(q) = N(q) - d - \tilde{d}T(q)$$

and setting to zero three denominators (ex:  $D_1 = 0, D_2 = 0, D_3 = 0$ )

## EXAMPLE: 4-PARTICLES PROCESS

$$N(q) - d - \tilde{d}(q) = [c(0) + \tilde{c}(q; 0)] D_0$$

We have infinite values of  $q$  for which

$$D_1 = D_2 = D_3 = 0 \quad \text{and} \quad D_0 \neq 0$$

→ Here we need 7 of them to determine  $c(0)$  and  $\tilde{c}(q; 0)$

- Let's go back to the integrand

$$A(\bar{q}) = \frac{N(q)}{\bar{D}_0 \bar{D}_1 \cdots \bar{D}_{m-1}}$$

- Insert the expression for  $N(q) \rightarrow$  we know all the coefficients

$$N(q) = \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} [d + \tilde{d}(q)] \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} D_i + \sum_{i_0 < i_1 < i_2}^{m-1} [c + \tilde{c}(q)] \prod_{i \neq i_0, i_1, i_2}^{m-1} D_i + \cdots$$

- Finally rewrite all denominators using

$$\frac{D_i}{\bar{D}_i} = \bar{z}_i, \quad \text{with} \quad \bar{z}_i \equiv \left(1 - \frac{\tilde{q}^2}{\bar{D}_i}\right)$$

# RATIONAL TERMS - II

$$\begin{aligned}
 A(\bar{q}) &= \sum_{i_0 < i_1 < i_2 < i_3}^{m-1} \frac{d(i_0 i_1 i_2 i_3) + \tilde{d}(q; i_0 i_1 i_2 i_3)}{\bar{D}_{i_0} \bar{D}_{i_1} \bar{D}_{i_2} \bar{D}_{i_3}} \prod_{i \neq i_0, i_1, i_2, i_3}^{m-1} \bar{Z}_i \\
 &+ \sum_{i_0 < i_1 < i_2}^{m-1} \frac{c(i_0 i_1 i_2) + \tilde{c}(q; i_0 i_1 i_2)}{\bar{D}_{i_0} \bar{D}_{i_1} \bar{D}_{i_2}} \prod_{i \neq i_0, i_1, i_2}^{m-1} \bar{Z}_i \\
 &+ \sum_{i_0 < i_1}^{m-1} \frac{b(i_0 i_1) + \tilde{b}(q; i_0 i_1)}{\bar{D}_{i_0} \bar{D}_{i_1}} \prod_{i \neq i_0, i_1}^{m-1} \bar{Z}_i \\
 &+ \sum_{i_0}^{m-1} \frac{a(i_0) + \tilde{a}(q; i_0)}{\bar{D}_{i_0}} \prod_{i \neq i_0}^{m-1} \bar{Z}_i
 \end{aligned}$$

The rational part is produced, after integrating over  $d^n q$ , by the  $\tilde{q}^2$  dependence in  $\bar{Z}_i$

$$\bar{Z}_i \equiv \left( 1 - \frac{\tilde{q}^2}{\bar{D}_i} \right)$$

The “Extra Integrals” are of the form

$$I_{s;\mu_1\cdots\mu_r}^{(n;2\ell)} \equiv \int d^n q \tilde{q}^{2\ell} \frac{q_{\mu_1} \cdots q_{\mu_r}}{\bar{D}(k_0) \cdots \bar{D}(k_s)},$$

where

$$\bar{D}(k_i) \equiv (\bar{q} + k_i)^2 - m_i^2, k_i = p_i - p_0$$

**These integrals:**

- have dimensionality  $\mathcal{D} = 2(1 + \ell - s) + r$
- contribute only when  $\mathcal{D} \geq 0$ , otherwise are of  $\mathcal{O}(\epsilon)$

Calculate  $N(q)$



## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly

## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly
- Calculate  $N(q)$  numerically via recursion relations

## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly
- Calculate  $N(q)$  numerically via recursion relations
- Just specify external momenta, polarization vectors and masses and proceed with the reduction!

## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly
- Calculate  $N(q)$  numerically via recursion relations
- Just specify external momenta, polarization vectors and masses and proceed with the reduction!

## Compute all coefficients

## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly
- Calculate  $N(q)$  numerically via recursion relations
- Just specify external momenta, polarization vectors and masses and proceed with the reduction!

## Compute all coefficients

- by evaluating  $N(q)$  at certain values of integration momentum

## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly
- Calculate  $N(q)$  numerically via recursion relations
- Just specify external momenta, polarization vectors and masses and proceed with the reduction!

## Compute all coefficients

- by evaluating  $N(q)$  at certain values of integration momentum

## Evaluate scalar integrals

## Calculate $N(q)$

- We do not need to repeat this for all Feynman diagrams. We can group them and solve for (sub)amplitudes directly
- Calculate  $N(q)$  numerically via recursion relations
- Just specify external momenta, polarization vectors and masses and proceed with the reduction!

## Compute all coefficients

- by evaluating  $N(q)$  at certain values of integration momentum

## Evaluate scalar integrals

- massive integrals  $\rightarrow$  FF [G. J. van Oldenborgh]
- massless integrals  $\rightarrow$  OneLOop [A. van Hameren]

- PV:
  
  
  
  
  
  
  
  
  
  
- Unitarity Methods:



- PV:
  - $N(q)$  or  $A(q)$  hasn't to be known analytically
  
- Unitarity Methods:

- PV:
  - $N(q)$  or  $A(q)$  hasn't to be known analytically
  - No computer algebra
  
- Unitarity Methods:

- PV:
  - $N(q)$  or  $A(q)$  hasn't to be known analytically
  - No computer algebra
  - Mathematica  $\rightarrow$  Numerica
- Unitarity Methods:

- PV:
  - $N(q)$  or  $A(q)$  hasn't to be known analytically
  - No computer algebra
  - Mathematica  $\rightarrow$  Numerica
- Unitarity Methods:
  - A more transparent algebraic method

- PV:
  - $N(q)$  or  $A(q)$  hasn't to be known analytically
  - No computer algebra
  - Mathematica  $\rightarrow$  Numerica
- Unitarity Methods:
  - A more transparent algebraic method
  - A solid way to get all rational terms

# THE MASTER EQUATION

## Properties of the master equation

## Properties of the master equation

- Polynomial equation in  $q$

## Properties of the master equation

- Polynomial equation in  $q$
- Highly redundant: the a-terms have a degree of  $m^2 - 2$  compared to  $m$  as a function of  $q$



## Properties of the master equation

- Polynomial equation in  $q$
- Highly redundant: the a-terms have a degree of  $m^2 - 2$  compared to  $m$  as a function of  $q$
- Zeros of (a tower of ) polynomial equations

## Properties of the master equation

- Polynomial equation in  $q$
- Highly redundant: the a-terms have a degree of  $m^2 - 2$  compared to  $m$  as a function of  $q$
- Zeros of (a tower of ) polynomial equations
- Different ways of solving it, besides 'unitarity method'

## Properties of the master equation


- Polynomial equation in  $q$
- Highly redundant: the a-terms have a degree of  $m^2 - 2$  compared to  $m$  as a function of  $q$
- Zeros of (a tower of ) polynomial equations
- Different ways of solving it, besides 'unitarity method'

## The $N \equiv N$ test

A tool to efficiently treat phase-space points with numerical instabilities

# 4-PHOTON AND 6-PHOTON AMPLITUDES

As an example we present 4-photon and 6-photon amplitudes  
(via fermionic loop of mass  $m_f$ )




Input parameters for the reduction:

- External momenta  $p_i$
- Masses of propagators in the loop
- Polarization vectors

# 4-PHOTON AND 6-PHOTON AMPLITUDES

As an example we present **4-photon and 6-photon amplitudes**  
(via fermionic loop of mass  $m_f$ )



Input parameters for the reduction:

- External momenta  $p_i \rightarrow$  in this example **massless**, i.e.  $p_i^2 = 0$
- Masses of propagators in the loop  $\rightarrow$  **all equal to  $m_f$**
- Polarization vectors  $\rightarrow$  various helicity configurations

$$\frac{F_{++++}^f}{\alpha^2 Q_f^4} = -8$$

Rational Part

$$\begin{aligned}
 \frac{F_{++++}^f}{\alpha^2 Q_f^4} = & -8 + 8 \left( 1 + \frac{2\hat{u}}{\hat{s}} \right) B_0(\hat{u}) + 8 \left( 1 + \frac{2\hat{t}}{\hat{s}} \right) B_0(\hat{t}) \\
 & - 8 \left( \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} \right) [\hat{t}C_0(\hat{t}) + \hat{u}C_0(\hat{u})] \\
 & - 4 \left[ \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} \right] D_0(\hat{t}, \hat{u})
 \end{aligned}$$

Massless four-photon amplitudes

$$\begin{aligned}
 \frac{F_{++++}^f}{\alpha^2 Q_f^4} &= -8 + 8 \left(1 + \frac{2\hat{u}}{\hat{s}}\right) B_0(\hat{u}) + 8 \left(1 + \frac{2\hat{t}}{\hat{s}}\right) B_0(\hat{t}) \\
 &\quad - 8 \left(\frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} - \frac{4m_f^2}{\hat{s}}\right) [\hat{t}C_0(\hat{t}) + \hat{u}C_0(\hat{u})] \\
 &\quad - 4 \left[4m_f^4 - (2\hat{s}m_f^2 + \hat{t}\hat{u})\frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} + \frac{4m_f^2\hat{t}\hat{u}}{\hat{s}}\right] D_0(\hat{t}, \hat{u}) \\
 &\quad + 8m_f^2(\hat{s} - 2m_f^2)[D_0(\hat{s}, \hat{t}) + D_0(\hat{s}, \hat{u})]
 \end{aligned}$$

Massive four-photon amplitudes




$$\begin{aligned}
 \frac{F_{++++}^f}{\alpha^2 Q_f^4} &= -8 + 8 \left( 1 + \frac{2\hat{u}}{\hat{s}} \right) B_0(\hat{u}) + 8 \left( 1 + \frac{2\hat{t}}{\hat{s}} \right) B_0(\hat{t}) \\
 &\quad - 8 \left( \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} - \frac{4m_f^2}{\hat{s}} \right) [\hat{t}C_0(\hat{t}) + \hat{u}C_0(\hat{u})] \\
 &\quad - 4 \left[ 4m_f^4 - (2\hat{s}m_f^2 + \hat{t}\hat{u}) \frac{\hat{t}^2 + \hat{u}^2}{\hat{s}^2} + \frac{4m_f^2\hat{t}\hat{u}}{\hat{s}} \right] D_0(\hat{t}, \hat{u}) \\
 &\quad + 8m_f^2(\hat{s} - 2m_f^2)[D_0(\hat{s}, \hat{t}) + D_0(\hat{s}, \hat{u})]
 \end{aligned}$$


Massive four-photon amplitudes

Results also checked for  $F_{+++-}^f$  and  $F_{+--+}^f$



Massless case:  $[++-- --]$  and  $[+- - + +-]$




Plot presented by Nagy and Soper hep-ph/0610028  
(also Binoth et al., hep-ph/0703311)

Massless case:  $[++--]$  and  $[+-+ -]$ 

Analogous plot produced with OPP reduction


Massless case:  $[++-- --]$  and  $[++-- +-]$ 

 Same plot as before for a wider range of  $\theta$

Massless case:  $[++--]$  and  $[++--+-]$




Same idea for a different set of external momenta

# SIX PHOTONS WITH MASSIVE FERMIONS




- Massless result [Mahlon]

# SIX PHOTONS WITH MASSIVE FERMIONS



- Massless result [Mahlon]
- $m = 0.5 \text{ GeV}$


# SIX PHOTONS WITH MASSIVE FERMIONS



- Massless result [Mahlon]
- $m = 0.5$  GeV
- $m = 4.5$  GeV




# SIX PHOTONS WITH MASSIVE FERMIONS



- Massless result [Mahlon]
- $m = 0.5$  GeV
- $m = 4.5$  GeV
- $m = 12.0$  GeV


# SIX PHOTONS WITH MASSIVE FERMIONS



- Massless result [Mahlon]
- $m = 0.5$  GeV
- $m = 4.5$  GeV
- $m = 12.0$  GeV
- $m = 20.0$  GeV

# $q\bar{q} \rightarrow ZZZ$ VIRTUAL CORRECTIONS

A. Lazopoulos, K. Melnikov and F. Petriello, [arXiv:hep-ph/0703273]



POLES  $1/\epsilon^2$  AND  $1/\epsilon$

$$\sigma^{\text{NLO, virt}}|_{\text{div}} = -C_F \frac{\alpha_s}{\pi} \frac{\Gamma(1+\epsilon)}{(4\pi)^{-\epsilon}} (s_{12})^{-\epsilon} \left( \frac{1}{\epsilon^2} + \frac{3}{2\epsilon} \right) \sigma^{\text{LO}}$$

# $q\bar{q} \rightarrow ZZZ$ VIRTUAL CORRECTIONS

A very naive implementation

## A very naive implementation

- Calculate the  $N(q)$  by brute (numerical) force namely multiplying gamma matrices !

## A very naive implementation

- Calculate the  $N(q)$  by brute (numerical) force namely multiplying gamma matrices !
- Calculate 4d and rational terms graph by graph

## A very naive implementation

- Calculate the  $N(q)$  by brute (numerical) force namely multiplying gamma matrices !
- Calculate 4d and rational terms graph by graph

## Comparison with LMP

## A very naive implementation

- Calculate the  $N(q)$  by brute (numerical) force namely multiplying gamma matrices !
- Calculate 4d and rational terms graph by graph

## Comparison with LMP

- Of course full agreement for the  $1/\epsilon^2$  and  $1/\epsilon$  terms



## A very naive implementation

- Calculate the  $N(q)$  by brute (numerical) force namely multiplying gamma matrices !
- Calculate 4d and rational terms graph by graph

## Comparison with LMP

- Of course full agreement for the  $1/\epsilon^2$  and  $1/\epsilon$  terms
- An 'easy' agreement for all graphs with up to 4-point loop integrals



## A very naive implementation

- Calculate the  $N(q)$  by brute (numerical) force namely multiplying gamma matrices !
- Calculate 4d and rational terms graph by graph



## Comparison with LMP

- Of course full agreement for the  $1/\epsilon^2$  and  $1/\epsilon$  terms
- An 'easy' agreement for all graphs with up to 4-point loop integrals
- A bit more work to uncover the differences in scalar function normalization that happen to show to order  $\epsilon^2$  thus influence only 5-point loop integrals.

# $q\bar{q} \rightarrow ZZZ$ VIRTUAL CORRECTIONS



# $q\bar{q} \rightarrow ZZZ$ VIRTUAL CORRECTIONS



# $q\bar{q} \rightarrow ZZZ$ VIRTUAL CORRECTIONS

Typical precision:

Typical precision:

- LMP: 9.573(66) about 1% error

## Typical precision:

- LMP: 9.573(66) about 1% error
- OPP:
  - $\left\{ \begin{array}{l} -26.45706742815552 \\ -26.457067428165503661018557937723426 \end{array} \right.$

## Typical precision:

- LMP: 9.573(66) about 1% error
- OPP:
  - $\left\{ \begin{array}{l} -26.45706742815552 \\ -26.457067428165503661018557937723426 \end{array} \right.$

Typical time: 200 times faster (for non-singular PS-points)



## Typical precision:

- LMP: 9.573(66) about 1% error
- OPP:  
$$\begin{cases} -26.45706742815552 \\ -26.457067428165503661018557937723426 \end{cases}$$

Typical time: 200 times faster (for non-singular PS-points)

Future perspectives: up to 4 orders of magnitude

Reduction at the integrand level

## Reduction at the integrand level

- changes the computational approach at one loop

## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Future

## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Future

- Understand potential stability problems

## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Future

- Understand potential stability problems
- Combine with the real corrections

## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Future

- Understand potential stability problems
- Combine with the real corrections
- Automate through Dyson-Schwinger equations



## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Future

- Understand potential stability problems
- Combine with the real corrections
- Automate through Dyson-Schwinger equations

A generic NLO calculator seems feasible

## Reduction at the integrand level

- changes the computational approach at one loop
- Numerical but still algebraic: speed and precision not a problem

## Future

- Understand potential stability problems
- Combine with the real corrections
- Automate through Dyson-Schwinger equations


A generic NLO calculator seems feasible

CUTTOOLS version 0. is ready !

# The Outlook

## ■ The DS equations to all orders


- Imagine a theory with 3- and 4- point vertices and just one field. Then it is straightforward to write an equation that gives the amplitude for  $1 \rightarrow n$



# The Outlook

## ■ The DS equations to all orders


- Imagine a theory with 3- and 4- point vertices and just one field. Then it is straightforward to write an equation that gives the amplitude for  $1 \rightarrow n$



# The Outlook


## ■ The DS equations to all orders

- Imagine a theory with 3- and 4- point vertices and just one field. Then it is straightforward to write an equation that gives the amplitude for  $1 \rightarrow n$



# The Outlook


- The DS equations to tree order



- Combine 2,3,...,n external particles


# The Outlook

- The DS equations to one loop order **linear** !



# The Outlook

- The DS equations to one loop order



- N+1 tree order sub-amplitudes