


Heavy Flavour Production in DIS

Two–Loop Massive Operator Matrix Elements and Beyond

Johannes Blümlein, DESY

in collaboration with I. Bierenbaum and S. Klein



1. Introduction
2. The Method
3. The Calculation
4. Results
5. Comparison to Former Results
6. Conclusion

based on:


J.B., A. De Freitas, W.L. van Neerven, and S. Klein,
Nucl. Phys. **B755** (2006) 272.

I. Bierenbaum, J.B., and S. Klein, Phys. Lett. **B648**
(2007) 195; Nucl. Phys. **B780** (2007) 40

and in preparation.;

I. Bierenbaum, J.B., S. Klein, and C. Schneider,
arXiv:0707.4759 [math-ph].

Deep-Inelastic Scattering (DIS):




$$\rightarrow L^{\mu\nu}$$

$$Q^2 := -q^2, \quad x := \frac{Q^2}{2pq} \quad \text{Björken-x}$$

$$\rightarrow W_{\mu\nu}$$

$$\frac{d\sigma}{dQ^2 dx} \sim W_{\mu\nu} L^{\mu\nu}$$

Heavy-flavor production: LO-process: photon-gluon fusion





Hadronic Tensor for heavy quark production via single photon exchange:

$$\begin{aligned}
 W_{\mu\nu}^{Q\bar{Q}}(q, P, s) &= \frac{1}{4\pi} \int d^4\xi \exp(iq\xi) \langle P, s | [J_\mu^{em}(\xi), J_\nu^{em}(0)] | P, s \rangle_{Q\bar{Q}} \\
 &= \frac{1}{2x} \left(g_{\mu\nu} - \frac{q_\mu q_\nu}{q^2} \right) F_L^{Q\bar{Q}}(x, Q^2) + \frac{2x}{Q^2} \left(P_\mu P_\nu + \frac{q_\mu P_\nu + q_\nu P_\mu}{2x} - \frac{Q^2}{4x^2} g_{\mu\nu} \right) F_2^{Q\bar{Q}}(x, Q^2) \\
 &\quad - \frac{M}{2Pq} \varepsilon_{\mu\nu\alpha\beta} q^\alpha \left[s^\beta g_1^{Q\bar{Q}}(x, Q^2) + \left(s^\beta - \frac{sq}{Pq} p^\beta \right) g_2^{Q\bar{Q}}(x, Q^2) \right].
 \end{aligned}$$

Björken scaling, F_i depends only on x , Q^2 -independent
 scaling violation, F_i becomes Q^2 -dependent

Goal of heavy flavour improved calculation:

- Increase accuracy of perturbative description of structure functions
- More precise definition of the Gluon and Sea Quark Distributions
- QCD analysis and determination of Λ_{QCD} from DIS data



Unpolarized DIS :

- LO : [Witten, 1976; Babcock & Sivers, 1978; Shifman, Vainshtein, Zakharov 1978; Leveille & Weiler, 1979]
- NLO : [Laenen, Riemersma, Smith, van Neerven, 1993, 1995]
asymptotic : [Buza, Matiounine, Smith, Migneron, van Neerven, 1996]

Polarized DIS :

- LO : [Watson, 1982; Glück, Reya, Vogelsang, 1991; Vogelsang, 1991]
- NLO : asymptotic: [Buza, Matiounine, Smith, van Neerven, 1997]

Mellin–Space Expressions:

[Alekhin, Blümlein, 2003].

massless RGE and Light-Cone Expansion in Björken-Limit $\{Q^2, \nu\} \rightarrow \infty, x$ fixed:

$$\lim_{\xi^2 \rightarrow 0} [J(\xi), J(0)] \propto \sum_{i, N, \tau} c_{i, \tau}^N(\xi^2, \mu^2) \xi_{\mu_1} \dots \xi_{\mu_N} O_{i, \tau}^{\mu_1 \dots \mu_N}(0, \mu^2)$$

Operators: Flavour non-singlet, singlet and pure singlet; consider leading twist-2 operators

mass factorization between Wilson coefficients and parton densities;

$$F_i(x, Q^2) = \sum_j \underbrace{C_i^j\left(x, \frac{Q^2}{\mu^2}\right)}_{\text{perturbative}} \otimes \underbrace{f_j(x, \mu^2)}_{\text{non-perturbative}}$$

with $[f \otimes g](z) = \int_0^1 dz_1 \int_0^1 dz_2 \delta(z - z_1 z_2) f(z_1) g(z_2)$.

(massless) RGE: Altarelli-Parisi evolution equations for pdfs ($\mu^2 = Q^2$):

$$\frac{d}{d \ln Q^2} f_g(x, Q^2) = \sum_{l=0}^{\infty} a_s^{(l+1)}(Q^2) \int_x^1 \frac{dz}{z} \left\{ P_{g \leftarrow q}^{(l)}(z) \sum_f \left[f_f\left(\frac{x}{z}, Q^2\right) + f_{\bar{f}}\left(\frac{x}{z}, Q^2\right) \right] + P_{g \leftarrow g}^{(l)}(z) f_g\left(\frac{x}{z}, Q^2\right) \right\}$$

$P_{i \leftarrow j}^{(l)}(z)$ are the splitting functions.

Heavy quark contribution: heavy quark Wilson coefficient, $H_{(2,L),i}^{S,NS} \left(\frac{Q^2}{\mu^2}, \frac{m^2}{\mu^2} \right)$

The **Renormalization Group Equations**[†] imply factorization for all non-power terms:

$$H_{(2,L),i}^{S,NS} \left(\frac{Q^2}{\mu^2}, \frac{m^2}{\mu^2} \right) = \underbrace{A_{k,i}^{S,NS} \left(\frac{m^2}{\mu^2} \right)}_{\text{massive OMEs}} \otimes \underbrace{C_{(2,L),k}^{S,NS} \left(\frac{Q^2}{\mu^2} \right)}_{\text{light-Wilson coefficients}}.$$

holds for polarized and unpolarized case in limit $Q^2 \gg m_Q^2$, which means $Q^2/m_Q^2 \geq 10$ for $F_2(x, Q^2)$.

Here $\langle i|A_l|j \rangle$ denote the partonic operator matrix elements,

OMEs obey expansion

$$A_{k,i}^{S,NS} \left(\frac{m^2}{\mu^2} \right) = \langle i|O_k^{S,NS}|i \rangle = \delta_{k,i} + \sum_{l=1}^{\infty} a_s^l A_{k,i}^{S,NS,(l)} \left(\frac{m^2}{\mu^2} \right), \quad i = q, g$$

[[†] Buza, Matiounine, Migneron, Smith, van Neerven, 1996;

Buza, Matiounine, Smith, van Neerven, 1997.]

Expansion up to $O(\alpha_s^2)$ of unpolarized Heavy Flavor Wilson Coefficient H_2 :

$$\begin{aligned}
 H_{2,g}^S \left(\frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) &= a_s \left[A_{Qg}^{(1)} \left(\frac{m^2}{\mu^2} \right) + \widehat{C}_{2,g}^{(1)} \left(\frac{Q^2}{\mu^2} \right) \right] \\
 &+ a_s^2 \left[A_{Qg}^{(2)} \left(\frac{m^2}{\mu^2} \right) + A_{Qg}^{(1)} \left(\frac{m^2}{\mu^2} \right) \otimes C_{2,q}^{(1)} \left(\frac{Q^2}{\mu^2} \right) + \widehat{C}_{2,g}^{(2)} \left(\frac{Q^2}{\mu^2} \right) \right], \\
 H_{2,q}^{PS} \left(\frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) &= a_s^2 \left[A_{Qq}^{PS,(2)} \left(\frac{m^2}{\mu^2} \right) + \widehat{C}_{2,q}^{PS,(2)} \left(\frac{Q^2}{\mu^2} \right) \right], \\
 H_{2,q}^{NS} \left(\frac{Q^2}{m^2}, \frac{m^2}{\mu^2} \right) &= a_s^2 \left[A_{qq,Q}^{NS,(2)} \left(\frac{m^2}{\mu^2} \right) + \widehat{C}_{2,q}^{NS,(2)} \left(\frac{Q^2}{\mu^2} \right) \right].
 \end{aligned}$$

- Polarized and longitudinal **Heavy Wilson coefficients** obey similar expansion.
- For H_L , $O(a_s^3)$ contributions have been derived recently.
[J. Blümlein, A. De Freitas, W. van Neerven, S. Klein (2006)].


Gluonic Massive Operator Matrix Elements have the same structure in the polarized and unpolarized case. Up to $O(a_s^2)$ they are given by:

$$\begin{aligned}
 A_{Qg}^{(1)} &= -\frac{1}{2} \widehat{P}_{qg}^{(0)} \ln \left(\frac{m^2}{\mu^2} \right) \\
 A_{Qg}^{(2)} &= \frac{1}{8} \left\{ \widehat{P}_{qg}^{(0)} \otimes \left[P_{qq}^{(0)} - P_{gg}^{(0)} + 2\beta_0 \right] \right\} \ln^2 \left(\frac{m^2}{\mu^2} \right) - \frac{1}{2} \widehat{P}_{qg}^{(1)} \ln \left(\frac{m^2}{\mu^2} \right) \\
 &\quad + \bar{a}_{Qg}^{(1)} \left[P_{qq}^{(0)} - P_{gg}^{(0)} + 2\beta_0 \right] + a_{Qg}^{(2)} \\
 A_{Qq}^{\text{PS},(2)} &= -\frac{1}{8} \widehat{P}_{qg}^{(0)} \otimes P_{gq}^{(0)} \ln^2 \left(\frac{m^2}{\mu^2} \right) - \frac{1}{2} \widehat{P}_{qq}^{\text{PS},(1)} \ln \left(\frac{m^2}{\mu^2} \right) + a_{Qq}^{\text{PS},(2)} + \bar{a}_{Qg}^{(1)} \otimes P_{gq}^{(0)} \\
 A_{qq,Q}^{\text{NS},(2)} &= -\frac{\beta_{0,Q}}{4} P_{qq}^{(0)} \ln^2 \left(\frac{m^2}{\mu^2} \right) - \frac{1}{2} \widehat{P}_{qq}^{\text{NS},(1)} \ln \left(\frac{m^2}{\mu^2} \right) + a_{qq,Q}^{\text{NS},(2)} + \frac{1}{4} \beta_{0,Q} \zeta_2 P_{qq}^{(0)} .
 \end{aligned}$$


with

$$\widehat{f} = f(N_F + 1) - f(N_F) .$$


Operator insertions in light-cone expansion:



$$\not\Delta \gamma_{\pm} (\Delta \cdot p)^{N-1} ,$$



$$g t_{ji}^a \Delta^{\mu} \not\Delta \gamma_{\pm} \sum_{j=0}^{N-2} (\Delta \cdot p_1)^j (\Delta \cdot p_2)^{N-j-2} ,$$



$$g^2 \Delta^{\mu} \Delta^{\nu} \not\Delta \gamma_{\pm} \sum_{0 \leq j < l}^{N-2} \left[(\Delta p_1)^{N-l-2} (\Delta p_1 + \Delta p_4)^{l-j-1} (\Delta p_2)^j (t^a t^b)_{ji} \right. \\ \left. + (\Delta p_1)^{N-l-2} (\Delta p_1 + \Delta p_3)^{l-j-1} (\Delta p_2)^j (t^b t^a)_{ji} \right] ,$$


$$\gamma_+ = 1 , \quad \gamma_- = \gamma_5 .$$

Δ : light-like momentum, $\Delta^2 = 0$.

γ_5 was treated in the 't Hooft-Veltman-Scheme:

$$\not\Delta \gamma_5 = \frac{i}{6} \varepsilon_{\mu\nu\rho\sigma} \Delta^{\mu} \gamma^{\nu} \gamma^{\rho} \gamma^{\sigma} .$$

Diagrams contributing
to the gluonic OME
 $\hat{A}_{Qg}^{(2)}$:



Calculation in Mellin-space: for **space-like** Q^2 : $0 \leq x \leq 1$:

$$\Rightarrow F(N) = \mathbf{M}[f, N] = \int_0^1 x^{N-1} f(x) dx$$

Convolution:

$$[f \otimes g](x) = \int_0^1 dx_1 \int_0^1 dx_2 \delta(x - x_1 x_2) f(x_1) g(x_2),$$

\Rightarrow Product:

$$\mathbf{M}[f \otimes g, N] = \mathbf{M}[f, N] \mathbf{M}[g, N] = F(N) G(N).$$

$$F_2^{Q\bar{Q}} = \sum_{k=1}^{n_f} e_k^2 \left[f_{k-\bar{k}}(N, \mu^2) H_{2,q}^{NS} \left(N, \frac{Q^2}{m^2}, \frac{Q^2}{\mu^2} \right) \right] \\ + e_Q^2 \left[\Sigma(N, \mu^2) H_{2,q}^{PS} \left(N, \frac{Q^2}{m^2}, \frac{Q^2}{\mu^2} \right) + G(N, \mu^2) H_{2,q}^S \left(N, \frac{Q^2}{m^2}, \frac{Q^2}{\mu^2} \right) \right]$$

$$f_{k-\bar{k}}(N, \mu^2) = f_k(N, \mu^2) - f_{\bar{k}}(N, \mu^2),$$

light-quark densities:


$$\Sigma(N, \mu^2) = \sum_{k=1}^{n_f} f_{k+\bar{k}}(N, \mu^2).$$

Our calculation:

- use of **Mellin-Barnes integrals**

$$\frac{1}{(A+B)^\nu} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} d\sigma A^\sigma B^{-\nu-\sigma} \frac{\Gamma(-\sigma)\Gamma(\nu+\sigma)}{\Gamma(\nu)}$$

↪ numerical check & some analytic results




- use of **hypergeometric functions** for general analytic results

$${}_P F_Q \left[\begin{matrix} (a_1) \dots (a_P) \\ (b_1) \dots (b_Q) \end{matrix} ; z \right] = \sum_{i=0}^{\infty} \frac{(a_1)_i \dots (a_P)_i}{(b_1)_i \dots (b_Q)_i} \frac{z^i}{\Gamma(i+1)}, \quad (c)_i = \frac{\Gamma(c+i)}{\Gamma(c)}.$$

- Summation of (new) infinite **one-parameter sums** into **harmonic sums**.
- Algebraic and structural simplification of the harmonic sums [J. Blümlein, 2003]

Calculating **scalar** Feynman diagrams by Mellin-Barnes integrals:

[I.Bierenbaum, S. Weinzierl, 2003 (massless case); I. Bierenbaum, J. Blümlein and S. Klein, 2006]




$$\begin{aligned}
 I_{e,\nu_1} &= \frac{(\Delta p)^{N-1}}{(4\pi)^D (2\pi i)^2} \frac{(m^2)^{D-\nu_{12345}} (-1)^{\nu_{12345}+1}}{\Gamma(\nu_2)\Gamma(\nu_3)\Gamma(\nu_5)\Gamma(D-\nu_{235})} \int_{\gamma_1-i\infty}^{\gamma_1+i\infty} d\sigma \int_{\gamma_2-i\infty}^{\gamma_2+i\infty} d\tau \Gamma(-\sigma)\Gamma(\nu_3+\sigma) \\
 &\times \frac{\Gamma(-\sigma+\nu_4+N-1)}{\Gamma(-\sigma+\nu_4)} \Gamma(-\tau)\Gamma(\nu_2+\tau) \frac{\Gamma(\sigma+\tau+\nu_{235}-D/2)\Gamma(\sigma+\tau+\nu_5)}{\Gamma(\sigma+\tau+\nu_{23})} \\
 &\times \Gamma(-\sigma-\tau+D-\nu_{23}-2\nu_5) \frac{\Gamma(-\sigma-\tau+\nu_{14}-D/2)}{\Gamma(-\sigma-\tau+\nu_{14}+N-1)},
 \end{aligned}$$

N	2	3	4	5
$I_{e,1}$	+0.49999	+0.31018	+0.21527	+0.16007
$I_{e,2}$	-0.09028	-0.04398	-0.02519	-0.01596

[package MB, M. Czakon, 2006]

Hypergeometric functions: Example, scalar Diagram e:



$$I_{e,1} := \iint \frac{d^D q d^D k}{(2\pi)^{2D}} \frac{(\Delta q)^{N-1}}{[q^2 - m^2]^a [(q-p)^2 - m^2] [k^2 - m^2] [(k-p)^2 - m^2] [(k-q)^2]}$$

- introduce Feynman parameters
- do momentum integrations

$$I_{e,1} := \frac{(\Delta p)^{N-1} \Gamma(1-\varepsilon)}{N(N+1)(4\pi)^{4+\varepsilon} (m^2)^{1-\varepsilon}} \int_0^1 dz dw \frac{w^{-1-\varepsilon/2} (1-z)^{\varepsilon/2} z^{-\varepsilon/2}}{(z+w-wz)^{1-\varepsilon}} \left[1 - w^{N+1} - (1-w)^{N+1} \right],$$

using $\Delta^2 = 0$.

$${}_2F_1 \left[\begin{matrix} a, b+1 \\ c+b+2 \end{matrix} ; z \right] = \frac{\Gamma(c+b+2)}{\Gamma(c+1)\Gamma(b+1)} \int_0^1 dx x^b (1-x)^c (1-zx)^{-a},$$

$$I_{e,1} = \frac{S_\varepsilon^2}{(4\pi)^4 (m^2)^{1-\varepsilon}} \frac{(\Delta p)^{N-1}}{N(N+1)} \exp\left\{ \sum_{i=2}^{\infty} \zeta_i \frac{\varepsilon^i}{i} \right\} \left\{ \begin{aligned} & B(\varepsilon/2 + 1, 1 - \varepsilon/2) B(1, -\varepsilon/2) {}_3F_2 \left[\begin{matrix} 1 - \varepsilon, 1, 1 + \varepsilon/2 \\ 2, 1 - \varepsilon/2 \end{matrix} ; 1 \right] \\ & - B(\varepsilon/2 + 1, 1 - \varepsilon/2) B(1, N + 1 - \varepsilon/2) {}_3F_2 \left[\begin{matrix} 1 - \varepsilon, 1, 1 + \varepsilon/2 \\ 2, N + 2 - \varepsilon/2 \end{matrix} ; 1 \right] \\ & - B(\varepsilon/2 + 1, 1 - \varepsilon/2) B(N + 2, -\varepsilon/2) {}_3F_2 \left[\begin{matrix} 1 - \varepsilon, N + 2, 1 + \varepsilon/2 \\ 2, N + 2 - \varepsilon/2 \end{matrix} ; 1 \right] \end{aligned} \right\}$$

with [Beta-function](#):

$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}, \quad \Gamma(1 - \varepsilon) = \exp(\varepsilon\gamma_E) \exp\left\{ \sum_{i=2}^{\infty} \zeta_i \frac{\varepsilon^i}{i} \right\}, \quad |\varepsilon| < 1.$$

$$\Psi(x) = \frac{1}{\Gamma(x)} \frac{d}{dx} \Gamma(x) \quad \Psi(N+1) = S_1(N) - \gamma.$$

[harmonic sums](#): [[J. Blümlein and S. Kurth, 1999](#); [J. Vermaseren, 1999](#)]

$$S_{a_1, \dots, a_m}(N) = \sum_{n_1=1}^N \sum_{n_2=1}^{n_1} \cdots \sum_{n_m=1}^{n_{m-1}} \frac{(\text{sign}(a_1))^{n_1}}{n_1^{|a_1|}} \frac{(\text{sign}(a_2))^{n_2}}{n_2^{|a_2|}} \cdots \frac{(\text{sign}(a_m))^{n_m}}{n_m^{|a_m|}}$$

$$N \in \mathbb{N}, \forall \ell, a_\ell \in \mathbb{Z} \setminus \{0\}$$

$$\begin{aligned}
 I_{e,1} &= \frac{-S_\varepsilon^2}{(4\pi)^4(m^2)^{1-\varepsilon}} \frac{(\Delta p)^{N-1}}{N(N+1)} \exp \left\{ \sum_{i=2}^{\infty} \zeta_i \frac{\varepsilon^i}{i} \right\} \\
 &\quad \times \sum_{s=0}^{\infty} \left\{ \frac{S_1(s) - S_1(1+N+s)}{(1+s)} + \frac{B(N+1, s+1)}{(1+s)} \right\} + O(\varepsilon) \\
 &= \frac{-S_\varepsilon^2}{(4\pi)^4(m^2)^{1-\varepsilon}} \frac{(\Delta p)^{N-1}}{N(N+1)} \sum_{s=1}^{\infty} \left\{ -\frac{1}{s^2} + \frac{S_1(s)}{s} - \frac{S_1(N+s)}{s} + \frac{B(N+1, s)}{s} \right\} + O(\varepsilon)
 \end{aligned}$$

$$I_{e,1} = \frac{S_\varepsilon^2}{(4\pi)^4(m^2)^{1-\varepsilon}} (\Delta p)^{N-1} \left\{ \frac{S_1^2(N) + 3S_2(N)}{2N(N+1)} + \frac{S_1^3(N) + 3S_1(N)S_2(N) + 8S_3(N)}{12N(N+1)} \varepsilon \right\}$$

More complicated sums \rightarrow solved both with combinations out of analytic and algebraic methods and also with [package SIGMA \[C. Schneider, 2007\]](#),

[\[I. Bierenbaum, J. Blümlein, S. Klein, C. Schneider, arXiv:0707.4659 \[math-ph\]\]](#).

Unpolarized case, examples for individual diagrams – numeric:

Diagram	N	$1/\varepsilon^2$	$1/\varepsilon$	1	ε	ε^2
b	2	-8	4.66666	-8.82690	2.47728	-5.69523
	6	-7.73333	0.81936	-8.89777	-1.84111	-7.25674
c	2	-8	39.6	-7.23431	34.66217	6.52891
	6	-2.66666	16.53968	-2.68048	14.25224	2.77564
d	2	-8	7.86666	-6.34542	4.71236	-2.18586
	6	-2.66666	-0.69523	-2.60657	-1.74990	-2.37611
e	2	8.88889	-11.2593	9.82824	-12.8921	2.39145
	6	2.93878	-4.24257	3.39094	-4.3892	0.826978
f	2	5.33333	-9.77777	18.34139	-2.52360	16.20210
	6	3.31428	-6.87289	12.25672	-1.63790	10.86956
g	2	2.66666	-9.55555	4.59662	-8.92015	1.07313
	6	0.57142	-2.00204	1.04814	-1.89142	0.32219

Polarized:
Individual diagrams
– numeric:

Diagram	moment	$1/\varepsilon^2$	$1/\varepsilon$	1	ε	ε^2
a	N = 3	-0.44444	0.12962	-0.26687	-0.30734	-0.12416
	N = 7	-0.06122	0.00819	-0.03339	-0.03800	-0.01278
b	N = 3	4.44444	-1.07407	4.45579	0.515535	3.13754
	N = 7	5.46122	0.74491	6.09646	2.97092	5.35587
c	N = 3	2.66666	-16.28888	0.26606	-13.11030	-5.29203
	N = 7	1.71428	-10.24659	0.28684	-8.21536	-3.19052
d	N = 3	2.66666	-0.02222	2.19940	1.03927	1.69331
	N = 7	1.71428	0.85340	1.78773	1.56227	1.80130
e	N = 3	-2.66666	4.99999	-2.27718	4.89956	0.73208
	N = 7	-1.71428	2.97857	-1.34709	2.83548	0.44608
f	N = 3	0	1.55555	-11.60184	-5.27120	-13.14668
	N = 7	0	2.80210	-7.08455	-1.57130	-7.44933
l	N = 3	-9.33333	0.25000	-8.83933	-3.25228	-6.84460
	N = 7	-6.73877	-1.86855	-7.09938	-4.56050	-6.50099
m	N = 3	-0.44444	1.42592	-0.82397	1.39877	-0.23237
	N = 7	-0.06122	0.22649	-0.11722	0.23939	-0.02415
n	N = 3	-2.22222	1.26851	-1.37562	0.69748	-0.36030
	N = 7	-3.19183	-0.50674	-3.39831	-1.76669	-2.97338

Results to order $O(1)$: [I. Bierenbaum, J. Blümlein, S. Klein, 2006 & 2007]

$$\begin{aligned}
A_e^{Qg} = T_R \left[C_F - \frac{C_A}{2} \right] & \left\{ \frac{1}{\varepsilon^2} \frac{16(N+3)}{(N+1)^2} + \frac{1}{\varepsilon} \left[-\frac{8(N+2)}{N(N+1)} S_1(N) - 8 \frac{3N^3 + 9N^2 + 12N + 4}{N(N+1)^3(N+2)} \right] \right. \\
& + \left[-2 \frac{9N^4 + 40N^3 + 71N^2 - 12N - 36}{N(N+1)^2(N+2)(N+3)} S_2(N) - 2 \frac{N^3 - N^2 - 8N - 36}{N(N+1)(N+2)(N+3)} S_1^2(N) + 4 \frac{(N+3)}{(N+1)^2} \zeta_2 \right. \\
& + \left. 4 \frac{4N^5 + 19N^4 + 31N^3 - 30N^2 - 44N - 24}{N^2(N+1)^2(N+2)(N+3)} S_1(N) + \frac{4P_4(N)}{N^2(N+1)^4(N+2)^2(N+3)} \right] \\
& + \varepsilon \left[-2 \frac{N+2}{N(N+1)} \left(2S_{2,1}(N) + S_1(N)\zeta_2 \right) - \frac{2}{3} \frac{13N^4 + 60N^3 + 111N^2 + 4N - 36}{N(N+1)^2(N+2)(N+3)} S_3(N) \right. \\
& - \frac{1}{3} \frac{N^3 - N^2 - 8N - 36}{N(N+1)(N+2)(N+3)} \left(3S_2(N)S_1(N) + S_1^3(N) \right) - 2 \frac{3N^3 + 9N^2 + 12N + 4}{N(N+1)^3(N+2)} \zeta_2 \\
& + \frac{P_{e1}}{N^2(N+1)^3(N+2)(N+3)} S_2(N) + \frac{4N^5 + 11N^4 + 15N^3 - 86N^2 - 92N - 24}{N^2(N+1)^2(N+2)(N+3)} S_1^2(N) \\
& \left. - 2 \frac{P_{e2}}{N^2(N+1)^3(N+2)^2(N+3)} S_1(N) - 2 \frac{P_{e3}}{N^3(N+1)^5(N+2)^3(N+3)} + \frac{4}{3} \frac{N+3}{(N+1)^2} \zeta_3 \right] \left. \right\}
\end{aligned}$$

Unpolarized case, Singlet O(1)

$$\begin{aligned}
a_{Qg}^{(2)}(N) = & 4C_F T_R \left\{ \frac{N^2 + N + 2}{N(N+1)(N+2)} \left[-\frac{1}{3} S_1^3(N-1) + \frac{4}{3} S_3(N-1) - S_1(N-1) S_2(N-1) \right. \right. \\
& \left. \left. - 2\zeta_2 S_1(N-1) \right] + \frac{N^4 + 16N^3 + 15N^2 - 8N - 4}{N^2(N+1)^2(N+2)} S_2(N-1) + \frac{3N^4 + 2N^3 + 3N^2 - 4N - 4}{2N^2(N+1)^2(N+2)} \zeta_2 \right. \\
& \left. + \frac{2}{N(N+1)} S_1^2(N-1) + \frac{N^4 - N^3 - 16N^2 + 2N + 4}{N^2(N+1)^2(N+2)} S_1(N-1) + \frac{P_1(N)}{2N^4(N+1)^4(N+2)} \right\} \\
& + 4C_A T_R \left\{ \frac{N^2 + N + 2}{N(N+1)(N+2)} \left[4\mathbf{M} \left[\frac{\text{Li}_2(x)}{1+x} \right] (N+1) + \frac{1}{3} S_1^3(N) + 3S_2(N) S_1(N) \right. \right. \\
& \left. \left. + \frac{8}{3} S_3(N) + \beta''(N+1) - 4\beta'(N+1) S_1(N) - 4\beta(N+1) \zeta_2 + \zeta_3 \right] - \frac{N^3 + 8N^2 + 11N + 2}{N(N+1)^2(N+2)^2} S_1^2(N) \right. \\
& \left. - 2 \frac{N^4 - 2N^3 + 5N^2 + 2N + 2}{(N-1)N^2(N+1)^2(N+2)} \zeta_2 - \frac{7N^5 + 21N^4 + 13N^3 + 21N^2 + 18N + 16}{(N-1)N^2(N+1)^2(N+2)^2} S_2(N) \right. \\
& \left. - \frac{N^6 + 8N^5 + 23N^4 + 54N^3 + 94N^2 + 72N + 8}{N(N+1)^3(N+2)^3} S_1(N) - 4 \frac{N^2 - N - 4}{(N+1)^2(N+2)^2} \beta'(N+1) \right. \\
& \left. + \frac{P_2(N)}{(N-1)N^4(N+1)^4(N+2)^4} \right\} .
\end{aligned}$$

Unpolarized case, Singlet $O(\varepsilon)$

$$\begin{aligned}
\bar{a}_{Qg}^{(2)} = & T_F C_F \left\{ \frac{2}{3} \frac{(N^2 + N + 2)(3N^2 + 3N + 2)}{N^2(N+1)^2(N+2)} \zeta_3 + \frac{P_1}{N^3(N+1)^3(N+2)} S_2 + \frac{N^4 - 5N^3 - 32N^2 - 18N - 4}{N^2(N+1)^2(N+2)} S_1^2 \right. \\
& + \frac{N^2 + N + 2}{N(N+1)(N+2)} \left(16S_{2,1,1} - 8S_{3,1} - 8S_{2,1}S_1 + 3S_4 - \frac{4}{3}S_3S_1 - \frac{1}{2}S_2^2 - S_2S_1^2 - \frac{1}{6}S_1^4 + 2\zeta_2S_2 - 2\zeta_2S_1^2 - \frac{8}{3}\zeta_3S_1 \right) \\
& - 8 \frac{N^2 - 3N - 2}{N^2(N+1)(N+2)} S_{2,1} + \frac{2}{3} \frac{3N+2}{N^2(N+2)} S_1^3 + \frac{2}{3} \frac{3N^4 + 48N^3 + 43N^2 - 22N - 8}{N^2(N+1)^2(N+2)} S_3 + 2 \frac{3N+2}{N^2(N+2)} S_2S_1 + 4 \frac{S_1}{N^2} \zeta_2 \\
& + \left. \frac{N^5 + N^4 - 8N^3 - 5N^2 - 3N - 2}{N^3(N+1)^3} \zeta_2 - 2 \frac{2N^5 - 2N^4 - 11N^3 - 19N^2 - 44N - 12}{N^2(N+1)^3(N+2)} S_1 + \frac{P_2}{N^5(N+1)^5(N+2)} \right\} \\
& + T_F C_A \left\{ \frac{N^2 + N + 2}{N(N+1)(N+2)} \left(16S_{-2,1,1} - 4S_{2,1,1} - 8S_{-3,1} - 8S_{-2,2} - 4S_{3,1} - \frac{2}{3}\beta'''' + 9S_4 - 16S_{-2,1}S_1 \right. \right. \\
& + \frac{40}{3}S_1S_3 + 4\beta''S_1 - 8\beta'S_2 + \frac{1}{2}S_2^2 - 8\beta'S_1^2 + 5S_1^2S_2 + \frac{1}{6}S_1^4 - \frac{10}{3}S_1\zeta_3 - 2S_2\zeta_2 - 2S_1^2\zeta_2 - 4\beta'\zeta_2 - \frac{17}{5}\zeta_2^2 \left. \right) \\
& + \frac{4(N^2 - N - 4)}{(N+1)^2(N+2)^2} \left(-4S_{-2,1} + \beta'' - 4\beta'S_1 \right) - \frac{2}{3} \frac{N^3 + 8N^2 + 11N + 2}{N(N+1)^2(N+2)^2} S_1^3 + 8 \frac{N^4 + 2N^3 + 7N^2 + 22N + 20}{(N+1)^3(N+2)^3} \beta' \\
& + 2 \frac{3N^3 - 12N^2 - 27N - 2}{N(N+1)^2(N+2)^2} S_2S_1 - \frac{16}{3} \frac{N^5 + 10N^4 + 9N^3 + 3N^2 + 7N + 6}{(N-1)N^2(N+1)^2(N+2)^2} S_3 - 8 \frac{N^2 + N - 1}{(N+1)^2(N+2)^2} \zeta_2S_1 \\
& - \frac{2}{3} \frac{9N^5 - 10N^4 - 11N^3 + 68N^2 + 24N + 16}{(N-1)N^2(N+1)^2(N+2)^2} \zeta_3 - \frac{P_3}{(N-1)N^3(N+1)^3(N+2)^3} S_2 - \frac{2P_4}{(N-1)N^3(N+1)^3(N+2)^2} \zeta_2 \\
& - \left. \frac{P_5}{N(N+1)^3(N+2)^3} S_1^2 + \frac{2P_6}{N(N+1)^4(N+2)^4} S_1 - \frac{2P_7}{(N-1)N^5(N+1)^5(N+2)^5} \right\}.
\end{aligned}$$

Unpolarized case, pure-singlet and non-singlet

$$a_{Qq}^{\text{PS},(2)} = C_F T_R \left\{ \left[-4 \frac{(N^2 + N + 2)^2}{(N-1)N^2(N+1)^2(N+2)} \left(2S_2(N) + \zeta_2 \right) + \frac{4P_5(N)}{(N-1)N^4(N+1)^4(N+2)^3} \right] \right. \\ \left. + \varepsilon \left[-2 \frac{(5N^3 + 7N^2 + 4N + 4)(N^2 + 5N + 2)}{(N-1)N^3(N+1)^3(N+2)^2} \left(2S_2(N) + \zeta_2 \right) \right. \right. \\ \left. \left. - \frac{4}{3} \frac{(N^2 + N + 2)^2}{(N-1)N^2(N+1)^2(N+2)} \left(3S_3(N) + \zeta_3 \right) + 2 \frac{P_9}{(N-1)N^5(N+1)^5(N+2)^4} \right] \right\}.$$

$$a_{qq,Q}^{\text{NS},(2)} = C_F T_R \left\{ \left[-\frac{8}{3} S_3(N) - \frac{8}{3} \zeta_2 S_1(N) + \frac{40}{9} S_2(N) + 2 \frac{3N^2 + 3N + 2}{3N(N+1)} \zeta_2 - \frac{224}{27} S_1(N) \right. \right. \\ \left. \left. + \frac{219N^6 + 657N^5 + 1193N^4 + 763N^3 - 40N^2 - 48N + 72}{54N^3(N+1)^3} \right] \right. \\ \left. + \varepsilon \left[\frac{4}{3} S_4(N) + \frac{4}{3} S_2(N) \zeta_2 - \frac{8}{9} S_1(N) \zeta_3 - \frac{20}{9} S_3(N) - \frac{20}{9} S_1(N) \zeta_2 + 2 \frac{3N^2 + 3N + 2}{9N(N+1)} \zeta_3 + \frac{112}{27} S_2(N) \right. \right. \\ \left. \left. + \frac{3N^4 + 6N^3 + 47N^2 + 20N - 12}{18N^2(N+1)^2} \zeta_2 - \frac{656}{81} S_1(N) + \frac{P_8}{648N^4(N+1)^4} \right] \right\}.$$


Polarized case, Singlet

$$\begin{aligned}
a_{Qg}^{(2)} = & C_F T_R \left\{ 4 \frac{N-1}{3N(N+1)} \left(-4S_3(N) + S_1^3(N) + 3S_1(N)S_2(N) + 6S_1(N)\zeta_2 \right) \right. \\
& - 4 \frac{N^4 + 17N^3 + 43N^2 + 33N + 2}{N^2(N+1)^2(N+2)} S_2(N) - 4 \frac{3N^2 + 3N - 2}{N^2(N+1)(N+2)} S_1^2(N) \\
& \left. - 2 \frac{(N-1)(3N^2 + 3N + 2)}{N^2(N+1)^2} \zeta_2 - 4 \frac{N^3 - 2N^2 - 22N - 36}{N^2(N+1)(N+2)} S_1(N) - \frac{2P_3(N)}{N^4(N+1)^4(N+2)} \right\} \\
& + C_A T_R \left\{ 4 \frac{N-1}{3N(N+1)} \left(12\mathbf{M} \left[\frac{\text{Li}_2(x)}{1+x} \right] (N+1) + 3\beta''(N+1) - 8S_3(N) - S_1^3(N) \right. \right. \\
& - 9S_1(N)S_2(N) - 12S_1(N)\beta'(N+1) - 12\beta(N+1)\zeta_2 - 3\zeta_3 \left. \right) - 16 \frac{N-1}{N(N+1)^2} \beta'(N+1) \\
& + 4 \frac{N^2 + 4N + 5}{N(N+1)^2(N+2)} S_1^2(N) + 4 \frac{7N^3 + 24N^2 + 15N - 16}{N^2(N+1)^2(N+2)} S_2(N) + 8 \frac{(N-1)(N+2)}{N^2(N+1)^2} \zeta_2 \\
& \left. + 4 \frac{N^4 + 4N^3 - N^2 - 10N + 2}{N(N+1)^3(N+2)} S_1(N) - \frac{4P_4(N)}{N^4(N+1)^4(N+2)} \right\}.
\end{aligned}$$

[J. Blümlein and S. Klein, 2007]

Heavy Flavor Wilson Coefficient for experimental use :

Inversion from Mellin-space to z-space: [J. Blümlein, ANCONT, 2000]



Continuation of harmonic sums:

$$S_1(N) = \Psi(N + 1) + \gamma,$$

etc.

$$F_2^{Q\bar{Q}}(x, Q^2) = \int_0^\infty dz \text{Im} [e^{i\Phi} x^{-c(z)} F_2^{Q\bar{Q}}(c(z), Q^2)],$$

$$c(z) = c_0 + ze^{i\Phi}$$

First Calculation to $O(\alpha_S^2)$: [Buza, Matiounine, Smith, Migneron, van Neerven, 1996]

↪ **Integration-by-parts method**

↪ direct integration of individual Feynman-parameter integrals in z-space

⇒ combinations of **Nielsen integrals**:
$$S_{p,n}(x) = \frac{(-1)^{n+p-1}}{(n-1)!p!} \int_0^1 \frac{dz}{z} \ln^{n-1}(z) \ln^p(1-zx)$$

$\delta(1-x)$	1	$\ln(x)$	$\ln^2(x)$	$\ln^3(x)$	$\ln(1-x)$
$\ln^2(1-x)$	$\ln^3(1-x)$	$\ln(x)\ln(1-x)$	$\ln(x)\ln^2(1-x)$	$\ln^2(x)\ln(1-x)$	$\ln(1+x)$
$\ln(x)\ln(1+x)$	$\ln^2(x)\ln(1+x)$	$\text{Li}_2(1-x)$	$\ln(x)\text{Li}_2(1-x)$	$\ln(1-x)\text{Li}_2(1-x)$	$\text{Li}_3(1-x)$
$S_{1,2}(1-x)$	$S_{1,2}(-x)$	$\frac{1}{1-x}$	$\frac{1}{1+x}$	$\frac{\ln(x)}{1-x}$	$\frac{\ln^2(x)}{1-x}$
$\frac{\ln^3(x)}{1-x}$	$\frac{\ln(x)}{1+x}$	$\frac{\ln^2(x)}{1+x}$	$\frac{\ln^3(x)}{1+x}$	$\frac{\ln(1+x)}{1+x}$	$\frac{\ln(x)\ln(1+x)}{1+x}$
$\frac{\ln(x)\ln^2(1+x)}{1+x}$	$\frac{\ln^2(x)\ln(1+x)}{1+x}$	$\frac{\ln(x)\ln(1-x)}{1-x}$	$\frac{\ln(x)\ln^2(1-x)}{1-x}$	$\frac{\ln(1-x)\text{Li}_2(x)}{1-x}$	$\frac{\text{Li}_2(1-x)}{1-x}$
$\frac{\ln(x)\text{Li}_2(1-x)}{1-x}$	$\frac{\ln(x)\text{Li}_2(1-x)}{1+x}$	$\frac{\ln(1+x)\text{Li}_2(-x)}{1+x}$	$\ln(1+x)\text{Li}_2(-x)$	$\text{Li}_2(-x)$	$\frac{\text{Li}_2(-x)}{1+x}$
$\frac{\ln(x)\text{Li}_2(-x)}{1+x}$	$\frac{\text{Li}_3(1-x)}{1-x}$	$\frac{\text{Li}_3(-x)}{1+x}$	$\frac{S_{1,2}(1-x)}{1-x}$	$\frac{S_{1,2}(1-x)}{1+x}$	$\frac{S_{1,2}(-x)}{1+x}$

Complexity of the results in Mellin space, unpolarized case to order $O(\varepsilon)$:

Diag	S_1	S_2	S_3	S_4	S_{-2}	S_{-3}	S_{-4}	$S_{2,1}$	$S_{-2,1}$	$S_{-2,2}$	$S_{3,1}$	$S_{-3,1}$	$S_{2,1,1}$	$S_{-2,1,1}$
a		++	+											
b	++	++	++	+				++			+		+	
c		++	+											
d	++	++	+					+						
e	++	++	+					+						
f	++	++	++	+				++					+	
g	++	++	+					+						
h	++	++	+					+						
i	++	++	++	+	++	++	+	++	++	+	+	+	+	+
j		++	+											
k		++	+											
l	++	++	++	+				++			+		+	
m		++	+											
n	++	++	++	+	++	++	+	++	++	+	+	+	+	+
o	++	++	++	+				++			+		+	
p	++	++	++	+				++			+		+	
s		++	+											
t		++	+											
PS _a		++	+											
PS _b		++	+											
NS _a														
NS _b	++	++	++	+										
Σ	++	++	++	+	++	++	+	+	++	+	+	+	+	+

van Neerven et al. to $O(1)$: unpolarized: 48 basic functions; polarized: 24 basic functions.

$O(1)$: $\{S_1, S_2, S_3, S_{-2}, S_{-3}\}$, $S_{-2,1} \implies 2$ basic objects.

$O(\varepsilon)$: $\{S_1, S_2, S_3, S_4, S_{-2}, S_{-3}, S_{-4}\}$, $S_{2,1}$, $S_{-2,1}$, $S_{-3,1}$, $S_{2,1,1}$, $S_{-2,1,1}$
 $S_{-2,2}$ depends on $S_{-2,1}$, $S_{-3,1}$
 $S_{3,1}$ depends on $S_{2,1}$
 $\implies 6$ basic objects

These objects are in common to all single scale higher order processes.

Str. Functions, DIS HQ, Fragn. Functions, DY, Hadr. Higgs-Prod., s+v contr. to Bhabha scatt., ...

- Structure of expression is given by

$$\beta(N+1) = (-1)^N [S_{-1}(N) + \ln(2)] ,$$

$$\beta^{(k)}(N+1) = \Gamma(k+1)(-1)^{N+k} [S_{-k-1}(N) + (1 - 2^{-k})\zeta_{k+1}] , \quad k \geq 2 ,$$

$$\mathbf{M} \left[\frac{\text{Li}_2(x)}{1+x} \right] (N+1) - \zeta_2 \beta(N+1) = (-1)^{N+1} [S_{-2,1}(N) + \frac{5}{8}\zeta_3]$$

- \implies harmonic sums with index $\{-1\}$ cancel (holds even for each diagram)

[cf. J.B., 2004; J.B. and V. Ravindran, 2005,2006; J.B. and S. Klein, arXiv: 0706.2426 [hep-ph],

J.B. and S. Moch in preparation.]

Calculation of quark-mass effects in QCD Wilson-coefficients in asymptotic regime $Q^2 \gg m^2$

- Calculation in **Mellin space**, **no use** of the IBP-Method
→ essential for simplification of calculation
- Use of **Mellin-Barnes integrals** (mainly numerical checks) and **generalized hypergeometric functions**, **new summation techniques**
- Results in term of **nested harmonic sums**
→ use of algebraic relations of harmonic sums for simplification of results
→ up to $O(\varepsilon)$ the usual **six basic harmonic sums** contribute
- Calculation of the constant term of the Operator Matrix Elements
→ **full agreement** with results of van Neerven et al. (**in a certain scheme**).
- **New:** Calculation of the **$O(\varepsilon)$ term** of the two-loop OMEs a_{Qg}, a_{qq} complete, necessary for the calculation of the Heavy Wilson coefficients up to **$O(\alpha_s^3)$**